

Cystoideen Kalke / Cystoid Limestone

FRITZ EBNER

Validity: Invalid; not formalized working term. First detailed description by FLAJS & SCHÖNLAUB (1976).

Type area: ÖK50-UTM, map sheet 4215 Eisenerz (ÖK50-BMN, map sheet 101 Eisenerz).

Type section: Ca. 30 m (altitude 1,575 m) above the bend (N 47°31'44" / E 14°58'03") of the Knappensteig (trail from Präbichl Pass to Leobner Hütte) at the ridge separating the Polsterkar and the Polster S-slope (ÖK50-UTM, map sheet 4215 Eisenerz, ÖK50-BMN, map sheet 101 Eisenerz).

Reference section(s): -

Derivation of name: Due to the occurrence of debris and some theca of cystoidea (FLAJS & SCHÖNLAUB, 1976).

Synonyms: -

Lithology: Light grey to pinkish spotted, indistinctly bedded, pure sparry limestones with weak flaser texture followed by 5 m calcareous sandstones (FLAJS & SCHÖNLAUB, 1976).

Fossils: Recrystallized cystoids (debris and theca), rich conodont fauna (FLAJS & SCHÖNLAUB, 1976).

Origin, facies: Marine shallow water environment.

Chronostratigraphic age: Upper Ordovician (upper Katian–Hirnantian) (FLAJS & SCHÖNLAUB, 1976).

Biostratigraphy: *amorphognathus ordovicicus* conodont zone (FLAJS & SCHÖNLAUB, 1976).

Thickness: 13 m.

Lithostratigraphically higher rank unit: -

Lithostratigraphic subdivision: -

Underlying unit(s): Polster Quartzite.

Overlying unit(s): "Red Sparitic Limestone" (FLAJS & SCHÖNLAUB, 1976: Fig. 3); ? transgressional contact.

Lateral unit(s): "Übergangsporphyroid" at Erzberg (KERN, 1927; SCHÖNLAUB, 1982a; not indicated in Text-Fig. 2): alternation of 6–8 m thick dm-bedded light limestones with greenish grey sandy layers followed by ankeritic material.

Geographic distribution: E-GWZ; Styria, Eisenerzer Alpen.

Remarks: -

Complementary references: TOLLMANN (1977), SCHÖNLAUB (1979, 1980a, 1982a), EBNER et al. (1989), SCHÖNLAUB & HEINISCH (1993).

Vulkanite / Volcanics

FRITZ EBNER

Validity: Invalid; not formalized working term.

Type area: ÖK50-UTM, map sheet 4215 Eisenerz (ÖK50-BMN, map sheets 131 Kalwang and 132 Trofaiach).

Type section: Not defined. Detailed descriptions derive from the mountain Kragelschinken (1,845 m, N 47°29'21" / E 14°49'26") and along the section in the upper Lange Teichen valley (starting NNE of the foot of Mt. Wildfeld, N 47°28'18" / E 14°48'29") (HIESSLEITNER, 1931; SCHÖNLAUB, 1977a, b, 1982a, b).

Reference section(s): -

Derivation of name: According to lithology.

Synonyms: "Kragelschinken Folge" (EBNER et al., 1989); "Basische Vulkanite" in the geological map 1:25,000 (SCHÖNLAUB, 1982a).

Lithology: Alternation of mottled schists ("Fleckenschiefer" = green schists with dark spots of chlorite), diabase schists, sandy-quartzite schists and subordinate layers of massive plagioclase-hornblende rocks (HIESSLEITNER, 1931; DAURER in SCHÖNLAUB, 1982a). Parts of the sequence are characterized as volcanoclastics (lapilli and ash tuffs), basaltic lavas with well preserved pillow structures and concordant swarms of gabbroidic sills (SCHLAEGEL-BLAUT, 1990).

Fossils: Conodonts, crinoids from intercalations of Crinoidal Limestone (SCHÖNLAUB, 1977a, b).

Origin, facies: Two sections (Finzenkogel, Schleichberg) NW Trofaiach (ÖK50-UTM, map sheet 4215 Eisenerz, ÖK50-BMN, map sheet 132 Trofaiach) were characterized by SCHLAEGEL-BLAUT (1990) in terms of volcanic islands (sensu FISHER, 1984). Section Finzenkogel: submarine proximal pyroclastic flow facies at the flank of a volcanic island at the transition of effusive to explosive eruptions below or above PCL (= pressure compensation level at water depths of ~ 500 m; stadium A and B₁ sensu FISHER, 1984). Section Schleichberg: upper part of a pillow volcano below PCL (stadium A sensu FISHER, 1984).

Chronostratigraphic age: Silurian (boundary Llandovery/Wenlock) (SCHÖNLAUB, 1982a).

Biostratigraphy: *amorphognathoides* Zone in crinoidal limestone intercalations within tuffitic volcanics as well as limestone intercalations within black schists/lydites below and above the volcanics (SCHÖNLAUB, 1976, 1977a, 1982a).

Thickness: 250–300 m at Paarenkogel and Kragelschinken (HIESSLEITNER, 1931). Along the Gößgraben (WSW Trofaiach, ÖK50-UTM, map sheet 4215 Eisenerz, ÖK50-BMN, map sheet 132 Trofaiach) the thickness of the entire volcanogenic sequences is between 180 and 550 m. There, the volcanoclastics include also intercalations of schists and up to 150 m thick lavas (SCHLAEGEL-BLAUT, 1990).

Lithostratigraphically higher rank unit: -

Lithostratigraphic subdivision: -

Underlying unit(s): Lydites, Arkoses, Schists.

Overlying unit(s): Crinoidal Limestones (SCHÖNLAUB, 1992) (not shown in ASC 2004).

Lateral unit(s): Lydites, Arkoses, Schists.

Geographic distribution: E-GWZ; Styria, Eisenerzer Alpen.

Remarks: Informal working term for thick basic volcanics (first description HIESSLEITNER, 1931) in the hanging parts of the "Lydites, Arkoses, Schists".

Complementary references: TOLLMANN (1977), SCHÖNLAUB (1979, 1980a), SCHÖNLAUB & HEINISCH (1993), LOESCHKE & HEINISCH (1993).

Crinoidenkalke / Crinoidal Limestone

(not shown in ASC 2004)

FRITZ EBNER

Validity: Invalid; working term (SCHÖNLAUB, 1976, 1979, 1982a).

Austrian Stratigraphic Chart 2004 - Paleozoic

(sedimentary successions)

Austrian Stratigraphic Commission



ERA	SYSTEM / PERIOD / SERIES / EPOCH	STAGE / AGE	DURATION Ma	Global Classification					
				ERATHM / ERA	SYSTEM / PERIOD / SERIES / EPOCH				
PALEOZOIC	PERMIAN	CHANGHSINGIAN / Dorashanian	251	PERMIAN	MID PERMIAN / GUADALUPIAN / LOPINGIAN				
		WUCHIAPINGIAN / Dabuffian	255						
		CAPITANIAN	260						
		WORDIAN	265						
		ROADIAN	270						
		PERMIAN	LOWER PERMIAN / CISURALIAN			KUNGURIAN	275		
						ARTINSKIAN	280		
						SAKMARIAN	285		
						ASSELIAN	290		
		PERMIAN	TRIAS			GZHELIAN	295	TRIAS	U. CARBONIFEROUS / PENNSYLVANIAN
KASIMOVIAN	300								
MOSKOVIAN	305								
BASHKIRIAN	310								
TRIAS	LOWER CARBONIFEROUS / MISSISSIPPIAN			SERPUKHOVIAN	315				
				VISEAN	320				
				TOURNAISIAN	325				
PERMIAN	DEVONIAN			FAMENNIAN	350	DEVONIAN	UPPER DEVONIAN		
				FRASNIAN	355				
				GIVETIAN	360				
		EIFELIAN	365						
		DEVONIAN	LOWER DEVONIAN	EMSIAN	370				
				LOCHKOVIAN	375				
		PERMIAN	DEVONIAN	LUDFORDIAN / GORSTIAN	380			DEVONIAN	MIDDLE DEVONIAN
				HOMERIAN / SHEINWOOD	385				
				TELYCHIAN	390				
				AERONIAN	395				
RHUDDANIAN	400								
DEVONIAN	LOWER DEVONIAN			PRAGIAN	405				
				LOCHKOVIAN	410				
PERMIAN	DEVONIAN			WEN-LOCK / LOW	415	DEVONIAN	LOWER DEVONIAN		
				HIRNANTIAN	420				
				LLANDOVERY	425				
		AERONIAN	430						
		RHUDDANIAN	435						
		DEVONIAN	LOWER DEVONIAN	PRAGIAN	440				
				LOCHKOVIAN	445				
		PERMIAN	DEVONIAN	WEN-LOCK / LOW	450			DEVONIAN	UPPER ORDOVICIAN
				LLANDOVERY	455				
				AERONIAN	460				
RHUDDANIAN	465								
DEVONIAN	LOWER DEVONIAN			PRAGIAN	470				
				LOCHKOVIAN	475				
PERMIAN	DEVONIAN			WEN-LOCK / LOW	480	DEVONIAN	MIDDLE ORDOVICIAN		
				LLANDOVERY	485				
				AERONIAN	490				
				RHUDDANIAN	495				
		DEVONIAN	LOWER DEVONIAN	PRAGIAN	500				
				LOCHKOVIAN	505				
		PERMIAN	DEVONIAN	WEN-LOCK / LOW	510			DEVONIAN	LOWER ORDOVICIAN
				LLANDOVERY	515				
				AERONIAN	520				
				RHUDDANIAN	525				
DEVONIAN	LOWER DEVONIAN			PRAGIAN	530				
				LOCHKOVIAN	535				
PERMIAN	DEVONIAN			WEN-LOCK / LOW	540	DEVONIAN	UPPER CAMBRIAN		
				LLANDOVERY	545				
				AERONIAN	550				
				RHUDDANIAN	555				
		DEVONIAN	LOWER DEVONIAN	PRAGIAN	560				
				LOCHKOVIAN	565				
		PERMIAN	DEVONIAN	WEN-LOCK / LOW	570			DEVONIAN	MIDDLE CAMBRIAN
				LLANDOVERY	575				
				AERONIAN	580				
				RHUDDANIAN	585				
DEVONIAN	LOWER DEVONIAN			PRAGIAN	590				
				LOCHKOVIAN	595				
PERMIAN	DEVONIAN			WEN-LOCK / LOW	600	DEVONIAN	LOWER CAMBRIAN		
				LLANDOVERY	605				
				AERONIAN	610				
				RHUDDANIAN	615				
		DEVONIAN	LOWER DEVONIAN	PRAGIAN	620				
				LOCHKOVIAN	625				



- Legend**
- pelagic, offshore, siliciclastic
 - pelagic, nearshore, calcareous
 - shallow marin, neritic
 - terrestrial-continental, coarse clastic
 - terrestrial-continental, fine clastic
 - evaporite (chloride, sulphate)
 - rhyolite, dacite
 - (basaltic) andesite, trachyandesite
 - basalt
 - phyllite
 - mixed-facies (in corresponding colors)
 - coal (may include several seams)
 - ? position/age doubtful/controversial
 - | equal units
 - \ older unit left \ younger unit right
 - hiatus
 - unconformity
 - GSSP
 - Fm. Formation
 - Ls. Limestone

© Commission for the Palaeontological and Stratigraphical Research of Austria (CPSA) of the Austrian Academy of Sciences and Austrian Stratigraphic Commission

Cutout and English adaptation of the "Die Stratigraphische Tabelle von Österreich 2004": Geological Survey of Austria

The Austrian Stratigraphic Chart 2004 - Paleozoic is a supplement of:
 Hubmann, B., Ebner, F., Ferretti, A., Kido, E., Krainer, K., Neubauer, F., Schönlaub, H.-P. & Suttner, T.J. (2014): The Paleozoic Era (them), 2nd edition. - In: Piller, W.E. (Ed.): The Lithostratigraphic units of the Austrian Stratigraphic Chart 2004 (sedimentary successions) - Vol. 1 - Abhandlungen der Geologischen Bundesanstalt, 66, 9-133, Wien.

Printing: Grasl Druck & Neue Medien GmbH, Bad Vöslau 2014

Landesmuseum Joanneum, OAW, Geologische Bundesanstalt, UNI GRAZ, OGG, Universität Wien, Naturhistorisches Museum Wien