

The Middle Miocene Fish Fauna (excl. otolithes) from Mühlbach am Manhartsberg and Grund near Hollabrunn, Lower Austria

by Ortwin SCHULTZ¹

(With 5 text figures)

Manuscript submitted on 25 October 2002,
the revised manuscript on 16 December 2002

Summary

The sites Mühlbach am Manhartsberg and Grund has yielded relatively little material. The samples from Mühlbach am Manhartsberg yielded 4 Chondrichthyes taxa as well as 6 marine and 1 freshwater-Osteichthyes taxa. 15 Chondrichthyes and 5 Osteichthyes taxa were documented from the Grund site. The marine material from both sites points to neritic conditions. In particular *Acanthurus*, but also *Sphyraena*, *Rhinoptera* as well as *Aetobatus* indicate tropical to subtropical waters. All remaining genera inhabit tropical to temperate waters.

Keywords: Central Paratethys, Gaiendorf Formation, Grund Formation, Middle Miocene, Badenian, fish teeth.

Zusammenfassung

Von den Fundkomplexen Mühlbach am Manhartsberg und Grund liegen jeweils nur relativ kleine Mengen an Belegen vor. Die Proben aus Mühlbach am Manhartsberg erbrachten 4 Chondrichthyes-, 1 Süßwasser- und 6 marine Osteichthyes-Taxa. Aus dem Raum Grund können 15 Chondrichthyes- und 5 Osteichthyes-Taxa nachgewiesen werden. Die marinischen Belege beider Fundgebiete indizieren neritische Verhältnisse. Insbesondere *Acanthurus*, aber auch *Sphyraena*, *Rhinoptera* sowie *Aetobatus*, sind Anzeiger für tropische bis subtropische Gewässer. Alle anderen Gattungen kommen in tropischen bis temperierten Gewässern vor.

Schlüsselwörter: Zentrale Paratethys, Gaiendorf-Formation, Grund-Formation, Badenium, Mittel Miozän, Fischzähne.

Introduction

Two sediment samples of the Gaiendorf Formation at Mühlbach am Manhartsberg, Lower Austria (Fig. 1) yielded fish remains. The individual taxa are primarily documented by isolated teeth or tooth plates and a few tail spines. The otolith faunas are treated in a separate contribution (REICHENBACHER, in prep.).

For comparative purposes, the fish fauna from Grund near Hollabrunn (Fig. 1), which is of the same age, was also analysed. Tab. 1 demonstrates that only small faunas with relatively few specimens are present. Nonetheless, a total of 24 taxa were found in the Lower Lagenidae Zone (Early Middle Miocene) in Lower Austria: 8 sharks/Squalomorphii, 7 rays/Batomorphii, 8 marine bony fishes/Osteichthyes as well as 1 freshwater bony fish/Osteichthyes.

¹ Dr. Ortwin SCHULTZ, Naturhistorisches Museum Wien, Geologisch-Paläontologische Abteilung, Postfach 417, A-1014 Wien, Austria / Österreich. – e-mail: ortwin.schultz@nhm-wien.ac.at




Fig. 1: Local position of the localities Mühlbach am Manhartsberg and Grund within Lower Austria during the Middle Miocene.

The fish faunas were not expected to provide evidence about their stratigraphic position. The study of the foraminifera faunas (RÖGL & SPEZZAFERRI 2003), of the calcareous nanoplankton (CORIC 2003) as well as of the rodent fauna (DAXNER-HÖCK 2003), however, demonstrated their position in Zone M5b/Mt5b or NN5, or in the higher MN5 or the higher Lower Lagenid Zone, Lower Badenian, Middle Miocene. The geological situation of the Gaindorf Formation at Mühlbach am Manhartsberg is documented by ROETZEL 2003.

All the material treated here is stored in the Geolog.-Paläontolog. Abteilung of the Naturhistorische Museum in Vienna: NHMWien: 2002z0067/0001-0010, 2002z0123/0001-0009, 2002z0124/0001-0010, 2002z0125/0001-0003, 2002z0126/0001-0009, 2002z0127/0001-0014, along with old material.

Faunal content

The determination of all marine material presented no greater difficulties². The determination of *Sphyrna*, however, remains uncertain because only a somewhat eroded tooth is available.

² determinations are based largely on the following literature: CAPPETTA 1970 and 1987, PROBST 1877, REINECKE et al. 2001, SCHULTZ 1971, 1978 and 1998

Tab. 1: The fish fauna of the Gaiendorf and the Grund Formation in Lower Austria³

	Mühlbach am Manhartsberg		Grund 1999			Vicinity of Grund
	Mü 1	Mü 2	GRU- B1-1	GRU- B1-3	GRU- F-11	
	NHM Wien		NHM Wien		NHM Wien	
Marine fishes						
<i>Squalomorphii</i> indet. ⁴					2	
<i>Notorynchus primigenius</i>						2
<i>Carcharias acutissimus</i>			1		9	21+3?
<i>Carcharias cuspidatus</i>						9
<i>C. acutissimus</i> or <i>C. cuspidatus</i>					2	
<i>Scyliorhinus distans</i>	1	1			1	
<i>Carcharhinus priscus</i>	1	2		1?	1+2?	11
<i>Galeocerdo aduncus</i>						2
<i>Sphyrna</i> ? sp.				1		
Batomorphii						
<i>Dasyatis</i> cf. <i>rugosa</i>		3		1	1	
<i>Dasyatis</i> sp.		1		1		2
<i>Myliobatis</i> sp.						1
<i>Rhinoptera</i> sp.						1
<i>Myliobatis/Rhinoptera</i> sp.				1	2	5
<i>Aetobatus arcuatus</i>				1		2
<i>Myliobatidae/Dasyatidae</i> ⁹ (Fig. 2)			1		2	
Osteichthyes						
<i>Diplodus incisivus</i>		1		2		
<i>Sparus umbonatus</i>					6	10
<i>Pagrus cinctus</i>					1	1+?1
<i>Sparidae</i> indet. ¹³	4	15		13	4	6
? <i>Labridae</i> indet.		1				
<i>Acanthurus haueri</i> (Fig. 3)	1					
<i>Sphyraena</i> sp. (Fig. 4)	2				1	
<i>Trichiurus miocaenicus</i> (Fig. 5)		1				
Freshwater fishes						
<i>Palaeoleuciscus</i> sp.	18					

³ Arrangement according to ESCHMEYER 1990⁴ vertebra⁵ from Guntersdorf 5 teeth, from Immendorf, Gaiendorf and from Niederleis 1 tooth each^{6, 7} from Windpassing^{8, 11} from Gaiendorf⁹ tail spine¹⁰ from Guntersdorf 1 fragment¹² from Mailberg¹³ with different tooth-types: small, round, dome-shaped tooth plates, or conical teeth, or elongate-conical teeth with curved tips = ? pharyngeal teeth¹⁴ from Guntersdorf




Fig. 2: **Myliobatidae** or **Dasyatidae**, tail spine, Grund (GRU-F-11); Badenian, Lower Lagenid Zone. – NHMWien 2002z0127/0008. – a: upper surface, b: lower surface.

The determination of the freshwater material relied on the excellent overview of morphotypes of the pharyngeal teeth ("Schlundzähne der Cyprinen" by HECKEL 1843: Plate 1), because only few publications have dealt with such fossil cyprinid teeth. In this connection the following authors deserve mention: RUTTE 1962, RUTTE & BECKER-PLATEN 1980, OBRHELOVA 1969, 1971 and 1990, RUTTE & VAN DE WEERD 1980, SYTCHEVSKAYA 1989, GAUDANT 1989-1997 as well as BÖHME 1993.

Four of the 18 isolated teeth from the Cyprinidae pharyngeal apparatus from Mühlbach am Manhartsberg (sample Mü 1), can be determined more precisely. The denticulation or grooves of the masticatory surface (compare RUTTE 1962: 177, Fig. 3) are represented by forms of the type "hooked teeth with masticatory surface" ("Hakenzähne mit Kauflächen"), or the "Drück- und Greifzähne" along with those of the type "hooked teeth without masticatory surface" ("Hakenzähne ohne Kauflächen") or so-called "Fangzähne" (HECKEL 1843): these are *Phoxinellus alepidotus* HECKEL, 1843, "*Leucos Basak*" – now *Rutilus basak* (HECKEL, 1843) –, "*Bliccopsis Buggenhagii*" – now recognized as a hybrid of *Aramis brama* x *Rutilus rutilus* – and *Blicca argyroleuca* – now *Aramis bjoerkna* (LINNAEUS, 1758) – as well as "*Squalius Dobula* BPT." and *Scardinius erythrophthalmus*¹⁵. A more detailed comparison based on the rich collection of pharyngeal apparatuses at the NHMWien¹⁶, which contains the specimens of HECKEL 1843, yielded a good correspondence for the four of the pharyngeal teeth with main row teeth (see RUTTE 1962: 177, Fig. 3) of "*Squalius Dobula*" – now *Leuciscus leuciscus* (LINNAEUS, 1758) –, "*Squ. Berag* HECKEL" – now *Leuciscus*

¹⁵ updated according to ESCHEMAYER 2002

¹⁶ comparisons were done with the following forms: *Aramis ballerus* (L.), *A. brama* (L.), *A. sapo* (PALLAS), *A. vetula* HECKEL, *Alburnoides bipunctatus* (BLOCH), *Alburnus iblis* HECKEL, *Aspius aspius* (L.), *Barbus barbus* (L.), *Blicca bjoerkna* (L.), *Bliccopsis abramorutilus* HECKEL, *Cyprinion macrostomus* HECKEL, *Gobio vulgaris* CUVIER, *Leuciscus cephalus orientalis* NORDM., *L. idus* (L.), *L. meidingeri* HECKEL, *L. pausingeri* HECKEL, *Leucos basak* HECKEL, *Pelecus cultratus* (L.), *Phoxinus laevis* AGASSIZ, *Rhodeus amarus* AGASSIZ, *Rutilus frisii* (NORDM.), *R. rutilus* (L.) x *Aramis brama* (L.), *Scardinius erythrophthalmus* BPT., *S. scardofa* BPT., *Squalius argenteus* SELYS, *Squ. balteatus* HECKEL, *Squ. berag* HECKEL, *Squ. cavedanus* BPT., *Squ. cephalus* L., *Squ. compressus* HECKEL, *Squ. dobula* BPT., *Squ. lepidus* HECKEL, *Squ. microlepis* HECKEL, *Squ. turskyi* HECKEL, *Squ. tyberinus* BPT., *Squ. delineatus* HECKEL, *Telestes savignyi* BPT., *Tinca tinca* (L.), *Vimba vimba* L.

cephalus (LINNAEUS, 1758)–, "Squ. Spurius HECKEL" – now *Leuciscus spurius* (HECKEL, 1843)– and "Squ. Turskyi HECKEL" – now *Leuciscus turskyi* (HECKEL, 1843). If the four available pharyngeal teeth, with their particularly characteristic denticulation, were from Recent material, then they would be determined as *Leuciscus* sp. (Leuciscinae) (compare also OBRHELOVA 1971: 593, Fig. 42/A-L).

Three additional, isolated pharyngeal teeth can be attributed to the pharyngeal apparatus of *Leuciscus* of the taxa just listed: 1st and 2nd tooth of the upper row (compare RUTTE 1962: 177, Fig. 3). The remaining 11 isolated smaller pharyngeal teeth lack distinctive features, but an attribution to *Leuciscus* is likely.

The genus *Leuciscus* is represented in the Miocene by *Palaeoleuciscus* OBRHELOVA, 1969 (compare OBRHELOVA 1969, 1971 and 1990, GAUDANT 1989-1995, as well as BÖHME 1993). A comparison with the pharyngeal teeth of this genus, for example of *Palaeoleuciscus chartaceus* (LAUBE) (compare GAUDANT 1993: Fig. 6 or NHMWien 1971/1405/10), confirms the strong morphological correspondence, so that all the pharyngeal teeth from Mühlbach can be attributed to *Palaeoleuciscus* sp.

Evaluation of the biogeographical and ecological conditions¹⁷

Although only a small number of fish findings are available (see Table 1 and 2) they do provide some general ecological information on the early marine Badenian of the Molasse Zone.

Tab. 2: Biogeographical and ecological data of fishes based on Recent distributions

Genera	tropical / subtropical	temperate	cold	littoral / neritic	pelagic	bathyal	benthonic	Fresh-water
<i>Notorynchus</i>	(+)	+		+			+	
<i>Carcharias</i> (2 taxa)	+	+		+	+			
<i>Scyliorhinus</i>	+	+		+		+	+	
<i>Carcharhinus</i>	+	(+)		+	+			
<i>Galeocerdo</i>	+	+		+	+			
<i>Sphyrna</i>	+	(+)		+				
<i>Dasyatis</i> (2 taxa)	+	+		+			+	
<i>Myliobatis</i>	+	+		+			+	
<i>Aetobatus</i>	+			+			+	
<i>Rhinoptera</i>	+			+			+	
<i>Palaeoleuciscus</i>		+						+
<i>Diplodus</i>	+	+		+				
<i>Sparus</i>	+	+		+				
<i>Pagrus</i>	+	+		+				
<i>Acanthurus</i> (Fig. 3)	+			+				
<i>Sphyraena</i> (Fig. 4)	+	(+)		+	+			
<i>Trichiurus</i> (Fig. 5)	+			+	+	+		

¹⁷ compiled largely from COMPAGNO 1984 & 1999, DECKERT 1974 and SMITH & HEEMSTRA 1986

tropical - subtropical (4): *Aetobatus, Rhinoptera; Acanthurus* (Fig. 3), *Trichiurus* (Fig. 5)

tropical - temperate (14): *Notorynchus, Carcharias* (2 taxa), *Scyliorhinus, Carcharhinus, Galeocerdo, Sphyrna, Dasyatis* (2 taxa), *Myliobatis; Diplodus, Sparus, Pagrus, Sphyraena* (Fig. 4)

littoral - neritic (16): *Notorynchus, Carcharias* (2 taxa), *Scyliorhinus, Carcharhinus, Sphyrna, Dasyatis* (2 taxa), *Myliobatis, Rhinoptera, Aetobatus; Diplodus, Sparus, Pagrus, Acanthurus* (Fig. 3), *Trichiurus* (Fig. 5)

littoral - pelagic (6): *Carcharias* (2 taxa), *Carcharhinus, Galeocerdo; Sphyraena* (Fig. 4), *Trichiurus* (Fig. 5)

littoral - bathyal (2): *Scyliorhinus, Trichiurus* (Fig. 5)

benthonic (7): *Notorynchus, Scyliorhinus, Dasyatis* (2 taxa), *Myliobatis, Rhinoptera, Aetobatus*

Freshwater (1): *Palaeoleuciscus*

Results

The fish fauna of Mühlbach am Manhartsberg consist of 11 taxa, of which 10 are exclusively marine and only one represents a freshwater form. The habitat of the marine taxa can be classified as neritic. The Recent genus *Trichiurus* (Fig. 5) is known from the continental shelf down to 350 m, but also advances into shallow coastal waters and harbors (PARIN 1986: 980; SMITH & HEEMSTRA 1986: 829); it has been documented also from shallow waters of the Badenian, for example from Gainfarn (SCHULTZ 1978: 213). This underlines that definitive bathymetric conclusions cannot be drawn based on a single tooth.

The *Acanthurus* find (Fig. 3) is remarkable. This genus is a superb indicator for tropical to subtropical marine waters. Also *Aetobatus, Rhinoptera* and *Trichiurus* are inhabitants of such waters. The remaining taxa inhabit tropical to temperate waters.

Palaeoleuciscus has been described from lacustrine depositional environments (OBRHELOVA 1971: 553), whereas the Recent genus *Leuciscus* can be found both in rivers and in lakes.

The fish fauna of the Grund Formation in and around Grund must be classified as poor, even though the site has been sampled for over 150 years. The total yield is only 8 shark/Squalomorphii-, 7 ray/Batomorphii- and 5 bony fish/Osteichthyes-taxa. The high contribution of benthonic taxa is conspicuous. These small-sized sharks as well as the rays and the Osteichthyes-taxa indicate a shallow sea for the Grund Formation in the vicinity of Grund. In particular the genus *Notorynchus* – in contrast to *Hexanchus*¹⁸ – is a Recent inhabitant of shallow waters; it is also known from Miocene shallow-water localities of the Paratethys, e.g. from the Eggenburgian (Kühnring), Ottangian (Höch, Rainbach, Prambachkirchen, Plesching) and the Badenian (Wien-Kalksburg, Gaaden, Devinska Nova Ves) (SCHULTZ 1971, SCHULTZ in BRZOBOHATY & SCHULTZ 1971 and 1973).

¹⁸ depending on species, known from shallow water to 1875 m or from 90 m - 600 m depth (compare COMPAGNO 1984: 20 und 21).




Fig. 3: *Acanthurus haueri* (v.MEYER, 1842), Mühlbach am Manhartsberg (Mü 2); Badenian, Lower Lagenid Zone. – NHMWien 2002z0123/0006.

Fig. 4: *Sphyraena* sp., Grund (GRU-F-11); Badenian, Lower Lagenid Zone. – NHMWien 2002z0127/0014.

Fig. 5: *Trichiurus miocaenicus* (DELFORTRIE, 1876), Mühlbach am Manhartsberg (Mü 2); Badenian, Lower Lagenid Zone. – NHMWien 2002z0124/0010.

Acknowledgements

Sampling was conducted in the framework of Project FWF-P15724 (Österr. Fonds zur Förderung der wissenschaftlichen Forschung; Project leader: G. Daxner-Höck).

It is my pleasant duty to thank Dr. Ernst Mikschi from the fish collection of the NHMWien and his colleagues Helmut Wellendorf, Matthias Reithofer and Christian Pollmann for lending a considerable number of dry mounts. I am grateful to Mrs. Alice Schumacher (NHMWien) for taking the photographs.

References

- BÖHME, E. (1993): Eine Untermiozäne Fischfauna (Teleostei, Cyprinidae) vom Dietrichsberg bei Vacha/Rhön. – Freiberger Forschungshefte, (C - Paläont., Stratigr., Fazies) **450**: 116-150, 52 textfigs., 2 tab., 2 pls. – Leipzig and Stuttgart.
- BRZOBOHATY, R. & SCHULTZ, O. (1971): Die Fischfauna der Eggenburger Schichtengruppe. – S. 719-759, 8 pls. – In: STEININGER, F., SENES, J. & Mitarbeiter (1971a): M1 Eggenburgien. Die Eggenburger Schichtengruppe und ihr Stratotypus. – Chronostratigraphie und Neostratotypen, Miozän der zentralen Paratethys, **2**: 827 pp., 37 textfigs., 9 tab., 1 map, 123 pls. - Bratislava (Vydavatel'stvo Slovenskej akad. vied).
- & — (1973): Die Fischfauna der Innviertler Schichtengruppe und der Rzehakia Formation. – pp. 625-693, 5 pls. – In: PAPP, A., RÖGL, F., SENES, J. & Mitarbeiter: M2 Ottnangien. Die Innviertler, Salgótarjaner, Bántapusztaer Schichtengruppe und die Rzehakia Formation. – Chronostratigraphie und Neostratotypen, Miozän der zentralen Paratethys, **3**: 841 pp., 45 textfigs., 5 tab., 82 pls. – Bratislava (Vydavatel'stvo Slovenskej akad. vied).
- CAPETTA, H. (1970): Les sélaciens du Miocène de la région de Montpellier. – Palaeovertebrata, Mém. extraord. 1970: 139 + 19 pp., 22 textfigs., 6 tab., 27 pls. – Montpellier.
- (1987): Chondrichthyes II. Mesozoic and Cenozoic Elasmobranchii. – In: SCHULTZE, H.-P. (Ed.): Handbook of Paleoichthyology, **3** B: 193 pp., 148 textfigs. – Stuttgart and New York (G. Fischer).
- COMPAGNO, L.J.V. (1984): Sharks of the world. – FAO species catalogue, 4/1 (Hexanchiformes to Lamniformes): VIII + 249 pp., pl. figs. - 4/2 (Carcarhiniformes): X + 251-655, pl. figs. – Rome (Food and Agriculture Organization of the United Nations).
- (2002): Checklist of living elasmobranchs. – In: HAMLETT, W.C. (Ed.): Sharks, skates, and rays: the biology of elasmobranch fishes . – 1-X + 1-555. – Baltimore & London (The John Hopkins University Press).
- CORIC, St. (2003): Calcareous Nannofossils biostratigraphy of the Mühlbach beds (Gaindorf Formation, Lower Badenian), Lower Austria. – Ann. Naturhist. Mus. Wien, **104/A**: 15-21. – Wien.

- DAXNER-HÖCK, G. (2003): *Cricetodon meinii* and other rodents from Mühlbach and Grund, Lower Austria (Middle Miocene, late MN5). – Ann. Naturhist. Mus. Wien, **104/A**: 267-291. – Wien.
- DECKERT, K. (1974): Fische. – 252 S., zahlr. textfigs. – In: DECKERT, K., FREYTAG, G.E., GÜNTHER, K., PETERS, G. & STERBA, G.: Fische, Lurche, Kriechtiere., 534 pp., pl. textfigs. – Reinbek b. Hamburg (rororo, Rowohlt).
- ESCHMEYER, W.N. (1990): Catalog of the Genera of Recent Fishes. – 697 pp. – San Francisco (California Academy of Sciences).
- (2002): Catalog of Fishes. – <http://www.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>.
- GAUDANT, J. (1989): Nouvelles observations sur l'ichthyofaune miocène de Steinheim am Albuch (Wurtemberg, Allemagne). – Stuttgarter Beitr. Naturkunde, (B) **151**: 33 pp., 21 textfigs., 4 pls. – Stuttgart.
- (1993): Révision des Cyprinidae (Poissons téléostéens) miocènes de Pochlovice (Bohème) et conséquences taxonomiques. – N. Jb. Geol. Paläont., Abh. **187/3**: 289-298, 6 textfigs. – Stuttgart.
- (1994a): Sur la présence du genre *Palaeoleuciscus* OBRHELOVÁ (Poissons téléostéens, Cyprinidae) dans le Miocène inférieur ou moyen (?) du Dietrichsberg, près de Vacha (Thuringe). – Paläont. Z., **68/1-2**: 223-243, 15 textfigs., 2 tab. – Stuttgart.
- (1994b): L'ichthyofaune du Pannonien supérieur de Götzendorf an der Leitha (Basse Autriche). – Ann. Naturhistor. Mus. Wien, **96/A**: 117-131, 1 textfig., 2 pls. – Wien.
- (1995): Mise au point sur les poissons Cyprinidae miocènes du Habichtswald, près de Cassel (Hesse, Allemagne). – N.Jb. Geol. Paläont., Mh. 1995/6: 363-374, 6 textfigs. – Stuttgart.
- (1997): L'ichthyofaune pliocène de Willershausen am Harz (Basse Saxe, Allemagne) - un réexamen. – Stuttgarter Beitr. Naturkunde, (B) **257**: 51 pp., 24 textfigs., 10 pls. – Stuttgart.
- HECKEL, J.J. (1843): Zahn-System der Cyprinen. – S. 1001-1043, pl. 1. – In: Ichthyologie. – In: RUSSEGGER, J.: Reise in Griechenland, Unteregypten, im nördlichen Syrien und südöstlichen Kleinasien mit besonderer Rücksicht auf die naturwissenschaftlichen Verhältnisse der betreffenden Länder unternommen in dem Jahre 1836. – Vol. 1/2: 417-1102, 3 maps, 35 pls. – Stuttgart (E. Schweizerbart'sche Verlagshandlung).
- OBRHELOVA, N. (1969): Die Karpfenfische im tschechoslowakischen Süßwassertertiär. – Casopis mineral. geol., **14/1**: 39-52, 5 textfigs. – Prague.
- (1971): Vergleichende Osteologie der Gattung *Leuciscus* (Pisces) aus tertiären Schichten der nördlichen und westlichen CSSR. – Paläont. Abh., (A) **4/3**: 549-660, 80 textfigs., 7 tab., 10 pls. – Berlin (Akademie-Verlag).
- (1990): Fische des Nordböhmischen Braunkohlenbeckens. – Sborník národ. Muz. v Praze (Acta Mus. Nat. Pragae), (B) **46/1990/1-2**: 1-35, 20 textfigs., 10 pls. – Prag.
- PARIN, N.V. (1986): Trichiuridae. – S. 976-980. – In: WHITEHEAD, P.J.P., BAUCHOT, M.-L., HUREAU, J.-C., NIELSEN, J. & TORTONESE, E. (Ed.): Fishes of the North-eastern Atlantic and the Mediterranean, **2**: 511-1008, pl. textfigs. – Paris (UNESCO).
- PROBST, J. (1877): Beiträge zur Kenntniss der fossilen Fische aus der Molasse von Baltringen. II. Batoidei A. Günther. – Jh. Ver. Naturkunde Württemberg, **33**: 69-103, pl. 1-2. – Stuttgart.
- REINECKE, T., STAPF, H. & RAISCH, M. (2001): Die Selachier und Chimären des Unteren Meeressandes und Schlechtsandes im Mainzer Becken (Rupelium, Unteres Oligozän). – Palaeontos, **1**: 73 pp., 8 textfigs., 63 pls. – Antwerpen.

- RÖGL, F. & SPEZZAFERRI, S. (2003): Foraminiferal paleoecology and biostratigraphy of the Mühlbach section (Gaindorf Formation, Lower Badenian), Lower Austria. – Ann. Naturhist. Mus. Wien, **104/A**: 23-75. – Wien.
- ROETZEL, R. (2003): Zur Geologie der mittelmiozänen Fossilfundstelle Mühlbach am Manhartsberg (Niederösterreich). – Ann. Naturhist. Mus. Wien, **104/A**: 3-13. – Wien.
- RUTTE, E. (1962): Schlundzähne von Süßwasserfischen. – Palaeontographica, A **120/4-6**: 165-212, 8 textfigs., 2 tab., pl. 13-21. – Stuttgart.
- & BECKER-PLATEN, J.D. (1980): Cypriniden-Schlundzähne (Pisces) aus dem Känozoikum der Türkei. – Newsl. Stratigr., **8/3**: 191-222, 3 textfigs., 3 tab., 3 pls. – Berlin, Stuttgart.
- & VAN DE WEERD, A. (1980): Cypriniden-Schlundzähne west- und südosteuropäischer Tertiär-Lokalitäten. – Newsl. Stratigr., **8/3**: 223-231, 2 textfigs., 1 pl. – Berlin, Stuttgart.
- SCHULTZ, O. (1971): Die Selachier-Fauna (Pisces, Elasmobranchii) des Wiener Beckens und seiner Randgebiete im Badenien (Miozän). – Ann. Naturhist. Mus. Wien, **75**: 311-341, 1 tab., 4 pls. – Wien.
- (1978): Neue und fehlinterpretierte Fischformen aus dem Miozän des Wiener Beckens. – Ann. Naturhist. Mus. Wien, **81**: 203-219, 1 pl. – Wien.
- (1998): Die Knorpel- und Knochenfischfauna (excl. Otolithen) aus dem Karpat des Korneuburger Beckens (Niederösterreich). – Beitr. Paläont., **23** (Das Karpat des Korneuburger Beckens, Teil 1): 295-323, 4 pls. – Wien.
- SMITH, M. & HEEMSTRA, Ph.C. (1986): Smith's Sea Fishes. – XX + 1047 pp., pl. figs., 144 pls. – Johannesburg (MacMillan South Africa Ltd).
- SYTCHEVSKAYA, E.K. (1989): Neogene Freshwater Fish Fauna of Mongolia. – Soviet-Mongolian Paleont. Expedition, **39**: 144 pp., 41 figs., 32 pls. – Moscow.