

Index of Living and Fossil Echinoids 1971-2008

By Andreas KROH¹

Manuscript submitted on November 20th 2009,
the revised manuscript on January 21st 2010

Abstract

All new taxa of fossil and living echinoids described from 1971 to 2008 are listed with their age, geographic and stratigraphic occurrence, repository of type material and bibliographic citation.

Keywords: Echinodermata, Echinoidea, bibliography, species list, type material.

Introduction

Comprehensive listings of living and fossil echinoids for the species and genera established before 1970 were published by LAMBERT & THIÉRY (1909-25) and KIER & LAWSON (1978) respectively. KIER & LAWSON's supplement for the years 1971-75 to their "Index of Living and Fossil Echinoids 1924-1970" has never been published. More than thirty years have passed since the latter publication and although the advent of information technology and the internet has made taxonomic research much easier, a comprehensive, up-to-date resource for echinoid species is still missing. At genus-level though, echinoids are described in detail in Andrew SMITH's Echinoid Directory (<http://www.nhm.ac.uk/research-curation/research/projects/echinoid-directory/index.html>), an indispensable resource for anyone working with this group.

This list was prepared utilizing a variety of resources, printed and online. The bulk of taxa was located by culling the current echinoderm literature for new taxa and by cross checking this list with the Zoological Record. Citations before 1971 are included if they were absent in LAMBERT & THIÉRY (1909-25) and KIER & LAWSON (1978).

In every case the original paper was consulted and checked for species that might have been missed. Where possible, information on the type material and its repository has been included in this index. No attempt has been made to revise the taxonomic assignment of the species. This index strictly follows the classification of the Treatise on Invertebrate Paleontology, even where I disagree with it. Higher-level taxa are also included in this index.

¹ Natural History Museum Vienna, Department of Geology & Palaeontology, Burgring 7, 1010 Vienna, Austria; e-mail: andreas.kroh@nhm-wien.ac.at

Information in this index is presented in the following way:

Family Name AUTHOR, Date

Genus *Name* AUTHOR, Date

Time horizon

G. species AUTHOR, Date: pages-pages; fig. x; pl. x: figs x-x. [Formation Fm., Locality, Region, Country] (Geological age) <HT(Holotype): ABBREVIATED INSTITUTION Specimen No.; PT(Paratype/s): ABBREVIATED INSTITUTION Specimen No.> {remarks}.

Acknowledgements

Special thanks to Mike REICH, Loïc VILLIER, and the librarians Andrea KOURGLI, Wolfgang BRUNNBAUER and Helga SCHMITZ for their help in locating even the most obscure papers. Johanna KOVAR-EDER, Inesa VISLOBOKOVA, and Oleg MANDIC helped translating Russian papers. Wenyu WU provided translations of Chinese texts.

Stephen K. DONOVAN, John W.M. JAGT, Mathias HARZHAUSER, Gudrun HÖCK, Christine LATAL, Kenneth McNAMARA, Rich MOOI, James H. NEBELSICK, Edeltraud PREIS, Werner E. PILLER, Ortwin SCHULTZ, Andrew B. SMITH, Herbert SUMMESBERGER, and numerous other persons helped in various ways during the preparation of this paper.

This research was supported by the Austrian Science Foundation, project No. P-14366-Bio to Werner E. PILLER (University of Graz) and the SYNTHESYS Project (<http://www.synthesys.info/>) which is financed by European Community Research Infrastructure Action under the FP 6 "Structuring the European Research Area" Programme.

Abbreviations

HT: Holotype

PT: Paratype/s

LT: Lectotype

PLT: Paralectotype/s

ST: Syntype/s

NT: Neotype

Collections:

AM Australian Museum, Sydney, Australia

AMNH American Museum of Natural History, New York, NY, USA

ANSP Academy of Natural Sciences of Philadelphia, Pennsylvania, USA

ANU Department of Geology, Australian National University, Canberra

ASU-GM	Ain Shams University, Geological Museum, Egypt
AU	Auckland University, New Zealand
AUGD	Adelaide Department of Geology and Mineralogy, Adelaide, Australia
AzINEFTEKHIM	Azerbaijan Institute of Gas and Oil Chemistry, Azerbaijan
AZIOC	Azerbaijan Institute of Oil Chemistry, Baku, Azerbaijan
BAS	British Antarctic Survey, Cambridge, UK
BEG	Texas Bureau of Economic Geology, Texas, USA (collections held by the TMM Nonvertebrate Paleontology Laboratory, see TMM)
BFU	Brigham Young University Repository, Provo, Utah, USA
BGS GSM	British Geological Survey, Geological Survey Museum, UK
BMNH	British Museum of Natural History, London, UK
BSPG	Bayerische Staatssammlung für Paläontologie und historische Geologie, München, Germany
BSIP	Museum, Birbal Sahni Institute of Palaeobotany, Lucknow, India
CASG	California Academy of Sciences geology collection, San Francisco, USA
C-EM	Collection COTTEAU, today included in the collection of the Ecole des Mines, Paris, France
CFP	Institut des Sciences de la Terre, Université de Dijon, Dijon, France
CGM	Central Geological Museum F.N.Chernysheva [Chernyshov], Leningrad [today Saint Petersburg], Russia
CGCPE	Colección del Grupo Cultural Paleontológico de Elche y del Museo de Paleontología de Elche, Spain
CGS	Museum of the Central Geological Survey, Republic of China
CGST	Central Geological Survey, Taipei, Taiwan
CM	Canterbury Museum, New Zealand
CNIGR	F.N. Chernyshev Central Research Geological Museum, Saint Petersburg, Russia
CPBA	Cátedra de Paleontología de la Facultad de Ciencias Exactas y Naturales de la Universidad de Buenos Aires, Argentine
CPC	Palaeontological Collection, Bureau of Mineral Resources, Canberra, Australia
CPUC	Colecciones Paleontológicas, Universidad de Concepción, Chile
CWUS	Collection Westlake, University of Southampton, UK

DGM do DNPM	Divisão de Geologia e Mineralogia do Departamento Nacional da Produção Mineral, Rio de Janeiro, Brazil
DGUL	Museum, Department of Geology, University of Lucknow, Lucknow, India
DNSM	Durban Natural Science Museum, Durban, South Africa
DPUW	Department of Paleontology, Faculty of Geology, University of Warsaw, Poland
ECG	Escuela Centroamericana Geología, Universidad de Costa Rica, San José, Costa Rica
EGSG	Department of Earth Sciences (Geology), Universities of Perugia (Italy) and Alicante (Spain) [as in the original publication]
EM	Ecole des Mines, Paris, France
FCDP	Facultad de Ciencias, Departamento de Paleontología, Montevideo, Uruguay
FGWG	FR Geowissenschaften, Ernst-Moritz-Arndt-Universität Greifswald, Germany
FPH	Fundação Paleontológica Phoenix, Aracaju, Brazil
GHUNL-Pam	Facultad de Ciencias Exactas y Naturales, Universidad de La Pampa, Argentina
GIH	Geological Institute, Hungary
GIUS	Museum of Department of Earth Sciences, Silesian University, Poland
GM-GDEMU	Geological Museum of the Geology Department, Faculty of Science, El Mina University, Egypt
GMM	A.P. & M.V. Pavlov Geological Museum, Moskva, Russia
GMUS	Geological Museum of the University of Seville, Spain
GPIH	Geologisch-Paläontologisches Institut Hamburg, Germany
GPIMUM	Geologisch-Paläontologisches Institut und Museum, Universität Münster, Münster, Germany
GPIUB	Geologisch-Paläontologisches Institut, Universität Bonn, Germany
GPIUF	Geologisch-Paläontologisches Institut der Universität Freiburg, Germany
GSI	Geological Survey of India, India
GSIJ	Geological Survey of Japan, Tsukuba, Japan
GSM	Geological Survey Museum (now British Geological Survey), UK
GSP	Geological Survey Pretoria, South Africa
GSWA	Geological Survey of Western Australia, Perth, Australia

GU/R/KE	Laboratory of Vertebrate Palaeontology, Department of Geology, HNB Garhwal University, Srinagar (Garhwal), Uttaranchal, India
HUJ	Zoological Museum, Hebrew University, Jerusalem, Israel
IGEO	Instituto de Geosciências da Universidade Federal do Rio de Janeiro, Brazil
IGM	Instituto de Geología, Universidad Nacional Autónoma de México, Mexico City
IGMCU	Museo de Paleontología del Instituto de Geología, en la Ciudad Universitaria, Mexico
IGN-ANURSR	Institute of Geological Sciences, Academy of Sciences URSR, Kiev, Ukraine
IGS	Institute of Geological Sciences, London, UK
IGPUW	Museum of the Faculty of Geology, University of Warsaw, Warsaw, Poland
IGUFP	Instituto de Geosciências da Universidade Federal de Pernambuco, Brazil
IGUFRJ	Instituto de Geosciências da Universidade Federal de Rio de Janeiro, Brazil
IIGA	Instituto de Investigaciones Geológicas de Antofagasta, Chile
IMBAS	Institute of Marine Biology, Academy of Sciences, Moscow, Russia
IMBFE	Institute of Marine Biology, Far East Center, Academy of Sciences, Vladivostok, Russia
IMGP-Gö	Institut und Museum für Geologie und Paläontologie, Universität Göttingen, Germany
IOANSSR	Institute of Oceanography, Academy of Sciences of the USSR, Moskva
IOAS	Institute of Oceanology, Academia Sinica, Qingdao, China
IORAS	Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia
IPM	Institut de Paléontologie du Muséum, Paris, France
IPPAS	Institute of Paleobiology of the Polish Academy of Science, Warszawa, Poland
KBIN	Koninklijk Belgisch Instituut voor Natuurwetenschappen, Brussels, Belgium (Royal Belgian Institut of Natural History)
LACMIP	Natural History Museum of Los Angeles County, Invertebrate Paleontology Section, California, USA
LGU	Leningrad State University, Saint Petersburg, Russia

LGUT	Laboratoire de Géologie sédimentaire et Paléontologie de l'Université Paul-Sabatier de Toulouse, France
LPB	Palaeontological Collection, University of Bucharest, Romania
LSJU	Stanford University, Regular Collection (transferred to the California Academy of Sciences, San Francisco, CA, USA)
LUGD	Lucknow University Geology Department, Lucknow, India
MACSG (MACS-G)	Maharashtra Association for Cultivation of Sciences, Research Institute, Department of Geology and Palaeontology, Pune, India
MÁFI	Magyar Állami Földtani Intézet Múzeum (Palaeontological Collection of the Museum of the Geological Survey of Hungary), Budapest, Hungary
MBE	Museum für Naturkunde, Berlin, Germany
MBSM	Musée de Boulogne-sur-mer, France
MCA	Museo di Cortina d'Ampezzo, Italy
MCZ	Museum of Comparative Zoology at Harvard University, Cambridge, MA, USA
MGB	Museo de Geologia, Barcelona, Spain
MGCA	Museo Geologico di Castell'Arquato, Italy
MGD-UAA	Museum of the Geological Department, University of Al Ain, United Arabian Emirates
MGIANGSSR	Geological Museum & Institute, Academy of Sciences of the Georgian SSR
MGPD	Dipartimento di Geologia, Paleontologia e Geofisica, Università degli Studi di Padova, Museo di Geologia e Paleontologia, Padova, Italy
MGSB	Museo Geológico del Seminario de Barcelona, Spain
MGU	Minno-Geoložski Universitet, Museum of Geology & Palaeontology, Sofia, Bulgaria
MHNN	Muséum du Havre, France
ML	Muséum d'Histoire naturelle de Lyon, Lyon, France
MMBS	Misaki Marine Biological Station, University of Tokyo, Japan
MMGL	Museo Mineralógico e Geológico da Faculdade de Ciências de Lisboa, Portugal
MMH	Mineralogisk Museum, Copenhagen, Denmark
MN	Museu Nacional do Rio de Janeiro, Brazil
MNA-CPO	Museo nacional de Antropología, Colección Paleontológica F. Oliveras, Uruguay

MNHN	Muséum National d'Histoire naturelle de Paris, France
MNHN-DHT	Muséum National d'Histoire naturelle de Paris, Département Sciences de la Terre, France
MNRJ	Museu Nacional do Rio de Janeiro, Brazil
MOZ	Museo Juan Olsacher, Dirección General de Minería de la Provincia del Neuquén, Neuquén, Argentina
MPEG	Museu Parmense Emílio Goeldi, Pará, Brazil
MSNV	Museo di Storia Naturale, Venice, Italy
MSU	Moscow State University, Moscow, Russia
MUGD	Melbourne University, Geological Department, Melbourne, Australia
MUGM	Geological Museum, El Minia University, Egypt
MV	Museum of Victoria, Melbourne, Australia
MZUB	Museum of Zoology, University of Bergen, Norway
MZUC	Museo de Zoología, Universidad de Concepción, Chile
NHM	Natural History Museum, London, UK
NHMM	Natuurhistorisch Museum at Maastricht, The Netherlands
NHMW	Naturhistorisches Museum Wien, Austria (Natural History Museum Vienna)
NIGP	Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing, China
NMB	Naturhistorisches Museum Basel, Switzerland
NMMNH	New Mexico Museum of Natural History, Albuquerque, New Mexico, USA
NMNH	National Museum of Natural History, USA
NMNZ	National Museum of New Zealand, Wellington, New Zealand
NMP	Czech National Museum, Prague, Czech Republic
NMV	National Museum of Victoria, Australia
NRF	Geological Survey of India, Northern Region, Lucknow, India
NSM	National Science Museum, Tokyo, Japan
NTUG	National Taiwan University, Geological Department, Taipei, Taiwan
NZGS	New Zealand Geological Survey Museum, New Zealand
NZOI	New Zealand Oceanographic Institute, Wellington, New Zealand
OU	Otago University, New Zealand
PASME	Polish Academy of Sciences, Museum of the Earth, Warszawa, Poland

PIASC	Paleontological Institute of the Academy of Sciences of Cuba, Havana, Cuba
PIN	Paleontological Institute, Academy of Sciences of the USSR
PMOZ	Museo Juan Olsacher, Neuquén, Argentina
PRI	Paleontological Research Institution, Ithaca, NY, USA
PUAB	Universitat Autònoma de Barcelona, Palaeontological Collections, Spain
PUPCE	Fossil echinoid collection at the Zoological Department, Punjab University, Lahore, Pakistan
QM	Queensland Museum, Australia
QVM	Queen Victoria Museum, Launceston, Australia
RSM	Royal Scottish Museum, Edinburgh, Scotland, UK
RUCA	Rijksuniversitair Centrum Antwerpen, Belgium
SAfM	South African Museum, Cape Town, South Africa
SAM	South Australian Museum, Adelaide, Australia
SGNP	Servicio Geológico Nacional, Paleontología, Buenos Aires, Argentina
SGWG	Sektion Geologische Wissenschaften der Ernst-Moritz-Arndt Universität Greifswald, Germany
SM	Sedwick Museum of Earth Sciences, University of Cambridge, Cambridge, UK
SMF	Senckenberg-Museum, Frankfurt am Main, Germany
SMNS	Staatliches Museum für Naturkunde Stuttgart, Germany
SNLfB	Niedersächsisches Landesamt für Bodenforschung, Hannover, Germany
SNM	Slovak National Museum, Bratislava, Slovakia
SUI	Repository of the Geological Department, The University of Iowa, Iowa City, USA
SUPTC	Stanford University, Paleontological Type Collection (transferred to the California Academy of Sciences, San Francisco, CA, USA)
TCE	Museum, Department of Geology, University of Lucknow, Lucknow
TGU	Geological & Palaeontological Department, Tbilisskogo Gosudavstevnogo Universiteta, Tbilisi, Georgia
TM	Tasmanian Museum & Art Gallery, Hobart, Australia
TMM	Texas Memorial Museum, Austin, Texas, USA
UA	University of Alaska Museum, Fairbanks, AK, USA

UCMP	University of California, Museum of Paleontology, Berkeley, CA, USA
UF	Florida Museum of Natural History, University of Florida, USA
UFS	Florida State Museum, University of Florida, Gainesville, USA
UI	University of Iowa, Iowa City, Iowa, USA
UILL	University of Illinois, Urbana, Illinois, USA
UIMG	University of Ife, Museum of Geology, Nigeria
UK	Geology Department, University of Kentucky, Lexington, KY, USA
UNAM	Universidad Nacional Autónoma de México, México City, Mexico
UNE	University of New England, Department of Geology, Armidale, N.S.W., Australia
UPNG	Department of Geology, University of Papua New Guinea, Port Moresby, Papua New Guinea
UPS	Université Paris-Sud, Orsay, France
USC	collection of the University of Sofia, Bulgaria
USGS	United States Geological Survey, USA
USNM	United States National Museum, Washington, USA
V	private collection of Alain Vadet, Boulogne-sur-mer, France
VNIGRI	Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta, Leningrad [today Saint Petersburg], Russia
WAM	Western Australian Museum, Perth, Australia
YCM	Yokosuka City Museum, Japan
YPM	Peabody Museum of Natural History, Yale University, New Haven, USA
ZGI	Zentrales Geologisches Institut, Berlin, Germany
ZI	Zoological Institute, Saint Petersburg, Russia
ZMA	Zoölogisch Museum, Universiteit van Amsterdam, The Netherlands
ZMTAU	Zoological Museum, Tel-Aviv University, Israel
ZMUC	Zoologisk Museum, Copenhagen, Denmark
ZPAL	Institute of Palaeobiology of the Polish Academy of Sciences, Warszawa, Poland

Class Echinoidea LESKE, 1778

Genus *Bromidechinus* SMITH & SAVILL, 2001

Bromidechinus SMITH & SAVILL, 2001: 138-141. Type-species: *B. rimaporus* SMITH & SAVILL, 2001.

Ordovician

B. rimaporus SMITH & SAVILL, 2001: 141; figs 3a-b, 4a-d, 5a-c. [Bromide Fm.; Dunn Quarry, Criner Hills, Carter County, Oklahoma, USA] (Blackriverian [= Harnagian, Burrellian, early Caradoc], Upper Ordovician) <HT: BMNH EE6607; PT: BMNH EE6632>.

Subclass Perischoechinoidea M'COY, 1849

Subclass Cidaroidea SMITH, 1984

Superorder Eognathostomata SMITH, 1984

Order Bothriocidaroida ZITTEL, 1879

Family Bothriocidaridae KLEM, 1904

Genus *Bothriocidaris* EICHWALD, 1859

Ordovician

B. kolatai KIER, 1982: 311-314; figs 74A-B; pl. 41, figs 1-2. [Poolville Mb., Bromide Fm., Culley Creek, Criner Hills, Carter County, Southern Oklahoma, USA] (Middle Ordovician) <HT: Graffham coll. 1122TX57>.

B. maquoketensis KOLATA, STRIMPLE & LEVORSON, 1977: 146-149; text-fig. 1a-c; pl. 1, figs 1-8. [Fort Atkinson Formation, Maquoketa Group, SE Fort Atkinson, Winneshiek County, and SW Eldorado, Fayette County, Iowa, USA] (Cincinnatian, Late Ordovician) <HT: SUI 42700; PT: SUI 42701>.

B. solemi KOLATA, 1975: 66; pl. 14, figs 14-15. [Walgreen Mb., Grand Detour Fm., Platteville Grp., Medusa Cement quarry, north side of Route 2, 0.5 miles N of Dixon, Lee County, Illinois, USA] (Middle Ordovician) <HT: UILL X-4882; PT: UILL X-4942 to X-4945>.

B. vulcani GUENSBURG, 1984: 67-68; pl. 14, figs 4-10. [Lebanon Limestone, locality Z-656, Maury County, Central Tennessee, USA] (Middle Ordovician) <HT: UI X-5842; PT: UI X-5841, X-5839>.

Genus *Neobothriocidaris* PAUL, 1967

Ordovician

N. templetoni KOLATA, 1975: 66-68; text-figs 18-19; pl. 14, figs 1-2. [Eagle Point Mb., Dunleith Fm., Galena Grp., Porter Brothers quarry, west of Highway 2, 1 mile S of Rockton, Winnebago County, Illinois, USA] (Middle Ordovician) <HT: UILL X-4882; PT: UILL X-4942 to X-4945>.

Genus *Unibothriocidaris* KIER, 1982

Unibothriocidaris KIER, 1982: 310-311. Type-species: *U. bromidensis* KIER, 1982.

Ordovician

U. bromidensis KIER, 1982: 311; fig. 73; pl. 41, fig. 3. [Poolville Mb., Bromide Fm., Culley Creek, Criner Hills, Carter County, Southern Oklahoma, USA] (Middle Ordovician) <HT: Graffham coll. 1122TX1>.

U. kieri GUENSBURG, 1984: 68; text-fig. 16; pl. 14, figs 12-15. [Lebanon Limestone, locality Z-651, Nashville, Davidson County, Central Tennessee, USA] (Middle Ordovician) <HT: UI X-5832; PT: UI X-5833-5838, X-6036-6044>.

Order Echinocystitioida JACKSON, 1912

Family Echinocystitidae GREGORY, 1897

Genus *Praepholidocidaris* FREST & STRIMPLE, 1977

Praepholidocidaris FREST & STRIMPLE, 1977: 101-102. Type-species: *P. pellaensis* FREST & STRIMPLE, 1977.

Carboniferous

P. pellaensis FREST & STRIMPLE, 1977: 102-105; text-fig. 1a-d, 2a-d; pl. 1, figs 3-9; pl. 2, figs 1-7. [Pella Formation, abandoned county quarry N of Oskaloosa, and inactive quarry NE of Givin, Mahaska County, Iowa, USA] (Mississippian (basal Chesterian), Early Carboniferous) <HT: SUI 39491; PT: SUI 39483-39490, 39492-39502>.

Genus *Pronechinus* KIER, 1965

Lower Permian

P. cretensis KÖNIG, 1982: 169-170; figs 1: 1-2, 2: 1-2. [Fodele Fm., Bali, Northern Crete, Greece] (Upper Carboniferous-Lower Permian, ?Asselian) <HT: GPIUF Pr. 1a+b>.

Family Lepidocentridae LÖVEN, 1874

Genus *Albertechinus* STEARN, 1956

Devonian

A. devonicus BOCZAROWSKI, 2001: 87-92; text-figs 32A, D-E, G-L, 33E, I, L-O, 34A-F, 35E-M; pl. 10, figs 1-4, 12-13, 16; pl. 11, figs 1-13, 15-16; pl. 12, figs 5-7, 12-13. [Marzysz, Holy Cross Mountains, Poland] (Late Givetian. Conodont Early *Mesotaxis falsiovalis* zone) <HT: GIUS 4-568 Mrz./980/1>.

Genus *Aptilechinus* KIER, 1973

Aptilechinus KIER, 1973: 652-654. Type-species: *Aptilechinus caledonensis* KIER, 1973.

Silurian

A. caledonensis KIER, 1973: 654-663; text-figs 1-4; pl. 80, figs 1-3; pl. 81, figs 1-4; pl. 82, figs 1-5; pl. 83, figs 1-3. [Starfish bed, Gutterborn Burn, Pentland Hills, Scotland] (latest Llandovery, Silurian, crenulata-zone) <HT: RSM 1897.32.537B; PT: RSM 1897.32.537A, 1897.32.538A, B, 1897.32.540A, B, 1897.32.541B, 1897.32.543, 1897.32.551, 1897.32.552>.

Genus *Elliptechinus* SCHNEIDER, SPRINKLE & RYDER, 2005

Elliptechinus SCHNEIDER, SPRINKLE & RYDER, 2005: 753-757. Type-species: *Elliptechinus kiwiaster* SCHNEIDER, SPRINKLE & RYDER, 2005.

Carboniferous

E. kiwiaster SCHNEIDER, SPRINKLE & RYDER, 2005: 757-759; figs 4, 5, 10.1-10.6, 11.1-11.5, 12.1-12.2. [Winchell Formation, Canyon Group, Lake Brownwood Spillway about 13.7 km north of Brownwood, Brown County, north-central Texas, USA] (Missourian (Kasimovian), Late Pennsylvanian, Late Carboniferous) <HT: TMM 1967TX23a; PT:

TMM 1967TX23b-d, 1967TX21, 1967TX22, 1967TX25, 1967TX26, 1967TX27, 1967TX54, 1967TX58, 1967TX98, 1967TX101>.

Genus *Kongielechinus* JESIONEK-SZYMANSKA, 1979

Kongielechinus JESIONEK-SZYMANSKA, 1979: 284-286. Type-species: *Kongielechinus magnituberculatus* JESIONEK-SZYMANSKA, 1979.

Devonian

K. magnituberculatus JESIONEK-SZYMANSKA, 1979: 286-289; figs 1C, 2A-E, 3A-C, 4A, 5, 6; pl. 17, figs 1-3; pl. 18, figs 1-3; pl. 19, figs 1-4; pl. 20, figs 1-4; pl. 21, figs 1-5; pl. 22, figs 1-3; pl. 23, figs 1-4. [Świętomierz-Śniadka profile, Błonie Valley, Holy Cross Mts., Poland] (Givetian, Middle Devonian) <HT: ZPAL ED 31; PT: ZPAL ED 22, 32>.

Genus *Pholiocidaris* MEEK & WORTHEN, 1869

Upper Carboniferous

P. hungaricus MIHÁLY, 1990b: 251-252; pl. 1: fig. 1. [Mályinka Fm., Nagyvisnyó, Bükk Mts., Hungary] (Moskovian) <HT: MÁFI Coll. Ech. C. 2363>.

Order Palaechinoida HAECKEL, 1866

Family Palaechinidae M'COY, 1849

Genus *Lovenechinus* JACKSON, 1912

Carboniferous

L. zunyii LI, HE & K.F.ZHANG, in LI, ZHANG, HE & ZHANG, 1985: 99 [Chinese], 103 [Engl.]; pl. 2: figs 5-7. [Xing-an, Guangxi Province, South China] (Early Carboniferous) <HT: GBG-LH-01 (repository not given; presumably Wuhan College of Geology)>.

Genus *Melonechinus* MEEK & WORTHEN, 1861

Carboniferous

M. guangxiensis HE & LI, in LI, ZHANG, HE & ZHANG, 1985: 100-101 [Chinese], 103 [Engl.]; figs 1-3; pl. 1: fig. 10; pl. 2: figs 8-12. [Gueilin, Guangxi Province, South

China] (Early Carboniferous) <HT: GBG-HL-07 (repository not given; presumably Wuhan College of Geology)>.

Genus *Palaechinus* M'COY, 1844

Carboniferous

P. jacksoni CHESNUT & ETENSOHN, 1988: 62-63; text-fig. 27; pl. 11, fig. 4. [Locality 1, Cincinnati-Southern Railroad cut near Sloans Valley, Pulaski County, Kentucky, USA] (Middle Chesterian, Upper Mississippian) <HT: USNM 372191>.

P. yangi X.P. ZHANG & LI, in LI, ZHANG, HE & ZHANG, 1985: 97-98 [Chinese], 102 [Engl.]; pl. 2: figs 1-2. [Hunan Province, South China] (Early Carboniferous) <HT: XBG-ZL-01 (repository not given; presumably Wuhan College of Geology)>.

Devonian

P. praematurus NESTLER, 1981: 556-557; figs 1-3; pl. {1}, figs 1-3. [Fleckenberg, Bl. Schmallenberg, Sauerland, Germany] (Early Devonian) <HT: ZGI > {no specimen no. given}.

Family Cravechinidae HAWKINS, 1946

Genus *Cravechinus* HAWKINS, 1946

Devonian

C.? *frankei* HAUDE, 1999: 152-158, 165-166; figs 2A-C, 3A-B, 4. [quarry Iberg-Winterberg, Harz Mts., NW Göttingen, Germany] (rhenana-zone, Adorf stage, Late Frasnian, Late Devonian) <HT: IMGP-Gö 1119-1> {based on a single, fragmented specimen}.

Family Proterocidaridae SMITH, 1984

Proterocidaridae SMITH, 1984: 147. Type-genus: *Proterocidaris* DE KONINCK, 1882 [USA, Europe] (Carboniferous to Permian).

Order Cidaroida CLAUS, 1880

Family Anisocidaridae VADET, 1999a

Anisocidaridae VADET, 1999a: 72. Type-genus: *Anisocidaris* THIÈRY, 1928.

Family Archaeocidaridae M'COY, 1844

Genus *Aarchaeocidaris* WEBSTER, 1997

Aarchaeocidaris WEBSTER, 1997: 37. Type-species: *Aarchaeocidaris strawberryensis* WEBSTER, 1997.

Carboniferous

A. strawberryensis WEBSTER, 1997: 37; pl. 4, figs 1, 4; pl. 8, figs 4-6, 9. [Lodgepole Formation and Brush Canyon Member, Henderson Canyon Formation, South wall of Strawberry Creek, Wyoming, USA and Beirdneau Hollow, Logan Canyon, Utah, USA] (Kinderhookian (= Tournaisian), Mississippian, Early Carboniferous) <HT: USNM 487247 (from Strawberry Creek, Wyoming, USA); PT: USNM 487248, 487249>.

Genus *Archaeocidaris* M'COY, 1844

Upper Permian

A. hamata MIHÁLY, 1980: 400-401; pl. 1: figs 1-5; pl. 2: figs 1-4 [Nagyvisnyó, Komitat Borsod-Abauj-Zemplén, Northern Hungary] <HT: MÁFI P. 1251> {based on isolated primary spines}.

A. schréteri MIHÁLY, 1980: 401-402; pl. 3: figs 1-4 [Nagyvisnyó, Komitat Borsod-Abauj-Zemplén, Northern Hungary] <HT: MÁFI P. 1359> {based on isolated primary spines}.

Carboniferous

A. apheles SCHNEIDER, SPRINKLE & RYDER, 2005: 753; figs 9.1-9.4. [Winchell Formation, Canyon Group, Lake Brownwood Spillway about 13.7 km north of Brownwood, Brown County, north-central Texas, USA] (Missourian (Kasimovian), Late Pennsylvanian, Late Carboniferous) <HT: TMM 1967TX20; PT: TMM 1967TX17, 1967TX18, TMM 1967TX55>.

A. brownwoodensis SCHNEIDER, SPRINKLE & RYDER, 2005: 747-753; figs 2-5, 6.1-6.6, 7.1-7.5, 8.1-8.3. [Winchell Formation, Canyon Group, Lake Brownwood Spillway about 13.7 km north of Brownwood, Brown County, north-central Texas, USA] (Missourian (Kasimovian), Late Pennsylvanian, Late Carboniferous) <HT: TMM 1967TX60, PT: TMM 1967TX1, 1967TX3, 1967TX2, 1967TX7, 1967TX16, 1967TX35, 1967TX62, 1967TX99, 1967TX61, 1967TX100, 1967TX101>.

A.? *diadematoides* HAUDE & THOMAS, 1994: 124-125; fig. 6; pl. 2: figs I-O. [Kulm-Tonschiefer, road cut "Kohleiche", Federal road B224, N Wuppertal, Germany] (cu III α 1,

Early Carboniferous) <HT: SMF [colln Thomas] Eko-667.1> {based on a single disarticulated specimen}.

- A. faassi* AISENBERG, 1974: 76-79; pl. XY: figs 1-63. [Limestone bed K5, River Volchya, Donbas, western part of the Donetz Basin, Ukraine] (Middle Carboniferous) <ST: IGN-ANURSR [no numbers given]>.
- A. hemispinifera* CHESNUT & ETTENSOHN, 1988: 63-64; text-fig. 28; pl. 11, figs 5-9. [Locality 3, Strunk Construction Company Quarry, near Tatesville, Pulaski County and locality 5, Laurel County Quarry, Laurel County Kentucky, USA] (Middle Chesterian, Upper Mississippian) <HT: UK 115989>.
- A. whatleyensis* LEWIS & ENSOM, 1982: 81-97; figs 1-19. [Whatley Quarry, Wathley, near Frome, Somerset, England] <HT: BMNH E.76888; PT: BMNH E.76887, E.76889>.

Family Miocidaridae DURHAM & MELVILLE, 1957

Genus *Miocidaris* DÖDERLEIN, 1887

Late Triassic

- M. adrianae* ZARDINI, 1973: 9; pl. 6: fig. 25. [Cassian Formation, Misurina, Cortina d'Ampezzo, Italy] (Carnian) <HT: MCA M12>.
- M. ampezzana* ZARDINI, 1973: 9; pl. 6: figs 23a, b. [Cassian Formation, Campo, Cortina d'Ampezzo, Italy] (Carnian) <HT: MCA C11>.
- M. cortinensis* ZARDINI, 1973: 9; pl. 6: figs 28, 29; pl. 19: fig. 7. [Cassian Formation, Mi-lieres and Sasso Stria, Cortina d'Ampezzo, Italy] (Carnian) <HT: MCA C13>.

Genus *Triadocidaris* DÖDERLEIN, 1887

Middle Triassic

- T. hungarica* MIHÁLY, 1981: 301-302; pl. 1: fig. 5. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI P. 5500> {based on test fragments}.

Family Cidaridae GRAY, 1825

Subfamily Histocidarinae MORTENSEN, 1928

Tribe Histocidarini MORTENSEN, 1928

Genus *Histocidaris* MORTENSEN, 1903

Recent

H. longicollis HOGGETT & ROWE, 1986: 2-4; pl. 1: figs A-C. [south of New Caledonia (23° S, 167°17' E, depth 420 m), South-west Pacific Ocean] (Recent) <HT: AM J19173>.

Pliocene

H. sicula BORGHI, 1999: 116-117; pl. 4, figs 1-4, 8, 10-11. [near Milazzo, Northern Sicily, Italy] (Pliocene) <HT: MGCA, specimen no. not given>.

Genus *Polycidaris* QUENSTEDT, 1857

Upper Jurassic

P. nusplingensis GRAWE-BAUMEISTER, SCHWEIGERT & DIETL, 2000:11-12; pl. 1: fig. 6; pl. 3: fig. 1; pl. 4: figs 1-2; pl. 5: figs 1-3. [Nusplingen Lithographic Limestone, Nusplingen, SW-Germany, Europe] (Upper Kimmeridgian) <HT: SMNS 64267/1; PT: SMNS 64104> {placed into the family Rhabdocidaridae LAMBERT, 1900 by GRAWE-BAUMEISTER et al., 2000: 11}.

Tribe Poriocidarini SMITH & WRIGHT, 1989

Poriocidarini SMITH & WRIGHT, 1989: 16. (Bajocian – Recent) {included genera: *Poriocidaris* MORTENSEN, 1909; *Plegiocidaris* POMEL, 1883}.

Subfamily Ctenocidarinae MORTENSEN, 1928

Genus *Aporocidaris* AGASSIZ & CLARK, 1907

Recent

A. eltaniana MOOI, DAVID, FELL & CHONÉ, 2000: 227-230; fig. 2. [R/V Eltanin stations (Cruise/Station: 4/138, 6/432), Region of Livingston Island, South Shetland Islands, Antarctica] <HT: NMNH E48122; PT: NMNH E11161, E11188, E11212, E11290, E14597, MCZ 8406>.

A. usarpi MOOI, DAVID, FELL & CHONÉ, 2000: 230-233; fig. 3. [R/V Eltanin station (Cruise 14, station 1212), mid-ocean point far to the southeast of New Zealand, North of the Ross Sea, Antarctica] <HT: NMNH E11134; PT: NMNH E11059, E14603>.

Genus *Austrocidaris* H. L. CLARK, 1907

Recent

A. pawsoni MCKNIGHT, 1974: 26-28; figs 1a-b. [NZOI Stat. E148 (44°30' S, 177°45' W to 44°30.2' S, 177°45.2' W, 880 m depth), F108 (48°19' S, 171°59' E, 1108 m depth), G700 (46°20' S, 171°15' E, 1116 m depth), New Zealand] (Recent) <HT: NZOI 181 (Stn F108); PT: NZOI P237 (Stn F148), P238 (Stn G700)>.

Eocene

A. seymourensis RADWANSKA, 1996: 120-122; figs 2-3; pl. 28: figs 1-4; pl. 29: figs 1-3; pl. 30: figs 1-5; pl. 31: figs 1-3. [ZPAL 1 (Bill Hill), Seymour Island, Antarctic Peninsula] <HT: IPPAS ZPAL E. VIII/1 PT: IPPAS ZPAL E.VII/2-5>.

Genus *Notocidaris* MORTENSEN, 1909

Recent

N. bakeri MCKNIGHT, 1974: 28-30; figs 2a-b. [NZOI Stat. D593a (41°00' S, 178°25' E, 3017-3109 m depth), New Zealand] (Recent) <HT: NZOI 182>.

N. lanceolata MOOI, DAVID, FELL & CHONÉ, 2000: 233-236; fig. 4. [R/V Eltanin stations (Cruise/Station: 27/1867, 27/1926, 27/1929, 32/2002, 32/2108, 32/2110, 32/2129), Ross Sea, Antarctica] <HT: MCZ 8336; PT: NMNH E21865, E21866, E22004, E22005, E22006>.

Genus *Ctenocidaris* MORTENSEN, 1910

Recent

C. aotearoa MCKNIGHT, 1974: 30-33; figs 3a-b. [NZOI Stat. D591 (42°28.8' S, 176°51.2' E, 1829 m depth), F123 (47°38' S, 178°57' W, 1280 m depth), F127 (49°22' S, 176°16' E, 1280 m depth), G701 (46°20' S, 171°30' E, 1400 m depth), G704 (46°17' S, 172°37' E, 1600 m depth), New Zealand] (Recent) <HT: NZOI 183 (Stn F123); PT: NZOI P239 (Stn F123), P240 (Stn F127), P241 (Stn D591)>.

Subfamily Goniocidarinae MORTENSEN, 1928

Genus *Goniocidaris* DESOR, in AGASSIZ & DESOR, 1846

Miocene

“*G.*” *noetlingi* ROMAN, 1976: 22-24; pl. 4, figs 1-12. [Upper Dam Fm., NNE of Qarn Abu Wail, Qatar] (Early-mid Miocene) <HT: MNHN 1975-16> {established for *Cidaris* sp. 1 of NOETLING, 1901}.

Subfamily Stereocidarinae LAMBERT, 1900

Genus *Hirudocidaris* SMITH & WRIGHT, 1989

Hirudocidaris SMITH & WRIGHT, 1989: 78. Type-species: *Cidaris hirudo* SORIGNET, 1850. [Central and NW Europe] (Cenomanian to Campanian) {placed into the Tribe Stereocidarini Mortensen, 1928 within the Subfamily Cidarinae GRAY, 1825 by SMITH & WRIGHT (1989); other species included: *Cidaris uniformis* SORIGNET, 1850; *Stereocidaris silesiaca* SCHLÜTER, 1892}.

Genus *Phalacrocidaris* LAMBERT, 1902

In the *Treatise* this genus is considered as junior synonym of *Stereocidaris* POMEL, 1883.

Campanian

P. catherinae SMITH & WRIGHT, 1989: 77-78; text-fig. 15D; pl. 19, fig. 5. [Danes Dyke, Sewerby, Humberside, England] (*Inoceramus lingua* Zone, Campanian) <HT: BMNH E82345>.

Genus *Stereocidaris* POMEL, 1883

Early Cretaceous

S. trigonodus AZIZ, 1991: 18; pl. 1, fig. 12. [Dalmiapuram Formation, Trichinopoly Subbasin, Southern India] (Late Aptian to Early Albian) <HT: MACSG 1936; PT: MACSG 1937-1939> {based on isolated spines}.

Genus *Temnocidaris* COTTEAU, 1863

Maastrichtian

T. schlueteri SALAH, 1982: 212-213; pl. 2, figs 1-8. [Hemmoor, North-western Germany] (Late Maastrichtian) <HT: SNLFB kma 281>.

Genus *Typocidaris* POMEL, 1883

In the Treatise, FELL (1966: U325) considered this genus as synonymous to *Stereocidaris* POMEL, 1883. GEYS (1987: 202) reestablished *Typocidaris* as separate genus. VADET (1988: 130) erected a separate family for this genus (→ Typocidaridae).

Campanian-Maastrichtian

T. ubaghsi GEYS, 1987: 206-207; pl. 2, figs 5-9. [Lower Gulpen Fm., Heure-le-Romain, Liège, Belgium and Rügen, Germany] (Upper Campanian-Lower Maastrichtian) <HT: KBIN IST 10253>.

Subfamily Rhabdocidarinae LAMBERT, 1900

Genus *Desoricidaris* GEYS, 1992

Desoricidaris GEYS, 1992: 140-141. Type-specie: *Rhabdocidaris pouyannei* COTTEAU, 1863. [Europe and Northern Africa] (Cretaceous) {other species included: *D. sanctaerucis* (COTTEAU, 1862); *D. salviensis* (COTTEAU, 1857); *D. venulosa* (AGASSIZ & DESOR, 1846); *Leiocidaris balli* FOURTAU, 1914; *D. bonolai* (GAUTHIER, in FOURTAU, 1900); *D. subvenulosa* (COTTEAU, PERON & GAUTHIER, 1880); *D. schweinfurthi* (GAUTHIER, in FOURTAU, 1901); *D. crameri* (DE LORIO, 1887)}.

Genus *Fellius* CUTRESS, 1980

Fellius CUTRESS, 1980: 116. Type-species: *Cidaris foveata* JACKSON, 1922. [Jamaica and Cuba] (Eocene).

Genus *Phyllacanthus* BRANDT, 1835

Miocene

P. priscus BRITO & RAMIRES, 1974: 264-266; pl. 1, figs 1, 5-7. [Pirabas Fm., Ilha de Fortaleza, Município de Primavera, Pará, Brazil] (Lower Miocene) <HT: MN 5238-I; PT: IGEO 302, 304>.

Genus *Prionocidaris* A. AGASSIZ, 1863

Recent

P. callista ROWE & HOGGETT, 1986: 252-256; figs 12A-B, 13B. [Bushy Island, Queensland to Montague Island, New South Wales, Australia; Lord Howe Island; Norfolk Island; Ker-madec Island; New Caledonia] (Recent) <HT: AM J15715; PT: AM J18919-18923>.

P. popeae HOGGETT & ROWE, 1986: 7-10; pl. 4: figs A-C; pl. 5: figs A-B. [off Ile des Pins (22°50' S, 167°34' E, depth 275 m), New Caledonia; 1.5 miles south of Aneityum Island (20°17' S, 169°48' E, depth 55-75 m), Vanuatu; and south of New Caledonia (22°48' S, 167°36.5' E, depth 85-100 m), South-west Pacific Ocean] (Recent) <HT: AM J19206 (from Ile des Pins); PT: AM J8192, J13887, J19207-J19210 (all from Ile des Pins)>.

Miocene-Pliocene

P. cookei CUTRESS, 1976: 192-197; figs 1A-F, 2A-B. [Chipola Formation, USGS loc. 2564, right bank of Chipola River, on the McClelland Farm, one mile below the bridge at the old Bailey's Ferry; USGS loc. 2213, one mile below Bailey's Ferry on the Chipola River; USGS loc. 3419, McClelland's Farm on the west bank of Chipola River, one mile below Bailey's Ferry; all 10 miles W of Blountstown, Calhoun County Florida, USA] (Middle Miocene) <HT: USNM 232503 (from USGS locality 2564); PT: USNM 232504-232506> {based on isolated spines and interambulacral plates}.

P. katherinae CUTRESS, 1980: 94-99; pl. 9, figs 4-10. [Cuba and Cibao Fm., Puerto Rico] (Middle Miocene to Pliocene) <HT: PRI 29661; PT: AMNH 18566/1, 2, ANSP 50983, MCZ 4102, PRI 29635, 29720, 29746, 29749-29751>.

Oligocene

[*P. placenta* SRIVASTAVA, 1988]: 152. [India] (*Lepidocyclina* (*Eulepidina*) Zone, Oligocene) <none defined> {nomen nudum}.

Maastrichtian

P.? *emiratus* SMITH, 1995: 133; fig. 9C; pl. 1: figs 3, 5-6. [Jebel Huwayyah, Oman/ United Arabian Emirates Border Region] <HT: BMNH EE3431> {SMITH, in SMITH &

JEFFERY (2000: 17-19) placed this species into the synonymy of *Phyllacanthus suleimani* (NOETLING, 1897)}.

Albian

P. neglecta SMITH & WRIGHT, 1989: 90-92; text-figs 17A-D, 18A; pl. 29, figs 3-4; pl. 30, fig. 1. [Gault of Folkestone, Kent, and Red Chalk of Hunstanton, Norfolk, England] (Upper Albian) <HT: BMNH E1078; PT: BMNH E82452, E82453, E82495, E19860-1>.

Genus *Prophyllacanthus* CUTRESS, 1980

Prophyllacanthus CUTRESS, 1980: 120-122. Type-species: *Leiocidaris leoni* LAMBERT & SÁNCHEZ ROIG, in SÁNCHEZ ROIG, 1926. [Jamaica and Cuba] (Maastrichtian).

Eocene

P. eocenicus CUTRESS, 1980: 127-131; pl. 12, figs 1-4. [Jabaco Fm., Cuba] (Eocene) <HT: SUPTC 10249a; PT: LSJU 53248, 53249, PRI 29633, 29634, SUPTC 10249b-e, 10250, 10251>.

Subfamily Cidarinae GRAY, 1825

Genus *Almucidaris* BLAKE & ZINSMEISTER, 1991

Almucidaris BLAKE & ZINSMEISTER, 1991: 632. Type-species: *Almucidaris durhami* BLAKE & ZINSMEISTER, 1991

Maastrichtian

A. durhami BLAKE & ZINSMEISTER, 1991: 632-635; text-fig. 2; pl. 1, figs 1-6. [Lopez de Bertodano Fm., Seymour Island, Antarctica] <HT: USNM 446322; PT: USNM 446323-446348>.

Genus *Balanocidaris* LAMBERT, 1910

Albian

B. clubis AZIZ, 1991: 21; pl. 1, fig. 14. [Dalmiapuram Formation, Trichinopoly Sub-basin, Southern India] (Late Aptian to Early Albian, Early Cretaceous) <HT: MACSG 1941> {based on isolated spines}.

B. fusiformis SMITH & WRIGHT, 1989: 25; pl. 8, figs 6-8. [Regularis Limestone, Acre Pit, Shenley Hill, Leighton Buzzard, Bedfordshire, England] (*L. tardefurcata* Zone, Lower Albian) <HT: BMNH E37948; PT: E82348-50>.

Genus *Calocidaris* H. L. CLARK, 1907

Miocene

C. palmeri CUTRESS, 1980: 55-59; pl. 3, fig. 1. [Güines Fm., Basal Yumurí Limestone, Palmer locality 978, Habana Province, Cuba] (Middle Miocene) <HT: ANSP 50974>.

Genus *Chesniericidaris* VADET, in VADET & CHESNIER, 2003

Chesniericidaris VADET, in VADET & CHESNIER, 2003: 27. Type-species: *Cidaris wrightii* DESOR, 1865. [Crickley Hill] (Bajocian, aujourd'hui Aalénien (zone à Murchinson)).

Jurassic

C. brangeri VADET & NICOLLEAU, 2005: 16; illustr. on p. 16. [Region of Rich, Morocco] (Late Domerian (= Late Pliensbachian), Jurassic) <Type-series: Roland Reboul coll'n 548a, 548b, Philippe Nicolleau coll'n 9178 (?8952), Alain Vadet coll'n V6725>.

Genus *Cidaris* LESKE, 1778

(= Genus *Cidarites* AUCTT.)

Oligocene-Miocene

C. bermudezi CUTRESS, 1980: 49-53; pl. 2, figs 2-8. [Cojímar Fm. and Güines Fm., Basal Yumurí Limestone, Cuba; Lares Fm. and Juana Dáz Fm., Puerto Rico] (Oligocene to Middle Miocene) <HT: PRI 29603; PT: PRI 29600-02, 29604-07, 29745, 29751>.

Upper Cretaceous

C. comptoni GLAUERT, 1923: 48-49; pl. 3. [Gingin Chalk, Gingin, Perth Basin, Western Australia] (Santonian-Campanian) <HT: WAM G3775> {placed into the genus *Goniocidaris* by McNAMARA (1986: 353-354)}.

Jurassic

Cidarites burckhardti LARRAIN, 1975: 38-42; figs 21-31, tab. 2. [Lonquimay, Malleco Province, Chile] (Rauracian, Late Jurassic) <HT: Colección del Departamento de Geología y Paleontología, Universidad de Concepcion, Chile (1 spine); PT: as HT (10 spines)> {based on isolated spines}.

C. leberti VADET, in CHEVET & RIGOLLET, 1989: 18; un-numbered fig. on p. 18. [Oolithe de Chemilly, Sarthe, France] (*K. jason* Zone, Middle Callovian) <HT: Alain VADET coll. no. V1558> {placed into the genus *Anisocidaris* THIÉRY, 1928 by VADET, 1991}.

Late Triassic

C. aialensis [*aculeata*] ZARDINI, 1973: 21; pl. 12: figs 24a-b, 25a-b, 26-35. [Cassian Formation, Campo, Forcella Giau, Milieres, Misurina, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 33> {mentioned as "*C. n.sp aculeata*" in part of the plate descriptor; based on isolated spines}.

C. aialensis fusiformis [*aculeata ailaensis*] ZARDINI, 1973: 21; pl. 18: fig. 28. [Cassian Formation, Sasso di Striae, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: repository and specimen no. not given > {mentioned as "*C. n.sp aculeata n.sp. ailaensis*" in the plate descriptor; based on isolated spines}.

C. alata brevicaulis ZARDINI, 1973: 19; pl. 9: fig. 25; pl. 10: figs 2a-c, 9, 10, 11a-e, 22a-c, 25; pl. 15: fig. 12; pl. 18: fig. 21. [Cassian Formation, Campo, Milieres, Misurina, Sasso Stria, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 28> {based on isolated spines}.

C. alata canaliculata ZARDINI, 1973: 20; pl. 9: figs 20a, b; pl. 10: figs 17a-d. [Cassian Formation, Campo, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 31> {based on isolated spines}.

C. alpina ZARDINI, 1973: 20; pl. 7: fig. 27; pl. 21: fig. 3. [Cassian Formation, Forcella Giau, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 32> {based on isolated spines}.

C. biconica ZARDINI, 1973: 18; pl. 9: figs 1-5, 6a, b, 7, 8, 9a, b; pl. 21: fig. 5. [Cassian Formation, Alpe di Specie, Misurina, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 25> {based on isolated spines}.

C. coralliophila ZARDINI, 1973: 23; pl. 18: figs 26-27. [Cassian Formation, Sasso Stria, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA SS 43> {based on isolated spines}.

C. costalarensis ZARDINI, 1973: 21; pl. 12: figs 4-11. [Cassian Formation, Costalaresc, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA CO 34> {based on isolated spines}.

C. costata ZARDINI, 1973: 19-20; pl. 10: figs 15a-b, 20a-b. [Cassian Formation, Alpe di Specie and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 29> {based on isolated spines}.

C. costata tuberculata ZARDINI, 1973: 20; pl. 10: figs 21a-b. [Cassian Formation, Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA T 30> {based on isolated spines}.

- C. costeanensis* ZARDINI, 1973: 21-22; pl. 10: figs 24a-b; pl. 17: figs 22, 23a-b; pl. 18: fig. 37. [Cassian Formation, Campo and Sasso Stria, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 35> {based on isolated spines}.
- C. dorsata alata* ZARDINI, 1973: 17-18; pl. 10: figs 1a-c; pl. 9: figs 18a-c, 19a-c. [Cassian Formation, Forcella Giau, and Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA MI 24> {based on isolated spines}.
- C. dorsata fusiformis* ZARDINI, 1973: 17; pl. 8: figs 16-19. [Cassian Formation, Milieres, Misurina, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA M 23> {based on isolated spines}.
- C. flexuosa brevicostata* ZARDINI, 1973: 26; pl. 15: figs 19, 23a-b. [Cassian Formation, Campo and Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 47> {based on isolated spines}.
- C. forminensis* [*forninensis*] ZARDINI, 1973: 18; pl. 9: figs 13a, b; pl. 17: figs 9a-c; pl. 20: fig. 35; pl. 21: fig. 8. [Cassian Formation, Forcella Giau, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA G 27> {named “*C. n.sp. forminensis*” in part of the plate descriptions; based on isolated spines}.
- C. giauensis* ZARDINI, 1973: 12; pl. 20: figs 34a-b. [Cassian Formation, Forcella Giau, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA G 17>.
- C. glabra* ZARDINI, 1973: 23; pl. 12: figs 37-39; pl. 18: fig. 33. [Cassian Formation, Alpe di Specie, Campo, Sasso Stria, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 42> {based on isolated spines}.
- C. lancedelli* ZARDINI, 1973: 12-13; pl. 5: fig. 24a-b. [Cassian Formation, Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA MI 18>.
- C. leonardi* ZARDINI, 1973: 11-12; pl. 5: fig. 13. [Cassian Formation, Cianzopè, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA R 15>.
- C. lineola campanulata* ZARDINI, 1973: 26-27; pl. 15: figs 17a-b, 18a-b. [Cassian Formation, Campo and Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA MI 48> {based on isolated spines}.
- C. milierensis* ZARDINI, 1973: 23; pl. 12: figs 19-20, 21a-b. [Cassian Formation, Campo and Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 41> {based on isolated spines}.
- C. montanaro* ZARDINI, 1973: 12; pl. 5: figs 14, 15a, b. [Cassian Formation, Alpe di Specie and Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 16>.
- C. ovata* ZARDINI, 1973: 18; pl. 8: figs 23a-c, 24a, b, 25a, b, 26a-c. [Cassian Formation, Campo and Tamari, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 26> {based on isolated spines}.

- C. plana* ZARDINI, 1973: 22; pl. 12: figs 45a-b; pl. 15: fig. 20; pl. 19: figs 2a-c; pl. 22: fig. 4. [Cassian Formation, Campo and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA MI 37> {based on isolated spines}.
- C. quadrialata* ZARDINI, 1973: 28; pl. 10; fig. 27. [Cassian Formation, Milieres, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA MI 49> {based on isolated spines}.
- C. quadriserrata* ZARDINI, 1973: 24-25; pl. 13: figs 1, 2, 4; pl. 22: fig. 6. [Cassian Formation, Alpe di Specie, Milieres, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 44> {based on isolated spines}.
- C. reticulata* ZARDINI, 1973: 25; pl. 13: figs 9, 10a-b, 11a-b, 12. [Cassian Formation, Alpe di Specie, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 45> {based on isolated spines}.
- C. rinbianchi* ZARDINI, 1973: 11; pl. 5: figs 1a-b, 2a-b; pl. 6: fig. 12. [Cassian Formation, Alpe di Specie, Campo, and Misurina, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA M 14>.
- C. seelandica* ZARDINI, 1973: 16-17; pl. 11: figs 1-5, 6a-b, 7a-c, 8, 9a-b, 10a-b, 11; pl. 21: fig. 4. [Cassian Formation, Alpe di Specie, Carbonin, and Misurina, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA M 21> {based on isolated spines}.
- C. seelandica misurinensis* ZARDINI, 1973: 17; pl. 7: figs 6a, b, 7a, b, 8a, b, 9a, b, 10a, b. [Cassian Formation, Misurina, Campo, and Rumerlo, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 22> {based on isolated spines}.
- C. spongiosa* ZARDINI, 1973: 22; pl. 12: figs 47a-b, 48a-b, 49a-b, 50; pl. 17: figs 5a-b, 6a-b, 7a-b, 8a-b; pl. 18: figs 10a-b, 11a-c, 12, 13a-b. [Cassian Formation, Forcella Giau, Milieres, Sasso Stria, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA SS 39> {based on isolated spines}.
- C. sulcata* ZARDINI, 1973: 22; pl. 12: figs 12a-b, 13-17. [Cassian Formation, Milieres, Misurina, and Tamarin, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA M 36> {based on isolated spines}.
- C. tenuicostata* ZARDINI, 1973: 25; pl. 13: figs 8a-b; pl. 19: figs 4a-b; pl. 22: fig. 5. [Cassian Formation, Campo, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA C 46> {based on isolated spines}.
- C. tetraedrica* ZARDINI, 1973: 16; pl. 7: figs 21, 22a, b, 23-26. [Cassian Formation, Alpe di Specie, Milieres, and Misurina, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA S 19> {based on isolated spines}.
- C. tetraedrica appendiculata* ZARDINI, 1973: 16; pl. 18: figs 14-15, 16a-b, 17a-d, 18a-c, 19a-d. [Cassian Formation, Sasso Stria, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA SS 20> {based on isolated spines}.

- C. trapezoidalis* ZARDINI, 1973: 22; pl. 12: figs 42a-b, 43a-b, 44a-b, 46a-b. [Cassian Formation, Forcella Giau, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA G 38> {based on isolated spines}.
- C. valparolae* ZARDINI, 1973: 28; pl. 18; figs 30-31. [Cassian Formation, Sasso Stria, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA SS 50> {based on isolated spines}.
- C. verticillata* [*spongiosa*] ZARDINI, 1973: 23; pl. 18: figs 6, 7a-c, 8, 9a-c. [Cassian Formation, Sasso Stria, Cortina d'Ampezzo, Italy] (Carnian, Late Triassic) <HT: MCA SS 40> {labelled as "*C. n.sp. spongiosa*" in the plate description; based on isolated spines}.

Middle Triassic

- C. adunca* MIHÁLY, 1981: 309-310; pl. 6: figs 3-5. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI T. 5441; PT: MÁFI T. 5440, T. 5442b> {based on isolated primary spines}.
- C. aggtelekensis* MIHÁLY, 1981: 308; pl. 5: figs 4-5. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI T. 5293c> {based on isolated primary spines}.
- "*C.*" *bangtoupensis* STILLER, 2001: 535-536; pl. 1, figs 1-10. [Leidapo Member, Yuqing Subformation, Qingyan Formation, Bangtoupou, NNE of Qingyan, Guizhou Province, south-western China] (early Late Anisian, Middle Triassic) <HT: GPIMUM B3A-6.E2.Fbt-1; PT: GPIMUM B3A-6.E2.F61-1, B3A-6.E2.F61-2>.
- C. batheri* MIHÁLY, 1981: 307-8; pl. 4: figs 8-9. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI T. 5253b> {based on isolated primary spines}.
- "*C.*" *fangchui* STILLER, 2001: 539-540; text-figs 6g-n. [Leidapo Member, Yuqing Subformation, Qingyan Formation, Bangtoupou, NNE of Qingyan, Guizhou Province, south-western China] (early Late Anisian, Middle Triassic) <HT: GPIMUM B3A-6.E6.F60-1; PT: GPIMUM B3A-6.E6.F53-1, B3A-6.E6.F61-1>.
- C. gladius* MIHÁLY, 1981: 304; pl. 2: fig. 5; pl. 3: fig. 1. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI T. 5340> {based on isolated primary spines}.
- "*C.*" *gu* STILLER, 2001: 541-546; text-figs 7a-j, 8a-l. [Leidapo Member, Yuqing Subformation, Qingyan Formation, Bangtoupou, NNE of Qingyan, Guizhou Province, south-western China] (early Late Anisian, Middle Triassic) <HT: GPIMUM B3A-6.E7.F30-1; PT: GPIMUM B3A-6.E7.F30-2, B3A-6.E7.F53-1, B3A-6.E7.F60-1, B3A-6.E7.F60-2, B3A-6.E7.F61-1>.
- "*C.*" *mafengpoensis* STILLER, 2001: 546; pl. 2, figs 16-19. [Mafengpo Member, Xiaoshan Subformation, Qingyan Formation, Mafengpo, NNE of Qingyan, Guizhou Province,

south-western China] (middle Late Anisian, Middle Triassic) <HT: GPIMUM B3A-6.E8.F92-1>.

C. palaformis MIHÁLY, 1981: 307; pl. 3: fig. 4. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI T. 5496> {based on isolated primary spines}

C. serradentata MIHÁLY, 1981: 309; pl. 6: figs 1-2. [Aggtelek, Northern Hungary] (Anisian [Pelsonian-Upper Illyrian]) <HT: MÁFI T. 5285> {based on isolated primary spines}

“*C.*” *venustespinoza* STILLER, 2001: 538-539; text-figs 6a-f. [Leidapo Member, Yuqing Subformation, Qingyan Formation, Wachangpo, and Bangtoupou, NNE of Qingyan, Guizhou Province, south-western China] (early Late Anisian, Middle Triassic) <HT: GPIMUM B3A-6.E4.F83-1; PT: GPIMUM B3A-6.E3.F43-1, B3A-6.E3.F43-2>.

“*C.*” *wachangpoensis* STILLER, 2001: 536-538; pl. 1, figs 11-16. [Leidapo Member, Yuqing Subformation, Qingyan Formation, Wachangpo, and Bangtoupou, NNE of Qingyan, Guizhou Province, south-western China] (early Late Anisian, Middle Triassic) <HT: GPIMUM B3A-6.E3.F83-1; PT: GPIMUM B3A-6.E3.F40-1>.

Genus *Cyathocidaris* LAMBERT, 1910

Miocene

C. avenionensis nerthensis PHILIPPE, 1989: 27. [Rhône Basin, France] (Miocene) <not given> {no description, figures or repository are given}.

C. avenionensis nerthensis PHILIPPE, 1998: 42-44; pl. 4, figs 2-7. [Anse du Petit-Nid, à Sauset-les-Pins, Bouches-du-Rhône, France] (Aquitanian) <HT: ML 53127; PT: ML 53128-31>.

Genus *Gymnocidaris* A. AGASSIZ, 1863

In the Treatise (p. U333) this genus is considered a synonym of *Eucidaris* POMEL, 1883.

Jurassic

G. boutini VADET & NICOLLEAU, 2002: 4-7; 3 figs [calcaire à Nérinées; vicinity of Angoulême, Charente, France] (Kimmeridgian) <HT: Boutin coll. 1459; PT: Coutin coll. 1460, Nicollet coll. 7710, Vadet coll. 6224, 6235-6237>.

Genus *Leurocidaris* KIER, 1977

Leurocidaris KIER, 1977: 20. Type-species: *Cidaris montanaro* ZARDINI, 1973. [St. Casian Beds, Italy] (Early Karnian, Late Triassic).

Genus *Lissocidaris* MORTENSEN, 1939

Recent

L. xanthe COPPARD & VAN NOORDENBURG, 2007: 55-63; figs 1A-C, 2A-I, 4Ai-Axii, 5Ai-Aii, 6Ai-Axii. [near Mactan Island, Philippines (depth 150-250 m)] (Recent) <HT: NHM 2007.4; PT: NHM 2007.5>.

Genus *Palmerius* CUTRESS, 1980

Palmerius CUTRESS, 1980: 63-64. Type-species: *Palmerius roberti* CUTRESS, 1980.

Eocene

P. roberti CUTRESS, 1980: 64-66; pl. 4, figs 1-4. [Jabaco Fm., Bermúdez 337, Palmer 52, Palmer 1102 and Palmer 1640 localities, Cuba] (Lowermost Upper Eocene) <HT: SUPTC 10242a; PT: PRI 29632, 29760, LSJU 53244; SUPTC 10242b-h, 10243>.

Genus *Paurocidaris* KIER, 1977

Paurocidaris KIER, 1977: 18. Type-species: *Cidaris rinbianchi* ZARDINI, 1973. [St. Casian Beds, Italy] (Early Karnian, Late Triassic) {placed into the family Triadocidariidae by SMITH (1994: 187)}.

Genus *Plegiocidaris* Pomel, 1883

Middle Jurassic

P. ? gemmata GERASIMOV, 1955: 17-18; pl. 5: figs 11a, б, 12a, б, 13a, б, 14a, б. [Open pit mine near Gzhel' and Amerevo, Klyazme River, Moskva Oblast, USSR] (Middle Callovian) <HT: GMM 860>.

Genus *Stylocidaris* MORTENSEN, 1909

Recent

S. ryukyuensis SHIGEL, 1975a: 321-327; figs 1-19. [East China Sea, off Ryukyu Islands, Soyo-Maru Stat. 93 (27°52'6" N, 125°34'8" E), depth 109 m] (Recent) <HT: MMBS Echi 1011; PT: MMBS Echi 1012>.

Genus *Tretocidaris* MORTENSEN, 1903

Miocene

T. anguillensis CUTRESS, 1980: 59-63; pl. 2, figs 9-12. [Anguilla Fm., Cleve locality (a), Anguilla] (Upper Lower Miocene) <HT: USNM 115399a; PT: USNM 115399b-e>.

Genus *Triassicidaris* SMITH, 1994

Triassicidaris SMITH, 1994: 184-186. Type-species: *Triassicidaris peruviansis* SMITH, 1994.

Triassic

T. peruviansis SMITH, 1994: 186; text-fig. 4; pl. 1, figs 1-8. [Huaninchocha and Huaricolca, Peru] (Norian) <USNM 398514; PT: USNM 398506, 398515-398517, 465250>.

Genus *Zardinechinus* KIER, 1977

Zardinechinus KIER, 1977: 16. Type-species: *Cidaris lancedelli* ZARDINI, 1973. [St. Cassian Beds, Italy] (Early Karnian, Late Triassic) {placed into the family Triadocidaridae by SMITH (1994: 187)}.

Triassic

Z. giulini KIER, 1984a: 11-12; pl. 12, figs 2-6; pl. 13, figs 1-4. [St. Cassian Beds, Stolla, St. Cassiano and Miurina, Italy] (Early Karnian, Late Triassic) <HT: MSNV 10088; PT: MSNV 10089 and one unnumbered specimen in a private collection>.

Z. pulchellus SMITH, 1994: 192-193; text-fig. 5; pl. 4, figs 1-6; pl. 5, figs 1-8. [Huaninchocha and Vinchuscancha, Peru] (Norian) <HT: USNM 465261; PT: USNM 398501, 398509, 398510>.

Z. regularis SMITH, 1994: 191; text-fig. 5; pl. 3, figs 1-10; pl. 4, figs 7-10. [Huaricolca, near Tarma and Huaninchocha, Peru] (?Norian) <HT: USNM 398503; PT: USNM 398504, 398505, 398513, 465255>.

Z. stanleyi SMITH, 1994: 187-190; text-fig. 5; pl. 2, figs 5-11. [Huaricolca, near Tarma, Peru] (Norian) <HT: USNM 398511; PT: USNM 398512>.

Tribe Cidarini GRAY, 1825

Subtribe Phyllacanthina SMITH & WRIGHT, 1989

Phyllacanthina SMITH & WRIGHT, 1989: 88. Type-genus: *Phyllacanthus* BRANDT, 1835. [cosmopolitan] (Albian to Recent) {genera included: *Acanthocidaris* MORTENSEN, 1903; *Chondrocidaris* A. AGASSIZ, 1863; *Eucidaris* POMEL, 1883; *Prionocidaris* A. AGASSIZ, 1863}.

Family Psychocidaridae IKEDA, 1936

Genus *Levicidaris* KIER, 1977

Levicidaris KIER, 1977: 21. Type-species: *Levicidaris zardinia* KIER, 1977. {placed into the family Triadocidaridae by SMITH (1994: 187)}.

Triassic

L. furlani KIER, 1984a: 10-11; pl. 10, figs 1-6; pl. 11, figs 1-4. [St. Cassian Beds, Stolla, Alpe di Specie and Miurina, Italy] (Early Karnian, Late Triassic) <HT: MSNV 10086; PT: MCA 5715>.

L. pfafferi KIER, 1984a: 11; pl. 11, figs 5-7; pl. 12, fig. 1. [St. Cassian Beds, Miurina, Italy] (Early Karnian, Late Triassic) <HT: Pfaffner coll. (private) 164>.

L. zardinia KIER, 1977: 21-22; pl. 14, figs 1-8; pl. 15, figs 1-5. [St. Cassian Beds, Alpe di Specie, Miurina, Bao Staolin and Tamarin, Italy] (Early Karnian, Late Triassic) <HT: MCA 148-S-So.-T; PT: MCA 147-S-L, 151-M-Z>.

Genus *Megaporocidaris* KIER, 1977

Megaporocidaris KIER, 1977: 22-23. Type-species: *Megaporocidaris mariana* KIER, 1977. {placed into the family Triadocidaridae by SMITH (1994: 187)}.

Triassic

M. mariana KIER, 1977: 23-24; pl. 15, figs 6-8; pl. 16, figs 1-10. [St. Cassian Beds, Alpe di Specie and Miurina, Italy] (Early Karnian, Late Triassic) <HT: MCA 2728-S-L; PT: MCA 176-S-Z, 2729-S-L>.

Genus *Reboulicidaris* VADET & NICOLLEAU, 2005

Reboulicidaris VADET & NICOLLEAU, 2005: 7. Type-species: *Reboulicidaris rebouli* VADET & NICOLLEAU, 2005. [Morocco] (Jurassic).

Early Jurassic

R. rebouli VADET & NICOLLEAU, 2005: 7-9; illustr. on p. 8. [Region of Rich, Morocco] (Late Domerian (= Late Pliensbachian)) <HT: MNHN J00951 (Lambert coll'n); PT: Philippe Nicolleau coll'n 8575, 8929; Roland Reboul coll'n 0480; Alain Vadet coll'n V6498, V6499-1 to -5, V6728, V6745, V6912-V6916, V6720-V6724, V6763-6766, V6502(1-14), V6753-1 to -7>.

Genus *Tylocidaris* POMEL, 1883

Eocene

T. bermudezi CUTRESS, 1980: 134-137; pl. 14, figs 1-4. [Loma Candela Fm. and Jabaco Fm., Cuba] (Middle to Upper Eocene) <HT: SUPTC 10252a; PT: ANSP 16682, LSJU 53250-58, PRI 29700-11, SUPTC 10252b-r, 10253>.

Maastrichtian

T. hemmoorensis SALAH & SCHMID, 1982: 187-188; pl. 7, figs 1-4. [Hemmoor/Niederelbe, Northern Germany] (*junior*-zone, Late Maastrichtian) <HT: SNLFB kma 108>.

T. (T.) inexpectata JAGT & VAN DER HAM, 1995: 234-237; fig. 2. [Meerssen Mb., Maastricht Fm., ENCI NV quarry, Maastricht, the Netherlands] (late Late Maastrichtian) <HT: NHMM 1993149>.

Early Cretaceous

T. safiense VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007: 7-8; pl. 1: figs 1-7. [Falaise de Borj Nador, N of Safi, Morocco] (Late Berriasian to Early Valanginian, Early Cretaceous) <Type series: Maignac coll. 1 to 7, Nicolleau coll. 5198, 7193 to 7197, Reboul coll. 42(a-b), Vadet coll. V7022 to V7024, 7440 to 7443, 7476 to 7479, 7479 to 7781; spines: Nicolleau coll. 5185, 7203, 8588, Reboul coll. 42(1-4b), Vadet coll. V7445(1-11)> {specimen Nicolleau 5185 is listed also in the type series of *Abelleicidaris borjnadorensis* VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007}.

Subgenus *Oedematocidaris* SMITH & WRIGHT, 1989

Oedematocidaris SMITH & WRIGHT, 1989: 33-34. Type-species: *Cidaris asperula* RÖMER, 1841. [France, Germany, Denmark and Britain] (Cenomanian to Maastrichtian) {fur-

ther species included: *Cidaris pleracantha* AGASSIZ & DESOR, 1846; *Cidaris baltica* SCHLÜTER, 1892}.

Subgenus *Sardocidaris* LAMBERT, 1907

Late Cretaceous

T. (S.) trempinus GALLEMÍ & SMITH, in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 91; pl. 1, fig. 2. [Trempe Basin and Santander, Spain] (Upper Campanian-Maastrichtian) <HT: PUAB-4321; PT: PUAB-4055, PUAB-43626-32>.

Family Diplocidaridae GREGORY, 1900

Genus *Smithicidaris* VADET, 1991

Smithicidaris VADET, 1991: 156. Type-species: *Diplocidaris jacquemeti* LAMBERT, 1910. (Jurassic).

Genus *Rolliericidaris* VADET, 1991

Rolliericidaris VADET, 1991: 160-161. Type-species: *Cidaris etalloni* DESOR, in DESOR & DE LORIO, 1869, by subsequent designation of VADET et al. (2002: 12). [Europe] (Jurassic) {originally two species were mentioned, but none selected as type: *Cidaris etalloni* DESOR and *Cidaris alternans* QUENSTEDT}.

Family Nenoticidaridae VADET, 1988

Nenoticidaridae VADET, 1988: 107. Type-genus: *Nenoticidaris* VADET, 1988. {other genera included: *Paracidaris* POMEL, 1883; *Plegiocidaris* POMEL, 1883; VADET (1991: 97) included the following further genera: *Abelleicidaris* VADET, 1991; *Couvelardicidaris* VADET, 1991; *Nudicidaris* VADET, 1991; *Philicidaris* VADET, 1991; *Romanocidaris* VADET, 1991}.

Genus *Nenoticidaris* VADET, 1988

Nenoticidaris VADET, 1988: 107-108. Type-species: *Cidaris parandieri* AGASSIZ, 1840. (Liassic and Dogger) {other species included: *Cidarites blumenbachii* MÜNSTER, in GOLDFUSS, 1836; *Cidaris moeschi* DE LORIO, 1869; *Cidaris smithii* WRIGHT, 1857; VA-

DET (1991: 100-104) included further species: *Plegiocidaris pichaueriensis* LAMBERT, in LAMBERT & THIÉRY, 1924; *Cidaris zschokkei* COTTEAU, 1875 ex DESOR; *Cidaris desori* COTTEAU, 1875}.

Genus *Abelleicidaris* VADET, 1991

Abelleicidaris VADET, 1991: 109-110. Type-species: *Cidaris bradfordensis* WRIGHT, 1856. [Europe] (Jurassic-Early Cretaceous) {other species included: *Cidaris saemanni* COTTEAU, 1857}.

Early Cretaceous

A. borjnadorensis VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007: 9-10; pl. 3: figs 1-6. [Falaise de Borj Nador, N of Safi, Morocco] (Late Berriasian, Early Cretaceous) <Type series: Maignac coll. 15 to 22, Nicolleau coll. 5185, 7204, 7205, 7635, Reboul coll. 570, Vadet coll. V6376 to 6379, 7446; spines: Maignac coll. 23 to 25, Nicolleau coll. 2027, 7365, Reboul coll. 570(1-5), Vadet coll. V7477(1-7)> {specimen Nicolleau 5185 is listed also in the type series of *Tylocidaris safiense* VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007}.

Genus *Couvelardicidaris* VADET, 1991

Couvelardicidaris VADET, 1991: 97-98. Type-species: *Cidaris moorei* WRIGHT, 1956. [Europe] (Jurassic).

Liassic

C. couvelardi VADET, 1991: 99-101; pl. 2, fig. 11. [May-sur-Orne, Normandie, France (type locality); Auxois, France; Somerset, England] (Hettngian-Lower Toarcian (type stratum)) <ST: C-EM 12491 (1-5b, 8-10, 14) (tests) C-EM 12489 (4-17-33-38-39-45) (spines)>.

Genus *Nudicidaris* VADET, 1991

Nudicidaris VADET, 1991: 113-114. Type-species: *Cidaris sublaevis* COTTEAU, 1875. [Europe] (Jurassic) {other species included: *Cidarites elegans* MÜNSTER, in GOLDFUSS, 1826}.

Genus *Philicidaris* VADET, 1991

Philicidaris VADET, 1991: 104-105. Type-species: *Rhabdocidaris blainvillei* DESOR ex DESMAREST, 1858. [Europe] (Jurassic) {other species included: *Cidaritis desnoyersi* COTTEAU, 1875}.

Genus *Remycidaris* VADET, 2003

Remycidaris VADET, 2003: 12. Type-species: *Remycidaris largiporus* VADET & REMY, 2004. [France, Germany, and Switzerland] (Callovian-Tithonian, Late Jurassic) {*Cidarites elegans* MÜNSTER, in GOLDFUSS, 1826}.

Late Jurassic

R. largiporus VADET, 2003: 11-12. [Oolithe de l'Antonnière and Calcaires Sableux de Teillé, environs de Degré, Sarthe, France] (Late Jurassic) <HT: Alain Vadet coll'n V2315; PT: Alain Vadet coll'n V2316-1636, 3102, 3592, 3593, 3594, 4439>.

Genus *Romanocidaris* VADET, 1991

Romanocidaris VADET, 1991: 117. Type-species: *Romanocidaris radeti* VADET, 1991.

Jurassic

R. dietli VADET, 1991: 119-120; pl. 5, figs 5-6. [Laufen, Germany] (Lower Bajocian) <HT: SMNS 62234>.

R. pseudofilograna NICOLLEAU & VADET, 1995: 70-71; 1 fig. on p. 70, 1 fig. on p. 71; pl. 32: figs 3A, B. [Marnes à spongiares, environs de Niort, Poitou, France] (*luciaeformis* Subzone, *transvesarium* Zone, Middle Oxfordian, Jurassic) <HT: Rigollet coll. no. 79.52; PT: Alain Vadet coll. 2721, 2722, Philippe Nicolleau coll. no. 2630>.

R. radeti VADET, 1991: 118-119; pl. 5, figs 3-4. [Marnes de Dives entre Houlgate et Villers sur-mer, Normandie, France] (zone à Lamberti, Upper Callovian) <HT: V 1906(1-2)> {other species included: *Cidarites ornatus* Quenstedt, 1858}.

Family Polycidaridae VADET, 1988

Polycidaridae VADET, 1988: 103-104. Type-genus: *Polycidaris* QUENSTEDT, 1858. [Europe] (Jurassic) {other genera included: *Procidaris* POMEL, 1883; later VADET (1991: 122-123) included further genera: *Anisocidaris* THIÉRY, 1928; *Panniericidaris* VADET, 1991; *Zbindenicidaris* VADET, 1991}.

Genus *Panniericidaris* VADET, 1991

Panniericidaris VADET, 1991: 129-130. Type-species: *Cidaris moraldina* COTTEAU, 1849. [Europe] (Early Jurassic) {other species included: *Miocidaris lorioli* LAMBERT & THIÉRY, 1909}.

Genus *Zbindenicidaris* VADET, 1991

Zbindenicidaris VADET, 1991: 133. Type-species: *Cidaris toucasi* COTTEAU, 1875. [Europe] (Late Triassic-Middle Jurassic) {other species included: *Rhabdocidaris varusensis* COTTEAU, 1875}.

Family Rhabdocidaridae LAMBERT, 1900

This family is not used in the Treatise, VADET (1988: 131-132) reinstalled it and placed the genera *Rhabdocidaris* DESOR, 1856 and *Dickesicidaris* VADET, 1988 into it. VADET (1991: 142 ff.) included further genera: *Callaudicidaris* VADET, 1991; *Guttonicidaris* VADET, 1991 and *Laurenticidaris* VADET, 1991}.

Genus *Callaudicidaris* VADET, 1991

Callaudicidaris VADET, 1991: 146-147. Type-species: *Cidaris fowleri* WRIGHT, 1851. [Europe] (Jurassic).

Genus *Dickesicidaris* VADET, 1988

Dickesicidaris VADET, 1988: 136-137. Type-species: *Cidarites maximus* MÜNSTER, 1826. [Europe] (Jurassic).

Genus *Guttonicidaris* VADET, 1991

Guttonicidaris VADET, 1991: 151-152. Type-species: *Rhabdocidaris bigoti* MERCIER, 1930. [Europe] (Jurassic).

Genus *Laurenticidaris* VADET, 1991

Laurenticidaris VADET, 1991: 143-144. Type-species: *Cidaris major* COTTEAU, 1875. [Europe] (Jurassic) {other species included: *Rhabdocidaris impar* DUMORTIER, 1875}.

Genus *Rhabdocidaris* DESOR, 1856

Late Jurassic

- R. reginae* BAUMEISTER, 1999: 320-321; figs 1, 3-5. [Günsberg Beds, Vigier company quarry, Pery/La Reuchenette, Switzerland] (Oxfordian) <HT: Promusée Paléontologique, Glovelier, Switzerland, No. 3018> {based on a single specimen}.
- R. schneideri* LAMBERT, 1926: 761-763; pl. 29: fig. 2. [Crenularis-Schichten, Mellikon, Switzerland] (Séquanien inférieur, Late Jurassic) <repository unknown, formerly coll. G. Schneider> {based on a single, apparently well preserved partial corona segment}.

Middle Jurassic

- R. lahuseni* GERASIMOV, 1955: 20; pl. 6: fig. 22. [Leninskie Mt., near Moskva, D'yakoo, Moskva Oblast; Ogarkovo, Unzhe River, Ivanovskog Oblast, USSR] (Middle Callovian) <n/a> {nomen novum pro *Rhabdocidaris remus* LAGOOZEI, 1895: p. 173, fig. 236, non DESOR, non TRAUTSCHOLD}.

Family Typocidaridae VADET, 1988

Typocidaridae VADET, 1988: 130. Type-genus: *Typocidaris* POMEL, 1883.

Order Roseicidaroida VADET, 1999a

Roseicidaroida VADET, 1999a: 78. {included families: Roseicidaridae VADET, 1991}.

Family Roseicidaridae VADET, 1991

Roseicidaridae VADET, 1991: 85-86. Type-genus: *Roseicidaris* VADET, 1991. [Europe] (Jurassic) {other genera included: originally: *Caenocidaris* THIÉRY, 1928; *Merocidaris* THIÉRY, 1928; later: *Leurocidaris* KIER, 1977, see VADET, 1999a}.

Genus *Roseicidaris* VADET, 1991

Roseicidaris VADET, 1991: 86. Type-species: *Cidaris morieri* COTTEAU, 1875. [Europe] (Toarcian).

Order Thielicidaroida VADET, 1999c

Thielicidaroida VADET, 1999c: 69. {included families: Thielicidaridae VADET, 1999c}.

Family Thielicidaridae VADET, 1999c

Thielicidaridae VADET, 1999c: 69. Type-genus: *Thielicidaris* VADET, 1999c.

Genus *Thielicidaris* VADET, 1999c

Thielicidaris VADET, 1999c: 69. Type-species: *Thielicidaris thieli* VADET, 1999c. [Europe] (Late Triassic)

Triassic

T. thieli VADET, 1999c: 69-70; figs 81-85. [Couches de Saint Cassian, Italy] (Carnian) <HT: Thiel K-1>.

Subclass Euechinoidea BRONN, 1860

Order Triadotiaroida HAGDORN, 1995

Triadotiaroida HAGDORN, 1995: 249. {included families: Triadotiaridae HAGDORN, 1995}.

Family Triadotiaridae HAGDORN, 1995

Triadotiaridae HAGDORN, 1995: 249. Type-genus: *Triadotiaris* HAGDORN, 1995. [Europe] (Anisian-Ladinian) {VADET (1999c: 70) included the following genera: *Lenticidaris* KIER, 1968; *Mesodiadema* NEUMAYR, 1889; *Anaulcidaris* ZITTEL, 1879}.

Genus *Triadotiaris* HAGDORN, 1995

Triadotiaris HAGDORN, 1995: 250. Type-species: *Cidarites grandaevus* v. ALBERTI, 1834. [Central Europe] (Anisian-Ladinian).

Order Serpianotiaroidea HAGDORN, 1995

Serpianotiaroidea HAGDORN, 1995: 258. {included family: Serpianotiaridae HAGDORN, 1995}.

Family Serpianotiaridae HAGDORN, 1995

Serpianotiaridae HAGDORN, 1995: 258. Type-genus: *Serpianotiaris* JEANNET, 1933. [Central Europe] (Anisian-Karnian).

Infraclass Acroechinoidea SMITH, 1981

Family Nicolleumatidae VADET, 1999a

Nicolleumatidae VADET, 1999a: 75. Type-genus: *Nicolleuma* VADET, 1999a.

Genus *Nicolleuma* VADET, 1999a

Nicolleuma VADET, 1999a: 75-76. Type-species: *Pseudodiadema collenoti* COTTEAU, 1882. [Europe] (Jurassic) {other species included: *Eodiadema lacostei* LAMBERT, 1933; *E. aff. minutum* SMITH, 1981; *Acrosalenia chartroni* LAMBERT, 1904}.

Order Glyptocidaroida JENSEN, 1982

Glyptocidaroida JENSEN, 1982: 90. {included families: Glyptocidaridae JENSEN, 1982}.

Family Glyptocidaridae JENSEN, 1982

Glyptocidaridae JENSEN, 1982: 90. Type-genus: *Glyptocidaris* A. AGASSIZ, 1853.

Superorder Echinothuriacea JENSEN, 1982

Echinothuriacea JENSEN, 1982: 92. {included orders: Echinothurioida CLAUS, 1880}.

Superorder Diadematacea DUNCAN, 1889

Order Echinothurioida CLAUS, 1880

JENSEN (1982: 92) placed this order into a new own super: Echinothuriacea JENSEN, 1982.

Family Phormosomatidae MORTENSEN, 1934

This family is not used in the *Treatise*, instead it runs as Subfamily Phormosomatinae MORTENSEN, 1934. SMITH & WRIGHT (1990: 104) proposed a new classification for the Echinothurioida, where this taxon is used at family level.

Subfamily Kamptosomatinae SMITH & WRIGHT, 1990

Kamptosomatinae SMITH & WRIGHT, 1990: 104. Type-genus: *Kamptosoma* MORTENSEN, 1903.

Family Echinothuriidae WYVILLE THOMPSON, 1872

Subfamily Echinothuriinae WYVILLE THOMPSON, 1872

Genus *Asthenosoma* GRUBE, 1868

Recent

A. marisrubis WEINBERG & DE RIDDER, 1998: 32-35; fig. 3A-C, 4A-C, 5A-E, 6a-g, 7a-g. [northern Red Sea] (Recent) <HT: ZMA V.ech.E 9365; PT: ZMA V.ech.E 9366; ZM-TAU EC 25123, NS 5091; HUJ ECHI 343; MNHN 6-9. EC ES 8376>.

Genus *Hapalosoma* MORTENSEN, 1903

Recent

H. pulchrum ROWE, 1989: 263-265; figs 2A-B, 3A-B. [off Norfolk Island] (Recent) <HT: AM J21699>.

Genus *Kamptosoma* MORTENSEN, 1903

Recent

K. abyssale MIRONOV, 1971: 321-322; figs 2, 3а-д, 4а-г. [R/V Vityaz Stat. 2074 (42°32' N, 150°41'05" E, depth 5140 m), 2119 (46°08' N, 155°16' E, depth 5070-5090 m), 3162 (43°15' N, 157°48' E, depth 5502 m), 3206 (30°53' N, 153°09' E, depth 5988-5998 m), 3575 (38°02' N, 146°33' E, depth 5495 m), 3825 (25°32' S, 175°27' W, depth 5123-5329 m), 4655 (16°07' S, 53°35' E, depth 4675 m), 4911 (1°55' S, 83°05' E, depth 4794-4809 m), 5065 (16°23' N, 146°36' W, depth 5363-5570 m), 5315 (8°22' S, 80°35' E, depth 5100-5120 m), 5322 (18°57' S, 56°05' E, depth 4374 m), 5609 (46°06' N, 153°18' E, depth 6090-6235 m), 5620 (44°48' N, 156°33' E, depth 5005-5045 m), 5623 (46°26' N, 154°59' E, depth 4995-5045 m), 5625 (45°28' N, 153°46' E, depth 6205-6215 m), 6143 (51°40' N, 163°00' W, depth 4860 m), Pacific Ocean and Indian Ocean] (Recent) <HT: IOANSSR no specimen no. given (from R/V Vityaz Stat. 5609)>.

Genus *Retzneiosoma* KROH, 2005

Retzneiosoma KROH, 2005: 14. Type-species: *Retzneiosoma jasenecki* KROH, 2005. [Europe] (Miocene).

Miocene

R. jasenecki KROH, 2005: 16-18; fig. 10; pl. 9, figs 1-12; pl. 10, figs 1-8. [grey marls overlying the Weissenegg Formation, Retznei (Lafarge quarry (formerly Perlmoser)), Styria, Austria] (Early Badenian [= Langhian], Middle Miocene) <HT: NHMW 2003z0071/0001>.

Genus *Sperosoma* KOEHLER, 1897

Recent

S. nudum SHIGEL, 1977a: 71-76; figs 1-16. [Soyo-Marú Stat. B2 (34°05'2" N, 140°00'7" E, off Miyake Island, depth 1410-1450 m), Stat. B3 (32°32'4" N, 140°18'7" E, off Aogashima Island, depth 2080 m) and Stat. 51 (34°3'5" N, 139°39'5" E, off Izu-Os-

hima Island, depth 1570 m), off central Japan, North Pacific Ocean] (Recent) <HT: MMBS Echi 1211; PT: MMBS Echi 1212-1216>.

Subfamily Hygrosomatinae SMITH & WRIGHT, 1990

Hygrosomatinae SMITH & WRIGHT, 1990: 104. Type-genus: *Hygrosoma* MORTENSEN, 1903.

Subfamily Paraphormosomatinae SMITH & WRIGHT, 1990

Paraphormosomatinae SMITH & WRIGHT, 1990: 104. Type-genus: *Paraphormosoma* MORTENSEN, 1934.

Subfamily Pelanodiademinae HESS, 1972

Pelanodiademinae HESS, 1972: 42. Type-genus: *Pelanodiadema* HESS, 1972.

Genus *Pelanodiadema* HESS, 1972

Pelanodiadema HESS, 1972: 42. Type-species: *Pelanodiadema oolithicum* HESS, 1972. [Central Europe] (Middle Jurassic).

Middle Jurassic

P. oolithicum HESS, 1972: 42-50; figs 48-63; pl. 16, figs 1-3; pl. 17, figs 1-3; pl. 18, fig. 3. [Upper Hauptrogenstein, Schinznach-Dorf, Switzerland] (Upper Bajocian) <HT: Hess coll. E300>.

Subfamily Sperosomatinae SMITH & WRIGHT, 1990

Sperosomatinae SMITH & WRIGHT, 1990: 104. Type-genus: *Sperosoma* KOEHLER, 1897. {other genera included: *Tromikosoma* MORTENSEN, 1903}.

Order Diadematoida DUNCAN, 1889

Family Diadematidae GRAY, 1855

Genus *Centrostephanus* PETERS, 1855

Recent

C. sylviae FELL, 1975: 182-190; figs 1, 2b-f, 4, 5. [San Felix Island, Chile and between Point Island and Padre Bay, west end of Mas a Tierra, Juan Fernández Islands, South Pacific Ocean; depth 9-70 m] (Recent) <HT: USNM E11379; PT: MCZ 8778-8780, 8783, NMNZ ECH.1500, National Museum Chile (Santiago), USNM E11380>.

Genus *Diadema* GRAY, 1825

Recent

D. setosum forma *depressa* DOLLFUS & ROMAN, 1981: 38-39; pl. 5, figs 4-6; pl. 6, figs 1-3. [Red Sea] (Recent) <HT: not given>.

Genus *Palaeodiadema* POMEL, 1887

Oligocene

P. reingardae KUTSCHER, 1985a: 5-6; pl. 1, fig. 5; pl. 2, figs 1-4. [Magdeburg, Germany] (Middle Oligocene) <HT: no specimen no. given>.

Family Aspidodiadematidae DUNCAN, 1889

Genus *Aspidodiadema* A. AGASSIZ, 1878

Recent

- A. intermedium* SHIGEI, 1977b: 79-84; figs 1-10. [East China Sea, off Shimo-Koshiki Island, 31°35' N, 129°42' E, depth 380 m] (Recent) <HT: MMBS Echi 1201>.
- A. montanum* MIRONOV, 1981: 132-133; figs 1.1, 1.3, 2.1, 2.3, 2.10. [Marcus Necker Ridge, North Pacific Ocean; 1270-1950 m depth] (Recent) <HT: IONASSR XY-65-17>.
- A. sinuosum* MIRONOV, 1981: 133-135; figs 1.2, 1.4, 2.2, 2.5, 2.8, 2.12. [Marcus Necker Ridge, North Pacific Ocean; 100-1350 m depth] (Recent) <HT: IONASSR XY-65-18>.

Genus *Cherreauma* VADET, NICOLLEAU & PINEAU, 1996

Cherreauma VADET, NICOLLEAU & PINEAU, 1996: 60-61. Type-species: *C. cherreai* VADET, NICOLLEAU & PINEAU, 1996. [Europe] (Jurassic).

Middle Jurassic

C. cherreai VADET, NICOLLEAU & PINEAU, 1996: 61-63; figs 74-77; pl. 15, figs 1a-d. [Oolithe de l'Antonnière, environs du Mans, Sarthe, France] (Callovian) <HT: Cherreau coll.; PT: Thiel coll., Dudicourt coll.> {no specimen numbers given}.

Genus *Culozoma* VADET & SLOWIK, 2001

Culozoma VADET & SLOWIK, 2001: 35. Type-species: *Culozoma baroini* VADET & SLOWIK, 2001. [Europe] (Jurassic).

Bajocian

C. baroini VADET & SLOWIK, 2001: 35-38; 2 text-figs (not numbered); pl. 6, 6 figs (not numbered). [Culoz dans l'Ain, France] (Lower? Bajocian) <HT: Vadet coll. V6006; PT: Vadet coll. V6120>.

Genus *Gymnotiara* POMEL, 1883

Middle Jurassic

G. kuhni HESS, 1971: 622-626; figs 11-14. [Anceps-Athleta-Beds, Schinboden ob Ramiswil, Solothurn, Switzerland] (Callovian, Jurassic) <HT: Hans HESS coll. E269>.

Family uncertain

Genus *Brangema* NICOLLEAU & VADET, 1995

Brangema NICOLLEAU & VADET, 1995: 77. Type-species: *Brangema brangeri* NICOLLEAU & VADET, 1995. [Europe] (Jurassic).

Jurassic

B. brangeri NICOLLEAU & VADET, 1995: 77-78; 2 figs on p. 77; pl. 30: figs 4A, B. [Marnes à spongiares, environs de Niort, Poitou, France] (stenocycloides Subzone, bifucatus Zone, Late Oxfordian, Late Jurassic) <HT: Philippe Nicolleau coll. no. 2532>.

Family Eodiadematidae SMITH, 1984

Eodiadematidae SMITH, 1984: 171, 172-173. Type-genus: *Eodiadema* DUNCAN, 1889.

Eodiadematida KINMAN, 1994: 31, 64. Type-genus: *Eodiadema* DUNCAN, 1889. (Jurassic)
{objective junior homonym of Eodiadematidae SMITH, 1984}.

Order Micropygoida JENSEN, 1982

Micropygoida JENSEN, 1982: 92. {included families: Micropygidae MORTENSEN, 1904}.

Superorder Pedinacea JENSEN, 1982

Pedinacea JENSEN, 1982: 93. {included orders Pedinoida MORTENSEN, 1939, Pygasteroida
DURHAM & MELVILLE, 1957}.

Order Pedinoida MORTENSEN, 1939

Family Pedinidae POMEL, 1883

Genus *Caenopedina* A. AGASSIZ, 1869

Recent

C. alanbakeri ROWE, 1989: 265-268; figs 4A-B, 5A-D. [off Norfolk Island] (Recent)
<HT: AM J21700>.

C. novaezealandiae PAWSON, 1964: 64-66; figs 1-5; plate 1. [16 miles south-east of Mayor
Island, Bay of Plenty, 180-240 fathoms (4/8/1963, prawn net), Australasia] (Recent)
<HT: Dominion Museum, Wellington, without no.>

C. otagoensis MCKNIGHT, 1968: 90-94; figs 1-3. [Stat. E 399 (46°00' S, 171°33' E, depth
1214-1222 m), E 427 (44°54' S, 172°54' E, depth 1207-1233 m), F 751 (45°23' S,
175°29' E, depth 1227-1258 m), Tasman Sea] (Recent) <HT: NZOI 22 (Stat. E 427,
female); PT: NZOI P 38, P 40 (2 females, Stat. E 427), P 39, P 41 (2 males, Stat. E
427)>.

Genus *Echinopedina* COTTEAU, 1866

Eocene

E. paucituberculata LEWIS, 1989: 9-13; figs 2 a-v, 3; pl. 1: figs 3-7. [Barton-on-Sea, Hampshire, England] (Bartonian) <HT: BMNH E 76581; PT: BMNH E 76425, E 76573-80, E 79701, E 79821>.

Genus *Hemipedina* WRIGHT, 1855

Triassic

H. hudsoni KIER, 1977: 30-32; fig. 11a-c; pl. 21, figs 5-9. [Elphinstone Group, probably Sumatra Fm., Wadi Milaha, Trucial Coast, Arabia] (Norian, Late Triassic) <HT: BMNH E-76154>.

Genus *Leiopedina* COTTEAU, 1866

Eocene

L. molinai SILLERO, 2002: 13-16; figs 1-4. [Carretera de Busot a Jijona, Busot, Province Alicante, Spain] (Middle Eocene) <HT: CGCPE-CI966; PT: CGCPE-CI967> {figured and re-described in LÓPEZ & SILLERO (2006: 153, fig. 126)}.

Genus *Pedina* L. AGASSIZ, 1838

Middle Jurassic

P. davoustiana COTTEAU, 1856 var. *mercieri* VADET, NICOLLEAU & PINEAU, 1996: 47-48; figs 53-54; pl. 1, figs 2a-c; pl. 11, figs 1a-2c. [Oolithe de l'Antonnière, Sarthe, France] (Callovian) <HT: V 4523; Type-series: V 3167-3178, 4523, Pineau coll. 1-3, 14>.

Genus *Thieulinipedina* VADET, NICOLLEAU & PINEAU, 1996

Thieulinipedina VADET, NICOLLEAU & PINEAU, 1996: 43. Type-species: *Pedina antiqua* COTTEAU, 1883. [Sarthe, France and Portugal] (Callovian) {other species included: *Pedina resecta* (LAMBERT, 1916)}.

Family uncertain

Genus *Farquharsonia* CURRIE, 1927

VADET, NICOLLEAU & PINEAU (1996: 57-59) place this genus into the family Aspidodiademataidae DUNCAN, 1889, within the order Diadematoidea DUNCAN, 1889, whereas KIER (1972a: 19) placed it into the family Diademataidae GRAY, 1855.

Middle Jurassic

F. crenulata KIER, 1972a: 23-24; fig. 7; pl. 1, figs 1-6; pl. 2, figs 1-2. [Middle Dhurma Fm., Locality KK9-43, Saudi Arabia] (Bathonian) <HT: USNM 170370>.

F. pineaui VADET, NICOLLEAU & PINEAU, 1996: 57-60; figs 69-73; pl. 14, figs 1a-3. [Oolithe de l'Antonnière, Sarthe, France] (Callovian) <HT: Pineau coll. 306; PT: Pineau coll. 305, Grignon coll.>.

Order Pygasteroidea DURHAM & MELVILLE, 1957

Family Pygasteridae LAMBERT, 1900

Genus *Jesionekechinus* VADET, 1997

Jesionekechinus VADET, 1997:128. Type-species: *Plesiechinus hawkinsi* JESIONEK, 1970. [Nevada, USA] (Sinemurian).

Genus *Plesiechinus* POMEL, 1883

Middle Jurassic

P. altus KIER, 1972a: 24-25; fig. 8; pl. 32, figs 1-4. [Middle Dhurma Fm., locality L-921, Saudi Arabia] (Bathonian) <HT: USNM 170419>.

Superorder Echinacea CLAUS, 1876

Order Acrosalenioidea VADET, 1999b

Acrosalenioidea VADET, 1999b: 78. {included families: Acrosaleniiidae GREGORY, 1900}.

Order Salenioidea DELAGE & HÉROUARD, 1903

According to SMITH & WRIGHT (1990: 117) the name *Calycina* GREGORY, 1900 has priority over *Salenioidea* DELAGE & HÉROUARD, 1903.

Family Acrosaleniidae GREGORY, 1900

Genus *Acrosalenia* L. AGASSIZ, 1840

Upper Jurassic

A. bowersi KIER, 1972a: 34-35; fig. 13; pl. 25, figs 1-5; pl. 26, figs 1-2. [Upper Dhurma Fm., locality S-1148, Saudi Arabia] (Callovian) <HT: USNM 170409; PT: USNM 170410>.

Middle Jurassic

A. arabica KIER, 1972a: 27-29; figs 9-10; pl. 28, figs 2-5; pl. 29, figs 1-4; pl. 30, figs 1-4. [Middle Dhurma Fm., localities S-1046, S-1056, S-1157, S-1160, S-1503, KK8-23, 30-35, 33-35, 34, 35-38, 37, 38.5, 40.5, 44, KK9-30-40, Saudi Arabia] (Bathonian) <HT: USNM 170412; PT: USNM 170413-170417>.

A. dhrumaensis KIER, 1972a: 29-34; figs 11-12; pl. 31, figs 1-8. [Middle Dhurma Fm., localities KK9-21-22.5, KK9-51.5, Saudi Arabia] (Bathonian) <HT: USNM 170418>.

Early Jurassic

A. marratensis KIER, 1972a: 26-27; pl. 27, figs 3-7; pl. 28, fig. 1. [Marrat Fm., localities S-1034, KK6-14, Saudi Arabia] (Toarcian) <HT: USNM 170411>.

Genus *Heterosalenia* COTTEAU, 1861

Late Jurassic

H. brocki KIER, 1972a: 36-37; fig. 14; pl. 7, figs 1-6; pl. 8, figs 1-6; pl. 9, fig. 1. [Upper Dhurma Fm., localities KK9-111, S-1167, KK9-112, Saudi Arabia] (Callovian) <HT: USNM 170378; PT: USNM 170379>.

H. ornata KIER, 1972a: 37-40; fig. 15; pl. 5, figs 2-6; pl. 6, figs 1-8. [Upper Dhurma Fm., locality KK9-112, S-1167, KK9-111, Saudi Arabia] (Callovian) <HT: USNM 170375; PT: 170376-170377>.

Middle Jurassic

H. dhrumaensis KIER, 1972a: 35-36; pl. 9, figs 2-4. [Middle Dhurma Fm., locality KK9-15, Saudi Arabia] (Bathonian) <HT: USNM 170380>.

Genus *Milnia* HAIME, 1849

In the *Treatise* the genus *Milnia* HAIME, 1849 is considered a junior synonym of *Acrosalenia* L. AGASSIZ, 1840.

Middle Jurassic

M. guttoni VADET, NICOLLEAU & PINEAU, 1996: 82-84; figs 97-100; pl. 22, figs 1a-2c. [Oolithe de l'Antonnière and Oolithe de Vivoin et l'Assise des Carreaux, Sarthe, France] (Callovian) <HT: V 4442; Type-series: V 2725, 2745-2748, 4072-4074, 4440-4450, Pineau coll. 110-111-117-118-285-361-362-363>.

Genus *Prosalenia* VADET, NICOLLEAU & PINEAU, 1996

Prosalenia VADET, NICOLLEAU & PINEAU, 1996: 84. Type-species: *Acrosalenia marcoui* COTTEAU, 1879. [Sarthe and Jura, France] (Callovian to Oxfordian).

Family Pseudosaleniidae VADET, 1999b

Pseudosaleniidae VADET, 1999b: 78. Type-genus: *Pseudosalenia* COTTEAU, 1859.

Genus *Pseudosalenia* COTTEAU, 1859

Upper Jurassic

P. malogostiana RADWAŃSKA, 1999: 309-310; pl. 11, figs 1a-f; pl. 12, figs 1a-f. [Małogoszcz Quarry, south-western margin of the Holy Cross Mountains, Central Poland] (Lower Kimmeridgian) <HT: DPUW EMa/34>.

Middle Jurassic

P. magniprocta KIER, 1972a: 40-42; fig. 16; pl. 9, figs 5-7; pl. 10, figs 1-4; pl. 11, figs 1-4. [Middle Dhurma Fm., localities KK8 30-35, 34, 35-38, KK9 30-40, S-1046, S-1160, Saudi Arabia] (Bathonian) <HT: USNM 170383; PT: USNM 170381-170382>.

Family Saleniidae L. AGASSIZ, 1838

Subfamily Saleniinae L. AGASSIZ, 1838

Genus *Salenia* GRAY, 1835Subgenus *Salenia* GRAY, 1835

Recent

- S. nudispina* MARKOV, 1988b: 568-569; figs 1a, г, з, 2a. [“Ихтиандр”- 6th cruise (“Ikhtiandr”-6th cruise) 23°25' S, 83°19' W, depth 390-560 m, East Pacific Ocean] (Recent) <HT: IOANSSR, no specimen no. given>.
- S. profundis brevispina* MARKOV, 1988a: 381-382; figs 1в, л, 2г. [“Академик Курчатов”-cruise (“Akademik Kurchatov”-cruise) Stat. 12 (0°08' N, 18°13' E, depth 2660-2800 m), “Витязь”-cruise (“Vityaz”-cruise) Stat. 7505 (34°17' N, 143°52' E, depth 1800-2400 m), “Дмитрий Менделеев”-cruise (“Dmitrig Mendelejev”-cruise) Stat. 529 (2°11' N, 102°31' E, depth 3190 m), Atlantic Ocean] (Recent) <HT: IOANSSR, no specimen no. given >.
- S. profundis intermedius* MARKOV, 1988a: 382. [“Витязь”-cruise (“Vityaz”-cruise) Stat. 8021 (34°41' N, 40°37' W, depth 3160-3080 m)] (Recent) <HT: IOANSSR, no specimen no. given>.
- S. profundis megalospina* MARKOV, 1988a: 382-383; figs 1д, н, 2у. [“Академик Курчатов”-cruise (“Akademik Kurchatov”-cruise) Stat. 23 (5°08' N, 62°18' E, depth 3040-3500 m), “Витязь”-cruise (“Vityaz”-cruise) Stat. 4634 (2°47' S, 65°43' E, depth 3530 m), “Дмитрий Менделеев”-cruise (“Dmitrig Mendelejev”-cruise) Stat. 1367 (33°63' S, 127°55' E, depth 2520 m)] (Recent) <HT: IOANSSR, no specimen no. given >.

Oligocene

- S. cascadiensis* LINDER, DURHAM & ORR, 1988: 946; figs 2.1, 2.3. [Scott Mills Fm., Abiqua Mb., Butte Creek, Marion County, Oregon, USA] (Late Oligocene) <HT: UCMP 38194; PT: UCMP 38195>.

Eocene

- S. tandoni* SRIVASTAVA, 1982: 23-25; figs 1-4. [about 2 km south-west of Guvar, Kutch, Gujarat, India] (*Nummulites beaumonti*-zone of TANDON, 1976, Middle Eocene) <HT: DGUL K500; PT: DGUL K501>.
- S. trisuranalis* LEWIS & JEFFERIES, 1980: 115-118; figs 1-4. [London Clay, Walton-on-the-Naze, Essex, England] (Lower Eocene) <HT: BMNH E76505>.

Maastrichtian

- S. microprocta* SMITH, 1995: 142; fig. 15, 16F; pl. 5: figs 13. [Jebel Huwayyah, Oman/ United Arabian Emirates Border Region] <HT: BMNH EE3657>.

S. sigillata pozaryskae GEYS & MACHALSKI, 1992: 136-138; pls. 1, figs 1a-e. [Kazimierz-on-Vistula, near Pulawy, Central Poland] (*Belemnella kazimiroviensis* zone, Uppermost Maastrichtian) <HT: ZPAL E. VI/1>.

Turonian

S. cherei GEYS, 1982: 10-11; pl. 3, figs 2-5; pl. 4, fig. 1. [Chercq (Cornet quarry), Hainaut, Belgium] (Turonian) <HT: KBIN IST 10187>.

Cenomanian

S. ammonitorum BANDEL & GEYS, 1985: 104-105; pl. 3, figs 1-6. [Salihi Fm., Wadi Salihi, near Amman, Jordan] (Cenomanian) <KBIN coll., IST 10232>.

S. (S.) baylissi SMITH & WRIGHT, 1990: 153-154; text-figs 41A, B; pl. 45, fig. 3. [Shapwick Quarry and between Beer and Branscombe, Devon, England] (*Calycoceras guerangeri* Zone, Upper Cenomanian) <HT: BMNH E82856; PT: BMNH E82873, GSM 5279>.

S. pentagonalis SHMIDT & SIMAKOV, 1953: 32-33; figs 3a-b; pl. 1: figs 15-17. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Cenomanian ?) <HT: VNIGRI 2/336>.

S. (S.) wilmingttonensis SMITH & WRIGHT, 1990: 152-153; text-figs 41C, D; pl. 46, fig. 1. [Waterworks Quarry, Wilmington, Devon, England] (*M. mantelli* Zone, *N. carcitanensis* Subzone, Basal Cenomanian) <HT: BMNH E82857>.

Aptian

S. banatica DRAGOMIR, 1997: 218-219; fig. 1, pl. 29.1, figs 3-8. [Valea Minişului Fm., Tisa valley, Moldova Novă, Banat, Romania] (Bedoulian, Early Aptian) <HT: LPB elt 138; PT: LPB elt 021 (8 specimens)>.

S. danubiana DRAGOMIR, 1997: 219; fig. 2, pl. 29.1, figs 9-14. [Valea Minişului Fm., Tisa valley, Moldova Novă, Banat, Romania] (Bedoulian, Early Aptian) <HT: LPB elt 025; PT: LPB elt 025a (7 specimens)>.

Cretaceous, undifferentiated

S. bella SZÖRÉNYI, 1955: 21-23, 163-164; fig. 6; pl. 1, figs 7-9. [marne glauconieuse, Bakonynána (carrière de la vallée Gaja), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/23>.

S. bella parva SZÖRÉNYI, 1955: 23, 164-165; pl. 1, figs 21-23. [marne glauconieuse, Pénezskút-Körisgyörpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/26>.

S. scutigera hungarica SZÖRÉNYI, 1955: 23-24, 165-166; pl. 1, figs 15-17. [calcaire à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/24>.

Subgenus *Pleurosalenia* POMEL, 1883

This taxon was considered as synonym of *Salenidia* POMEL, 1883 in the *Treatise* (p. U378).

Paleocene

S. (P.) vanbirgeleni SMITH, in SMITH & JEFFERY, 2000: 59-60; text-fig. 21D-F. [Geulhem Mb., Albertkanaal, near Kesselt, Belgium] (Early Danian) <HT: NHMM MB 432J>.

Genus *Leptosalenia* SMITH & WRIGHT, 1990

Leptosalenia SMITH & WRIGHT, 1990: 134-135. Type-species: *Salenia pretensis* DESOR, 1856. (Cretaceous) {other species included: *S. arabica* FOURTAU; *S. texana* CREDNER; *S. mexicana* SCHLÜTER; *S. volana* WHITNEY; *S. plana* FOURTAU; ?*S. humei* FOURTAU; ?*S. mathuri* CHIPLONKER}.

Aptian

L. faringdonensis SMITH & WRIGHT, 1990: 137-139; text-fig. 33A-B; pl. 37, fig. 1. [Faringdon Sponge Gravels, Faringdon, Berkshire, England] (*P. nutfieldensis* zone, Upper Aptian) <HT: BMNH E82847; PT: BMNH E79226, E78961, E82871>.

L. qishnensis NESTLER, 1998: 569-570; figs 4c-d, 5a-d. [Qishn Formation, Wadi Hajr, östl. Jawl Ba Hawa, Hadramawt, South Yemen] (Aptian, Early Cretaceous) <HT: FGWG 107/1> {based on a single specimen}.

Genus *Novasalenia* ŽITT & GEYS, 2003

Novasalenia ŽITT & GEYS, 2003: 24. Type-species: *Novasalenia predbojensis* ŽITT & GEYS, 2003. [Europe] (Late Cretaceous).

Cenomanian

N. plananyensis ŽITT & GEYS, 2003: 27-29; figs 5a-d. [Plaňany, Bohemian Cretaceous Basin, Czech Republic] <boundary interval of the *Rotalipora cushmani* and *Praeaectinocamax plenus* biozones, Late Cenomanian> (HT: NMP O 6313).

N. predbojensis ŽITT & GEYS, 2003: 25-27; figs 2a-d, 3, 4- [Předboj, Bohemian Cretaceous Basin, Czech Republic] <boundary interval of the *Rotalipora cushmani* and *Praeaectinocamax plenus* biozones, Late Cenomanian> (HT: NMP O 6312).

Genus *Perisalenia* VALETTE, 1906

Jurassic

P. hemicidaroides var. *lenoiri* VADET, 2005a: 68-69; illustr. on p. 68. [membre A, formation de Leulinghen, couches à Modioles, Boulonnais, France] (Late Bajocian-Early Bathonian) <HT: Laurent Lenoir coll'n L234b; PT: Laurent Lenoir coll'n L234a, L235, L236a-d, Alain Vadet coll'n V5276, V6441-V6444>.

Genus *Plesiosalenia* SMITH & WRIGHT, 1990

Plesiosalenia SMITH & WRIGHT, 1990: 134. Type-species: *Salenia depressa* GRAS, 1848. [Europe] (Early Cretaceous) {junior homonym of *Plesiosalenia* Valette, 1906}.

Genus *Prandinia* VADET, 2000

Prandinia VADET, 2000: 93; figs on pp. 94, 95; pl. 1 (5 figs). Type-species: *Cidarites interpunctata* QUENSTEDT, 1852. [Southern Germany] (Malm, Late Jurassic).

Genus *Salenocidaris* A. AGASSIZ, 1869

Maastrichtian

S. gallemii SMITH & JEFFERY, 2000: 65-66; text-fig. 23A-C. [Partida del Matet, Polop de la Marina, and Aspe, Alicante Province, Spain] (?Early Maastrichtian) <HT: BMNH EE6626; PT: BMNH EE6267-8, EE6785-90>.

Genus *Salenidia* POMEL, 1883

Subgenus *Salenidia* POMEL, 1883

Maastrichtian

S. sanctipetri GEYS, 1979: 303-306; figs 5.1-5.6, 6.1-6.2. [St. Pietersberg, near Maastricht, Dutch Limberg, the Netherlands] (Upper Maastrichtian) <HT: KBIN 10160>.

Subgenus *Platysalenia* SMITH & WRIGHT, 1990

Platysalenia SMITH & WRIGHT, 1990: 134, 167. Type-species: *Salenia dux* WRIGHT, 1967. [England] (Late Albian).

Tribe Salenocidarini SMITH & WRIGHT, 1990

Salenocidarini SMITH & WRIGHT, 1990: 134, 168-169. Type-genus: *Salenocidaris* AGASSIZ, 1869.

Tribe Holosaleniini SMITH & WRIGHT, 1990

Holosaleniini SMITH & WRIGHT, 1990: 134. Type-genus: *Holosalenia* SMITH & WRIGHT, 1990.

Genus *Holosalenia* SMITH & WRIGHT, 1990

Holosalenia SMITH & WRIGHT, 1990: 134. Type-species: *Salenia batnensis* COTTEAU, PERON & GAUTHIER, 1879. [Europe, Northern Africa, Middle East] (Mid to Late Cretaceous).

Early Cretaceous

H. bahiensis MANSO & SOUZA-LIMA, 2007: 30-32; figs 4A-I, 5A-C. [Germânia Member, Algodões, Boipeba 2 (UTM 8.499.000N, 508.600E), Boipeba 3 (type locality, UTM 8.499.000N, 508.600E), Boipeba 4 (UTM 8.499.050N, 508.800E), and Tassimirim 1 (UTM 8.499.000N, 509.150E), Boipeba Island, NW Velha Boipeba, Camamu Bay, and Cangaíba 1 (UTM 8.462.150N, 498.100E), and Cangaíba 2 (UTM 8.462.150N, 498.250E), Cangaíba Island, Camamu Bay, Eastern Brazil] (Late Albian) <HT: FPH-462-I (Boipeba 3); PT: FPH-449-I to FPH-456-I (Boipeba 2), FPH-460-I, FPH-461-I, FPH-463-I, FPH-464-I, FPH-467-I to FPH-503-I, FPH-507-I, FPH-508-I, MN 8251-I, MN 8252-I (Boipeba 3), FPH-868-I, FPH-869-I (Boipeba 4), FPH-889-I, FPH-890-I (Tassimirim 1), FPH-870-I, FPH-871-I, FPH-873-I to FPH-876-I (Cangaíba 1), FPH-879-I to FPH-882-I, FPH-885-I, FPH-886-I to FPH-888-I (Cangaíba 2)>.

Subfamily Hyposaleniinae MORTENSEN, 1934

Genus *Hyposalenia* DESOR, 1856.

Albian

H. radians SMITH & WRIGHT, 1990: 191-193; text-figs 62, 63; pl. 69, fig. 2. [Shenley Limestone, Double Arches pit, Shenley Hill and Arnold's pit, Billington Crossingm Leighton Buzzard, Bedfordshire, England] (*L. regularis* Subzone, Lower Albian) <HT: BMNH E83526; PT: BMNH E82866>.

Genus *Glyphopneustes* POMEL, 1869

Maastrichtian

G. hattaensis ALI, 1992a: 68-70; fig. 3a-e. [Simsima Fm., Gebel El Rowdah, Hatta area, Oman-U.A.E.] (Late Maastrichtian) <HT: MGD-AAA> {no specimen numbers given}.

Tribe Hyposaleniini SMITH & WRIGHT, 1990

Hyposaleniini SMITH & WRIGHT, 1990: 123, 174. Type-genus: *Hyposalenia* DESOR, 1856.

Tribe Goniophorini SMITH & WRIGHT, 1990

Goniophorini SMITH & WRIGHT, 1990: 123, 193. Type-genus: *Goniophorus* AGASSIZ, 1838. {other genera included: *Glyptopneustes* POMEL, 1869}.

Order Hemicidaroida BEURLEN, 1937

Family Hemicidaridae WRIGHT, 1857

Genus *Gymnotiara* POMEL, 1883

In the *Treatise* FELL & PAWSON (1966: U387) considered this genus as synonymous to *Hypodiadema* DESOR, 1858.

Middle Jurassic

G. kuhni HESS, 1971: 622-626; figs 11-14. [Anceps-Athleta-Beds, Schinboden ob Ramiswil, Solothurn, Switzerland] (Callovian, Jurassic) <HT: Hans Hess coll. E269>.

Genus *Hemicidaris* L. AGASSIZ, 1838

Cenomanian-Turonian

H. bandeli GEYS, 1989: 128; pl. 1, figs 1-4. [Atrash Fm., Wadi Qena, Eastern Desert, Egypt] (Cenomanian-Turonian) <HT: KBIN IST 10483>.

H. depressus ABDELHAMID & AZAB, 2003: 856-857; pl. 1, figs M-O. [Galala Fm., Saint Paul, north Eastern Desert, Egypt] (Cenomanian) <HT: Geol. Dept., Fac. Sci., Ain Shams Univ., Cairo>.

Aptian-Albian

H. psiloses AZIZ, 1991: 15-16; pl. 1, fig. 8. [Dalmiapuram Formation, Trichinopoly Subbasin, Southern India] (Late Aptian to Early Albian, Early Cretaceous) <HT: MACSG 1921> {based on isolated spines}.

H. rhabdoses AZIZ, 1991: 16-17; pl. 1, fig. 9. [Dalmiapuram Formation, Trichinopoly Subbasin, Southern India] (Late Aptian to Early Albian, Early Cretaceous) <HT: MACSG 1927> {based on isolated spines}.

H. teraniensis AZIZ, 1991: 13-14; figs 2d, pl. 1, figs 3-4. [Dalmiapuram Formation, Teran, Trichinopoly Subbasin, Southern India] (Late Aptian to Early Albian, Early Cretaceous) <HT: MACSG 1945>.

Berriasian

H. sidibouzidense VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007: 14-15; pl. 5: figs 1-7. [Falaise de Borj Nador, N of Safi, Morocco] (Late Berriasian, Early Cretaceous) <Type series: Maignac coll. 63 to 67, Nicollet coll. 7271, Reboul coll. 602, Vadet coll. V7482, 7483; spines: Maignac coll. 36 to 41, Nicollet coll. 5197, 7206, Reboul coll. 321, Vadet coll. V7449(1-3)>.

Bathonian

H. ederlei VADET, EDERLÉ & ROBERT, 1995: 104-105; pl. 5; pl. 6, fig. 2. [Calcaires compacts, Yonne, France] (Upper Bathonian) <HT: Robert Ederlé coll. 123-485; PT: Robert Ederlé coll. 123-1734>.

Genus *Heterodiadema* COTTEAU, 1864

SMITH & WRIGHT (1993: 218) placed this genus in its own, new family Heterodiadematidae SMITH & WRIGHT, 1993, which they placed within the order Phymosomatoida MORTENSEN, 1904

Maastrichtian

H. buhaysensis SMITH, 1995: 133-136; fig. 10, 11; pl. 2: figs 1-3. [Simisima Fm., Jebel Buhays, Jebel Rawdah and Jebel Thanais, Oman/United Arabian Emirates Border Region] <HT: BMNH EE3441; PT: BMNH EE3442-5, EE5019>.

Genus *Pseudocidaris* ÉTALLON, 1859

SMITH & WRIGHT (1993: 199) placed this genus in its own, new family Pseudocidaridae
SMITH & WRIGHT, 1993.

Upper Jurassic

P. raratuberculata KIER, 1972a: 42-43; fig. 17; pl. 11, figs 5-6; pl. 12, figs 1-8. [Upper Dhurma Fm., locality KK9-111, Saudi Arabia] (Callovian) <HT: USNM 170384; PT: USNM 170385>.

P. romani KIER, 1972a: 45-46; fig. 19; pl. 13, figs 1-5; pl. 14, figs 1-4; pl. 15, figs 1-2. [Upper Dhurma Fm., localities KK9-112, KK9-111, Saudi Arabia] (Callovian) <HT: USNM 170386; PT: USNM 170387-170388>.

P. santacruzensis RADWAŃSKA, 1999: 318-319; pl. 23, figs 9-16. [Wierzbica, north-eastern margin of the Holy Cross Mountains, Central Poland] (Lower Kimmeridgian) <HT: DPUW EWi/127>.

Middle Jurassic

P. depressa KIER, 1972a: 43-45; fig. 18; pl. 15, figs 3-6; pl. 16, figs 1-4. [Middle Dhurma Fm., locality KK9-21-21.5, Saudi Arabia] (Bathonian) <HT: USNM 170389; PT: USNM 170390>.

Family Pseudocidaridae SMITH & WRIGHT, 1993

Pseudocidaridae SMITH & WRIGHT, 1993: 199. Type-genus: *Pseudocidaris* ÉTALLON, 1859.
{other genera included: *Cidaropsis* COTTEAU, 1863}.

Family Pseudodiademmatidae POMEL, 1883

Genus *Acrocidaris* L. AGASSIZ, 1840

VADET & NICOLLEAU (2000: 106) placed this genus into the family Acropeltidae LAMBERT
& THIÉRY, 1915.

Early Cretaceous

A. marocense VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007: 23-24; pl. 11: figs 1-5. [Falaise de Borj Nador, N of Safi, Morocco] (Late Berriasian, Early Cretaceous) <Type series: Maignac coll. 1071, Reboul coll. 640>.

Callovian

A. nicolleaui VADET & NICOLLEAU, 2000: 106-109; 5 figs (not numbered); pl. 3; pl. 4. [Fatima, Portugal] <HT: Nicolleau 4148; PT: Nicolleau 4147, 7004, 7005>.

Genus *Diplopodia* MACCOY, 1848

Jurassic

D. atlasensis VADET & NICOLLEAU, 2005: 28-29; illustr. on p. 28. [Region of Rich, Morocco] (Late Domerian (= Late Pliensbachian), Jurassic) <Type-series: Alain Vadet coll'n V6903, Philippe Nicolleau coll'n 8937, 9180>.

D. thieulini VADET, 1993: 90-91; fig. 74; pl. 10, figs 2-3. [Falaise de Saint Honorine les Pertes, Normandie, France] (Upper Bajocian, zone à *Parkinsoni*) <HT: MBSM coll. Thieulin>.

Genus *Girardema* VADET, 1993

Girardema VADET, 1993: 88. Type-species: *Diadema depressum* COTTEAU, 1850. [Normandie, France] (Bajocian) {other species included: *Diplopodia bipuncata* DESOR, 1857}.

Genus *Hypodiadema* DESOR, 1858

Upper Jurassic

H. nanituberculata KIER, 1972a: 46-48; fig. 20; pl.16, figs 5-10; pl. 17, figs 1-2. [Upper Dhurma Fm., localities KK9-111 and KK9-112, Saudi Arabia] (Callovian) <HT: USNM 170391>.

Genus *Pedinopsis* COTTEAU, 1863

Cenomanian

P. (P.) sphaerica SMITH, SIMMONS & RACEY, 1990: 50-54; figs 12a-e, 13. [Echinoid Marker Bed, Natih Fm., Jebel Madamar and Jebel Madar, Oman] (late middle Cenomanian) <HT: BMNH E83127; PT: BMNH E83128>.

Subgenus *Sinaiopsis* SMITH, SIMMONS & RACEY, 1990

Sinaiopsis SMITH, SIMMONS & RACEY, 1990: 54. Type-species: *Pedina sinaica* (AGASSIZ & DESOR, 1847). [Algeria, Egypt, Israel and Oman] (Cenomanian).

Genus *Polydiadema* LAMBERT, 1888

Cenomanian

P. bosei CHIPLONKAR & BADVE, 1972: 139-140; pl. 11: figs 8-10. [Mongra (22°00'30" N, 74°20'30" E), India] (Cenomanian) <HT: MACSG Mor. 123/69> {based on a single specimen}.

Valanginian

P. korotkovi PORETSKAYA, 1983: 113-115; figs 1, 2a-l. [vicinity of Kugusem Wells, Mangyshlak Peninsula, U.S.S.R.] (Lower Valanginian) <HT: LGU 346/1> {pagination of English translation, Russian original not seen}.

Middle Jurassic

P. ambiguum HESS, 1972: 58-62; figs 81-87, 89; pl. 2, fig. 1; pl. 4, fig. 2; pl. 14, fig. 2; pl. 17, fig. 4; pl. 18, fig. 4; pl. 19, fig. 2. [Upper Hauptrogenstein, Schinznach-Dorf, Switzerland] (Upper Bajocian) <HT: Hess coll. E 321>.

Genus *Pseudodiadema* MCCOY, 1848

Cretaceous

P. bakonyense SZÖRÉNYI, 1955: 28-30, 169-170; figs 9-11; pl. 1, figs 28-32. [groupes de manes argileuses, Zirc-Tündérmajor, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/30>.

Jurassic

P. vadeti NICOLLEAU & VADET, 1995: 74-75; 1 fig. on p. 74; pl. 30: figs 2A, B. [Marnes à spongiares, environs de Niort, Poitou, France] (*luciaiformis* Subzone, *transvesarium* Zone, Middle Oxfordian, Jurassic) <HT: Philippe Nicolleau coll. no. 2191>.

Genus *Tetragramma* L. AGASSIZ, 1840

Cretaceous

- T. gloriae* BUITRÓN, 1973: 38-39; pl. 1: figs 1-7. [San Lucas Formation, San Nicolás, Huétamo Region, south-south-eastern Michoacán, Mexico] (Hauterivian-Aptian, Early Cretaceous) <HT: IGM-2456; PT: IGM-2455> {designation of holotype and paratype added in handwriting in some copies of the paper}.
- T. variolare baconicum* SZÖRÉNYI, 1955: 33-34, 172-173; figs 12-13; pl. 1, figs 33-35. [marne glauconieuse, Pénzeskút-Körisgyörpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/56>.

Genus *Trochotiara* LAMBERT, 1901

RADWAŃSKA (1999: 319) placed this genus into the order Phymosomatoida MORTENSEN, 1904.

Coniacian

- T. kiiensis* TANAKA, 1984b: 189-190; text-fig. 1; pl. 1, figs 1a-b, 2a-b; pl. 2, figs 5. [Matsubara Fm., Ikadachi, Kanaya-cho, Arida-gun, Wakayama Prefecture, Kii Province, Japan] (?Coniacian) <HT: GSJ F6073A, B; PT: GSJ F6074>.

Cenomanian

- T. moabitorum* BANDEL & GEYS, 1985: 107-108; pl. 5, figs 5-7; pl. 6, figs 1-3. [Rumeimin Fm., Rumeimin, near Amman, Jordan] (Cenomanian) <HT: KBIN coll., IST 10214>.

Kimmeridgian

- T. kongieli* RADWAŃSKA, 1999: 319-321; pl. 24, figs 1-5; pl. 25, figs 1-6. [Czarnogłowy near Szczecin, Western Pomerania, Poland] (Lower Kimmeridgian) <HT: DPUW ECz/128>.
- T. suleiovense* RADWAŃSKA, 1999: 321; pl. 26, figs 1a-c. [Sulejów, south-western margin of the Holy Cross Mountains, Central Poland] (Lower Kimmeridgian) <HT: DPUW ESI/133>.

Subfamily Polydiadematinae HESS, 1972

Polydiadematinae HESS, 1972: 58. Type-genus: *Polydiadema* LAMBERT, 1888. {placed into the family Pseudodiademataidae by HESS (1972)}.

Family Emiratidae ALI, 1990

Emiratidae ALI, 1990: 103. Type-genus: *Emiratus* ALI, 1990. {emended to Emiratiidae by SMITH & WRIGHT, 1993: 248; they included also the genera *Allomma* POMEL, 1883 and *Polydiadema* LAMBERT, 1888 in this family}.

Genus *Emiratus* ALI, 1990

Emiratus ALI, 1990: 103. Type-species: *Emiratus raskhaimahensis* ALI, 1990. [UAE] (Cenomanian) {emended to *Emiratia* by SMITH & WRIGHT (1993: 248)}.

Cenomanian

E. raskhaimahensis ALI, 1990: 103-105; figs 3 (6-11). [Mauddud Fm., Wadi Kated al Qa's, Ras al Kaimah, United Arab Emirates] (Cenomanian) <HT: MGD-UAA > {no specimen no. given}.

Family Uncertain

Genus *Allomma* POMEL, 1883

SMITH & WRIGHT (1993: 254-255) included this genus into the family Emiratiidae ALI, 1990.

Cretaceous

A. kalon SZÖRÉNYI, 1955: 26-27, 167-168; figs 7-8; pl. 1, figs 24-26. [marne glauconieuse, Bakonynána (carrière de la vallée Gaja), Pézseskut, and Körisgyörpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/27 (from Bakonynána)>.

A. wrighti SMITH & WRIGHT, 1993: 261-262; text-figs 88, 89F-H; pl. 90, figs 1-8; pl. 91, figs 1-3. [Grey Chalk, Folkestone and Dover, Kent, England] (Cenomanian) <HT: SM B50355; PT: SM B32755-58, BMNH 75381, E1084, EE1295>.

Genus *Trochodiadema* DE LORIO, 1900

ROMAN (1991: 53) placed this genus into the family Pseudodiadematidae POMEL, 1883.

Albian

T. ? dhofarensis ROMAN, 1991: 53-56; fig. 2; pl. 1, figs 1-3, 7-9. [Dhalqut Fm. and Kharfot Fm., Umbaraaf, Dhofar, Oman] (Middle to Late Albian) <HT: MNHN RO9027; PT: 8 specimens, not numbered>.

Order Phymosomatoida MORTENSEN, 1904

Family Phymosomatidae POMEL, 1883

Genus *Gauthieria* LAMBERT, 1888

Campanian

G. ? mosae GEYS, 1980: 215-218; figs 5/1-3, figs 6/1. [Lower Gulpen Chalk, Heure-le-Romain, Liège, Belgium] (Upper Campanian) <HT: KBIN 10178> {JAGT (2000: 233) confirmed the placing of this species in the genus *Gauthieria*}.

Genus *Gauthiosoma* KUTSCHER, 1985c

Gauthiosoma KUTSCHER, 1985c: 524. Type-species: *Phymosoma princeps* v. HAGENOW, 1840. [Northern Germany] (Campanian-Maastrichtian).

Genus *Hemithylus* ARNAUD, 1895

Maastrichtian

H. alternus KUTSCHER, 1985b: 239-240; pl. 1, figs 11-12; pl. 3, figs 5-12. [Quarry Wittenfelde, Rügen, Germany] (Late Early Maastricht) <HT: Kutscher coll. 1493/1> {transferred to *Gauthieria* by JAGT et al. (1998: 23) and SMITH & JEFFERY (2000: 89)}.

Genus *Kachchhia* SRIVASTAVA, GUPTA & JAUHRI, 2008

Kachchhia SRIVASTAVA, GUPTA & JAUHRI, 2008: 107-108. Type-species: *Kachchhia krohi* SRIVASTAVA, GUPTA & JAUHRI, 2008. [India] (Eocene).

Eocene

K. krohi SRIVASTAVA, GUPTA & JAUHRI, 2008: 108-110; pl. 1: figs 1-9. [Fulra Limestone, Three km SSE of Harudi Village, Kachchh, Gujarat, India (23°29'21.6" N, 68°41'48.9" E)] (NP17 Zone, Bartonian, late Middle Eocene) <HT: LUGD/I/2028> {based upon a single specimen}.

Genus *Micropsis* COTTEAU, 1856

Maastrichtian

M. (?) caementum JAGT & VAN DER HAM, in JAGT, 2000: 239-240; pl. 12, figs 12-13. [Emael Mb., Maastricht Fm., ENCI-Maastricht Quarry, Maastricht, The Netherlands] <HT: NHMM K 2681>.

Genus *Phymosoma* HAIME, 1853

Maastrichtian

P. maastrichtensis ENGEL, 1972: 540-543; pl. 1. [Belvédère, Caberg, Prov. Zuid-Limburg, The Netherlands] (Maastrichtian) <HT: NHMM 1340> {JAGT (2000: 242) referred this species to the genus *Circopeltis*}.

Cenomanian

P. mongraensis CHIPLONKAR & BADVE, 1972: 140-141; pl. 11: figs 5-6. [Mongra (22°00'30" N, 74°20'30" E) and Guneri, 4 miles E of Walpur (22°07'30" N, 74°29' E), India] (Cenomanian) <HT: MACSG Mor. 114/69>.

Aptian-Albian

P. dalmiapurensis AZIZ, 1991: 12-13; figs 2b, pl. 1, figs 1-2. [Dalmiapuram Formation, Dalmiapuram Mine, N of Kallakkud, Trichinopoly Subbasin, Southern India] (Late Aptian to Early Albian, Early Cretaceous) <HT: MACSG 1944>.

P. paynei SMITH & WRIGHT, 1996: 276-277; text-fig. 94; pl. 94, fig. 6; pl. 95, figs 1-2; pl. 110, figs 1-3. [Shenley Limestone, Double Arches Pit, Shenley Hill, Leighton buzzard, Bedfordshire, England] (*L. tardefurcata* zone, *L. regularis* subzone, Lower Albian) <HT: BMNH EE5514; PT: BMNH EE5515-16>.

Genus *Pleurodiadema* DE LORIOU, 1870

Middle Jurassic

P. raboeufi VADET, NICOLLEAU & PINEAU, 1996: 98-100; figs 116-119; pl. 25, figs 1a-2.. [Oolithe de l'Antonnière, Sarthe, France] (Callovian) <HT: Raboef coll.; PT: Pineau coll. 311> {no specimen number for the holotype given}.

Genus *Pomeliosoma* VADET, 2005b

Pomeliosoma VADET, 2005b: 136-137. Type-species: *Cyphosoma legayi* RIGAUD, 1882. [France & Germany] (Kimmeridgian-Tithonian, Late Jurassic) {other species included: *Cyphosoma duplicatum* COTTEAU, 1885}.

Genus *Porosoma* COTTEAU, 1856

Paleocene

P. fifei WAGNER, 1972: 652-655; text-figs 1-4; pl. 1, figs 1-3, 5-7. [Sepultura Fm., El Cardon area, Baja California, Mexico] (Late? Paleocene) <HT: UCMP 10782; PT: UCMP 10783-10785, USNM 178626>.

Campanian

P. kaspicum GEYS, 1984: 28-29; figs 2-6. [Shhakh-Bogota, Mangyshlak, Kazachstan, U.S.S.R.] (*Belemnitella langei*-zone, Upper Campanian) <HT: KBIN IST 10210>.

Genus *Rachiosoma* POMEL, 1883

Maastrichtian

R. gigasei GEYS, 1983: 255-256; pl. 1, figs 6. [Upper Gulpen Chalk, Lixhe, Liège, Belgium] (Lower Maastrichtian) <HT: KBIN IST 10204>.

Genus *Thylechinus* POMEL, 1883

Paleocene

T. (T. ?) tessieri ADEGOKE, 1977: 60-61; pl. 4: figs 18-21. [Ewekoro Fm., Ewekoro quarry, Ewekoro, 50 km N Lagos, Nigeria] (Paleocene) <HT: UIMG 166> {based on single specimen}.

T. vanderhami SMITH & JEFFERY, 2000: 124. [Geulhem Mb., Maastricht, The Netherlands] (Early Danian) <NHMM MB432N>.

Subfamily Circopeltinae SMITH & JEFFERY, 2000

Circopeltinae SMITH & JEFFERY, 2000: 110-111. Type-genus: *Circopeltis* POMEL, 1883. {other genera included: *Phymechinus* DESOR, 1856; and *Phymotaxis* LAMBERT & THIÉRY, 1914}.

Family Diplopodiidae SMITH & WRIGHT, 1993

Diplopodiidae SMITH & WRIGHT, 1993: 221. Type-genus: *Diplopodia* MCCOY, 1848. {other genera included: *Tetragramma* L. AGASSIZ, 1840; *Tiaromma* POMEL, 1883}.

Family Heterodiadematidae SMITH & WRIGHT, 1993

Heterodiadematidae SMITH & WRIGHT, 1993: 218. Type-genus: *Heterodiadema* COTTEAU, 1864. {further genera included: *Trochodiadema* DE LORIO, 1900}.

Family Polydiadematidae VADET, 1999b

Polydiadematidae VADET, 1999b: 79. Type-genus: *Polydiadema* LAMBERT, 1888.

Family Stomechinidae POMEL, 1883

Genus *Baronechinus* VADET & NICOLLEAU, 2005

Baronechinus VADET & NICOLLEAU, 2005: 29. Type-species: *Baronechinus baroni* VADET & NICOLLEAU, 2005. [Morocco] (Early Jurassic).

Jurassic

B. baroni VADET & NICOLLEAU, 2005: 29-30; illustr. on p. 29 and 30. [surroundings of Rich, Morocco] (Late Domerian (= Late Pliensbachian), Jurassic) <no types designated; Material studied: Philippe Nicolleau coll'n 8922, 8923, 8925, 9179, Roland Reboul coll'n 535, Alain Vadet coll'n V6715, V6734-V6736, V6920, V6923>.

Genus *Circopeltis* POMEL, 1883

Maastrichtian

C.? *emiratus* SMITH, 1995: 172; fig. 37; pl. 15: figs 1-3; pl. 17: figs 1-2. [Jebel Buhays, Oman/United Arabian Emirates Border Region] <HT: BMNH EE3584; PT: BMNH EE3582, EE3583, EE3585, EE3596>.

Genus *Codechinus* DESOR, 1856

Barremian

C. prozorovskyi PORETZKAJA, 1989: 161-165; figs a-b; unnumbered pl., figs 1a-6, 2a-b, 3. [Lesser Balkan, Turkmenistan] (Barremian) <HT: ЭКЗ. No. 305/21>.

Genus *Diplotagma* SCHLÜTER, 1870

SMITH & JEFFERY (2000: 87) placed this genus into the subfamily Phymosomatinae POMEL, 1883, within the family Phymosomatidae POMEL, 1883.

Maastrichtian

D. snellingsi SMITH & JEFFERY, 2000: 87-88; text-fig. 33A-B. [Maastricht Fm., Maastricht district, The Netherlands or Belgium] (Late Maastrichtian) <HT: BMNH E3186>.

Genus *Leioechinus* KIER, 1972a

Leioechinus KIER, 1972a: 48. Type-species: *Leioechinus namus* KIER, 1972a. [Saudi Arabia] (Middle to Late Jurassic).

Upper Jurassic

L. amplus KIER, 1972a: 52-53; figs 24-25; pl. 22, figs 1-7. [Upper Dhurma Fm., localities KK9-111-112, S-1176 and L-926, Saudi Arabia] (Callovian) <HT: USNM 170397; PT: USNM 170398-170401>.

Middle Jurassic

L. namus KIER, 1972a: 48-52; figs 21-23; pl. 23, figs 1-6; pl. 24, figs 1-11. [Middle Dhurma Fm., localities KK8-30-35, KK7-131, KK8-6-40.5, KK9-20-40, S-1064, S-1154, S-1160 and L-921, Saudi Arabia] (Bathonian) <HT: USNM 170402; PT: USNM 170403-170408>.

Genus *Noetlingaster* VREDENBURG, 1911

Maastrichtian

N. monotuberculatus SMITH, in SMITH & JEFFERY, 2000: 77; text-fig. 28J, L. [Simsima Fm., United Arab Emirates-Oman border region] (*A. fresvillensis* zone, mid Maastrichtian) <not given> {new name for *Hattopsis paucituberculatus* SMITH, 1995}.

N. emiratescus ALI, 1989a: 398-400; figs 2(6), 3. [Simsima Fm., Gebel El Rowdah, United Arab Emirates] (Late Maastrichtian) <MGD-UAA > {no type specimen defined}.

Genus *Phymechinus* DESOR, 1856

Maastrichtian

P.? *perplexus* SMITH, 1995: 172-175; fig. 38, 39; pl. 15: figs 4-10. [Jebel Rawdah, Oman/ United Arabian Emirates Border Region] <HT: BMNH EE3579; PT: BMNH EE3581, EE3591, EE3593, EE3619>.

Genus *Polycyphus* L. AGASSIZ & DESOR, 1846

Upper Jurassic

P. parvituberculatus KIER, 1972a: 58-59; pl. 18, figs 4-8. [Upper Dhurma Fm., locality KK9-112, Saudi Arabia] (Bathonian) <HT: USNM 170394>.

Middle Jurassic

P. arabicus KIER, 1972a: 56-58; fig. 26; pl. 19, figs 1-7; pl. 20, figs 1-4. [Middle Dhurma Fm., localities KK8-46 and KK9-10-20, Saudi Arabia] (Bathonian) <HT: USNM 170395; PT: USNM 170396>.

Genus *Psephechinus* POMEL, 1883

Middle Jurassic

P. pavyi ROBERT, 1994: 128-129; pl. 20: figs 1-5. [Calcaires à chailles; Brétignelles, près Druyes, Dept. Yonne, France] (*Antecedens* subzone, *Plicatilis* zone, Middle Oxfordian) <HT: Philippe Robert coll'n No. 287>.

Genus *Stomechinus* DESOR, 1856

Oligocene

S. ? dissimilaris LINDER, DURHAM & ORR, 1988: 948; fig. 2.6. [Scott Mills Fm., Abiqua Mb., Butte Creek, Marion County, Oregon, USA] (Late Oligocene) <HT: UCMP 38196>.

Jurassic

S. tinginatus VADET & NICOLLEAU, 2005: 33-34; illustr. on p. 33 and 34. [surroundings of Rich, Morocco] (Late Domerian (= Late Pliensbachian), Jurassic) <no types designated; Material studied: Philippe Nicolleau coll'n 8922, Roland Reboul coll'n 534, Alain Vadet coll'n V6714, V6733, V6751, V6769-V6775>.

Genus *Trochalosoma* LAMBERT, 1897

T. (T.) gharamulensis ABDELHAMID & AZAB, 2003: 860-862; pl. 3, figs B-D. [Galala Fm., G. Gharamul, north Eastern Desert, Egypt] (Cenomanian) <HT: Geol. Dept., Fac. Sci., Ain Shams Univ., Cairo>.

Family Uncertain

Genus *Boletechinus* COOKE, 1955

LEWIS (1986: 61) placed this genus into the Family Zeuglopleuridae LEWIS, 1986.

Maastrichtian

B. delawaricus LEWIS, 1986: 71-74; figs 2a, 3a-f, 5a-d, 6, 7. [Navesink Marl, Monmouth Group, Delaware Canal, Delaware, U.S.A.] (Maastrichtian) <HT: BMNH E76803; PT: BMNH E7604-05>.

Senonian

B. rowei anglicus LEWIS, 1986: 78-81; figs 2a, 3a-f, 9a-e, 10. [Charlton, Kent, England] (Senonian) <HT: BMNH E75556a; PT: BMNH E75556b, E39376-8; GSM 118257-9>.

Order Stomechinoida VADET, 1999b

Stomechinoida VADET, 1999b: 79. {included families: Atopechinidae THIÉRY, in THIÉRY, LAMBERT & COLLIGNON, 1928; Stomechinidae POMEL, 1883}.

Order Arbacioida GREGORY, 1900

Genus *Masrouraster* VADET, NICOLLEAU & REBOUL, 2008

Masrouraster VADET, NICOLLEAU & REBOUL, 2008: 15-16. Type-species: *Masrouraster ouhouissi* VADET, NICOLLEAU & REBOUL, 2008. [Morocco] (Early Jurassic).

Early Jurassic

M. ouhouissi VADET, NICOLLEAU & REBOUL, 2008: 16-17; unnumbered figs on p. 16 and 17. [environs d'Amellago, Haut Atlas, Maroc] (Domerian, Late Pliensbachian) <ST: Reboul coll'n 718, Nicolleau coll'n 9996>.

Family Arbaciidae GRAY, 1855

Genus *Arbacia* GRAY, 1835

Oligocene

A. abiquaensis LINDER, DURHAM & ORR, 1988: 948; figs 3.1-3.6. [Scott Mills Fm., Abiqua Mb., Butte Creek, Marion County, Oregon, USA] (Late Oligocene) <HT: UCMP 39198; PT: UCMP 38199-A, 38199-B>.

Genus *Acropeltis* L. AGASSIZ, 1840

Lower Cretaceous

A. atlantica REY, TAJ-EDDINE & WITAM, 1989: 603-604; fig. 3; pl. 1, figs 1-11. [Borj Nador, 6 km north of Safi, Morocco] (Late Berriasian-Early Valanginian) <HT: LGUT BN 13; PT: LGUT BN 9 – BN 12, BN 14 – BN 22>.

Genus *Baueria* NOETLING, 1885

Eocene

B. angelae CARRASCO, 2006: 23-32; figs 3-4; pl. 1, figs 1a-d, 2a-d. [Serraduy Fm., El Carrasquero, Tremp-Graus, Zona Central Surpirenaica, NW Barcelona, Spain] (Middle Illerian, Early Eocene) <HT: MGSB 73.406a; PT: MGSB 73.406b, 73.407>

Genus *Codiopsis* L. AGASSIZ, 1840

Maastrichtian

C. lehmannae SMITH, 1995: 152-153; fig. 23; pl. 9: figs 1-2; pl. 12: figs 1-3. [Simisima Fm., Jebel Buhays and Jebel Rawdah, Oman/United Arabian Emirates Border Region] <HT: BMNH EE5033; PT: BMNH EE3439, EE3440>.

Cenomanian

C. smiseri GEYS, 1985: 142-143; pl. 5, figs 3-7. [Tournai, Hainaut, Belgium] (Cenomanian) <HT: KBIN IST 9121>.

Jurassic

C. rupellensis VADET, NICOLLEAU & RIGOLLET, 2002: 60-61; 7 figs [Le Chay, près de La Rochelle, Charente Maritime, France] (Kimmeridgian) <HT: Bertrand coll. 31K ; PT: Rigollet coll. CY 110, Guenne coll. KI27>.

Genus *Coelopleurus* L. AGASSIZ, 1840

Recent

C. exquisitus COPPARD & SCHULTZ, 2006: 4-14; figs 1A-C, 2A-C, 3A-N, 4A-K, 5; tab. 1a. [N.O. "Vauban" MUSORSTOM 4, Stat. DW181 (18°57' S 163°22' E, depth 350 m; C. Vadon Coll. 18th September, 1985), N.O. "Jean-Charcot" BIOCAL, Stat. DW50 (23°07' S, 167°54' E, depth 240–260 m; Guille and Menau Collection, 31th August 1985), and Coreolus Expedition, South of Isles of Pines (23°06' S, 167°05' E, depth 520 m); all off New Caledonia] (Recent) <HT: MNHN EcEh 1281 (from MUSORSTOM 4, Stat. DW181); PT: MNHN EcEh 1282 (same locality as the HT), BMNH 2006.599>.

Palaeogene

C. ulugqatensis YANG SHENGQIU, 1991: 114; pl. 5, figs 1-5; pl. 6, figs 3-4. [Wulagen Fm. Bashibulake, Wuqia County, Tarim Basin, China] (Early Tertiary) <HT: NIGP 88338>.

Genus *Cottaldia* DESOR, 1856

Upper Jurassic

C. paquettei VADET, 1995: 114-115; figs 53-54; pl. 10, figs 1-3. [Marnes à Spongiaires, Poitou, France] (Oxfordian) <HT: V 3495>.

Genus *Glypticus* L. AGASSIZ, 1840

Middle Jurassic

G. icaunense ROBERT, 1994: 88-89; pl. 18: figs 1-4. [Calcaires blancs; Andryes, Yonne, France] (*Bifurcatus* subzone, *Transversarium* zone, Middle to Late Oxfordian) <HT: Philippe Robert coll'n No. 874>.

Genus *Goniopygus* L. AGASSIZ, 1838

Maastrichtian

G. arabicus SMITH, 1995: 142-47; fig. 17, 18A, B, D, E, G; pl. 6: figs 3-10; pl. 7: figs 1, 3, 5-6. [Simisima Fm., Jebel Buhays, Jebel Faiyah, Jebel Rawdah and Jebel Thanaïs, Oman/United Arab Emirates Border Region] <HT: BMNH EE4012; PT: BMNH EE3983-84, EE3986, EE3992, EE3997, EE4005, EE4007, EE4015, EE4017, EE4019>.

Genus *Hattopsis* ALI, 1992b

Hattopsis ALI, 1992b: 694. Type-species: *Hattopsis sphericus* ALI, 1992b. [Middle East] (Maastrichtian) {SMITH, in SMITH & JEFFERY (2000: 75) placed this genus into the synonymy of *Noetlingaster* VREDENBURG, 1911}.

Maastrichtian

H. sphericus ALI, 1992b: 694-695; figs 3-5. [Simsima Fm., Gebel El Rowdah, Hatta area, Oman-U.A.E.] (Late Maastrichtian) <HT: MGD-UAA 910401; PT: MGD-UAA 910402>.

H. paucituberculatus SMITH, 1995: 157-159; fig. 24, 25A, 26A, 27B; pl. 9: figs 9-11; pl. 10: figs 3, 6; pl. 11: fig. 9. [Simisima Fm., Jebel Aqabah, Jebel Buhays, Jebel Faiyah and Jebel Rawdah, Oman/United Arab Emirates Border Region] <HT: BMNH EE3683; PT: BMNH EE3682, EE3678, EE3688, EE3684-85> {SMITH, in SMITH & JEFFERY (2000: 77) proposed the name *Noetlingaster monotuberculatus* for this species, since the name became a subjective homonym through the synonymy of *Hattopsis* and *Noetlingaster*}.

Genus *Magnosia* MICHELIN, 1858

Lower Cretaceous

M. densituberculata REY, TAJ-EDDINE & WITAM, 1989: 604-605; pl. 2, figs 5-9. [Borj Nador, 6 km north of Safi, Morocco] (Late Berriasian-Early Valanginian) <HT: LGUT BN 36; PT: LGUT BN 34>.

Genus *Magnosiopsis* ŽITŤ, 1986

Magnosiopsis ŽITŤ, 1986: 375. Type-species: *Magnosia suessi* LORIOLE, 1901. [Kopřivnice Fm., Štramberk, Czechoslovakia] (Late Valanginian).

Valanginian

M. ornata ŽITŤ, 1986: 379-381; text-figs 2F, 8C; pl. 2, figs 4-6. 5. [Kopřivnice Fm., Kotouč massif, Štramberk, Czechoslovakia] (Late Valanginian) <HT: NMP 0 5429; PT: NMP 0 5430-5434>.

Genus *Mimiosalenia* SMITH, 1995

Mimiosalenia SMITH, 1995: 147. Type-species: *Mimiosalenia quinquetuberculata* SMITH, 1995 [Oman/UAE] (Maastrichtian) {SMITH (1995: 147) placed this genus in the family Goniopygidae SMITH & WRIGHT, 1993}.

Maastrichtian

M. quinquetuberculata SMITH, 1995: 148-149; fig. 19-20; pl. 5: figs 4-10, 12. [Jebel Faiyah, Oman/United Arabian Emirates Border Region] <HT: BMNH EE3981; PT: BMNH EE3971, EE3974, EE3977-78, EE3980, EE3982, EE5014-17>.

Genus *Sexpyga* SHIGEL, 1975b

Sexpyga SHIGEL, 1975b: 329. Type-species: *Sexpyga soyoae* SHIGEL, 1975b. [North-west Pacific] (Recent).

Recent

S. soyoae SHIGEL, 1975b: 330-331; figs 1-9. [SW of 20°N, Hachijo Island, North Pacific Ocean, Soyo-Marun Stat. B4 (32°04'0" N, 140°21'5" E), depth 1940-1980 m] (Recent) <HT: MMBS Echi 1013>.

Family Coelopleuridae MÄRKEL, 1969

Coelopleuridae MÄRKEL, 1969: 24. Type-genus: *Coelopleurus* L. AGASSIZ, 1840.

Family Glyphopneustidae SMITH & WRIGHT, 1993

Glyphopneustidae SMITH & WRIGHT, 1993: 209. Type-genus: *Glyphopneustes* POMEL, 1869. {other genera included: *Arbia* COOKE, 1948}.

Family Glypticidae LAMBERT & THIÉRY, 1914

Glypticidae VADET, 1999a: 83. Type-genus: *Glypticus* L. AGASSIZ, 1840 {preoccupied by Glypticidae LAMBERT & THIÉRY, 1914}

Family Atopechinae THIÉRY, in THIÉRY, LAMBERT & COLLIGNON, 1928

Atopechinae [Atopechinaeæ] THIÉRY, in THIÉRY, LAMBERT & COLLIGNON, 1928: 101. Type-genus: *Atopechinus* THIÉRY, in THIÉRY, LAMBERT & COLLIGNON, 1928 [Europe] (Jurassic) {apparently used only rarely subsequently; elevated to family rank by VADET et al. (2007)}.

Order Temnopleuroida MORTENSEN, 1942

Family Glyphocyphidae DUNCAN, 1889

Genus *Ambipleurus* LAMBERT, 1932

Eocene

A.? *quaylei* LEWIS, 1989: 17-19; figs 2 f-h; pl. 2: figs 6 a-b, 7. [Barton-on-Sea, Hampshire, England] (Bartonian) <HT: BMNH E 76822; PT: BMNH E 76823-4, E 76923>.

A. viladensis CARRASCO, 2007: 11-15; fig. 6; pl. 1: figs a-g. [Vilada and Sant Llorenç de Morunys, N of Barcelona, Spain] (Lutetian, Middle Eocene) <HT: MGSB 10.969 (from Sant Llorenç de Morunys); PT: Pere Barniol colln 1785, 1768, 253a, 253b, s/n (from Vilada; to be deposited at the Museu Municipal de Berga), Joan Maria Viader colln 730 (from Vilada)>.

Genus *Bandelicyphus* GEYS, 1992

Bandelicyphus GEYS, 1992: 145. Type-species: *Bandelicyphus genaensis* GEYS, 1992. [Egypt] (Turonian).

Turonian

B. genaensis GEYS, 1992: 145-146; pl. 2; figs 1-7. [Tarfa Fm., Wadi Qena, Eastern Desert, Egypt] (Turonian) <HT: KBIN IST-10498>.

Family Temnopleuridae A. AGASSIZ, 1872

Genus *Arbacina* POMEL, 1869

Pliocene

A. hugueti DUDICOURT, NÉRAUDEAU, NICOLLEAU, CEULEMANS & BOUTIN, 2005: 553-556; pl. 4, figs 1-7. [La Minotière, Challans, Vendée, western France] (Pliocene) <HT: Musée de Niort, Nicollet coll. no. 8279; PT: MNHN-DHT R64551, A24829>.

A. pareyni ROMAN, 1983: 18-20; pl. 2, figs 9-12. [Saint-André de Bohon (Manche) and E of Rougeville, Western France] (Redonien récent {Late Pliocene}) <HT: IPM R 50799>.

Miocene

A. emmae NÉRAUDEAU, BARBE, MERCIER & ROMAN, 2003: 162-163; fig. 2; pl. 1: fig. 5; pl. 2: figs 4-6. [Lilion, Saint-Jacques-de-la-Lande, Ille-et-Vilaine, Western France] (Messinian (?), Late Miocene) <HT: MNHN coll. Barbe, no specimen no. given>.

Genus *Brochopleurus* FOURTAU, 1920

Pliocene

B. pliocenicus DEVIRIES, 1973: 84-86; pl. 5: figs 13-17. [Algeria] (Pliocene) <no type nos. mentioned, repository unknown>.

Genus *Echinocyphus* COTTEAU, in COTTEAU & TRIGER, 1860

SMITH & WRIGHT, 1996: 313 placed this genus into the Plesion (Family) Zeuglopleuridae LEWIS, 1986.

Cenomanian

E. intermedius SMITH & WRIGHT, 1996: 323-324; text-figs 117A, B, 119; pl. 112, figs 3-5, 8. [Plenus Marls, Betchworth, Surrey, England] (*M. geslinianum* zone, Upper Cenomanian) <HT: BMNH EE83314>.

Genus *Genocidaris* A. AGASSIZ, 1869

Pliocene

G. maculata pliorecens BORGI, 1995: 6; pl. 1, figs 4; pl. 2, figs 4-6; pl. 3, figs 1-3; pl. 4, figs 1-3; pl. 5, figs 1-2, 5-6. [near Castell'Arquato and Salsomaggiore, Northern Italy] (Pliocene – Pleistocene) <not given>.

Genus *Microcyphus* L. AGASSIZ, in AGASSIZ & DESOR, 1846

Quarternary

M. iglahensis ELATTAAR, 2001a: 645-646; fig. 6A; pl. 1: figs 4-9. [Raised reed at Wadi Igla, S of Wadi Wizr, 41 km SW Quseir, Red Sea coast, Egypt] (Quarternary) <3 syntypes, no numbers mentioned, AUSGM E collection>.

Genus *Nannoglyphus* NESTLER, 1978

Nannoglyphus NESTLER, 1978: 621. Type-species: *Nannoglyphus wehrlii* NESTLER, 1978. [Northern Europe] (Maastrichtian).

Maastrichtian

N. wehrlii NESTLER, 1978: 621-624, figs 2-3a-d. [Quarry Wittenfelde, Rügen, Germany] (Early Maastrichtian) <HT: SGWG 56/1> {this genus is placed into the synonymy of *Zeuglopleurus* by SMITH & JEFFERY (2000:129)}

Genus *Paradoxechinus* LAUBE, 1869

Oligocene

P. granulosus PHILIP & FOSTER, 1971: 674-676; pl. 129: figs 4, 5, 9, 10, 18, 19; pl. 134: fig. 3. [Point Addis Limestone, Airey's Inlet, Anglesea District, Victoria, Australia] (Janjukian, Upper Oligocene) <HT: NMV P27947; PT: UNE 11757, 11761>.

P. profundus PHILIP & FOSTER, 1971: 676-677; pl. 128: fig. 1; pl. 129: fig. 8; pl. 134, fig. 2. [Port Willunga Beds, Seaford, south of Port Noarlunga, South Australia] (Middle Oligocene to Lower Miocene) <HT: NMV P27945; PT: NMV P 27946, UNE 11754>.

Eocene

P. stellatus PHILIP & FOSTER, 1971: 677-678; pl. 126: fig. 2; pl. 129: figs 16, 20-23; pl. 134, fig. 1. [Port Willunga Beds, Onkaparinga River, Port Noarlunga, South Australia] (Upper Eocene) <HT: NMV P27944; PT: UNE 11753>.

Genus *Pentechinus* PHILIP & FOSTER, 1971

Pentechinus PHILIP & FOSTER, 1971: 678. Type-species: *Pentechinus mirabilis* PHILIP & FOSTER, 1971. [Victoria, Australia] (Oligocene).

Oligocene

P. mirabilis PHILIP & FOSTER, 1971: 678-681; text-figs 3-4; pl. 124: figs 1-3; pl. 129: figs 1-3, 11, 15. [Point Addis Limestone, Airey's Inlet, Anglesea District, Victoria, Australia] (Janjukian, Late Oligocene) <HT: NMV P27942; PT: UNE 11512>.

Genus *Temnotrema* A. AGASSIZ, 1863

Recent

T. scillae var. *eythraea* DOLLFUS & ROMAN, 1981: 56-57; figs 7-11; pl. 13, figs 7-9. [Station 11; Red Sea] (Recent) <HT: not given>.

T. xishaensis LIAO, 1978: 112-114 [Chinese], 126-127 [Engl.]; figs 5.1-5.2, 6; pl. 2: figs 2-4; pl. 5: figs 2, 3, 6-8. [Xisha Island, Yongxingdao, South China Sea] (Recent) <HT: IOAS E.00813>.

Pliocene

T. greifatensis ELATTAAR, 2001a: 646-647; fig. 6B-C; pl. 1: figs 10-14. [Sharm El Arab Member, Shagra Formation, Mersa Um Greifat, S of Wadi Wizr, 41 km SW Quseir, Red Sea coast, Egypt] (Early Pliocene) <7 syntypes, no numbers mentioned, AUSGM E collection>.

T. wizrensis ELATTAAR, 2001b: 83-86; figs 3A-C, 4A-D; pl. 1, figs 1-5. [Sharm El Arab Member, Shagra Formation, Wadi Wizr, 41 km SW Quseir, and Wadi Gassus, 10 km SW Safaga, Red Sea coast, Egypt] (Late Pliocene) <2 specimen, repository unknown> {based on a single specimen}.

Subgenus *Viaudechinus* ROMAN, 1983

Viaudechinus ROMAN, 1983: 4. Type-species: *Dicoptella bigoti* LAMBERT & THIÉRY, 1911. [Western France] (“Helvetian” {Miocene} – Redonian {Late Pliocene}).

Genus *Tremaster* DUDICOURT, NÉRAUDEAU, NICOLLEAU, CEULEMANS & BOUTIN, 2005

Tremaster DUDICOURT, NÉRAUDEAU, NICOLLEAU, CEULEMANS & BOUTIN, 2005: 552-553. Type-species: *Tremaster romani* DUDICOURT, NÉRAUDEAU, NICOLLEAU, CEULEMANS & BOUTIN, 2005. [Western Europe] (Pliocene).

Pliocene

T. romani DUDICOURT, NÉRAUDEAU, NICOLLEAU, CEULEMANS & BOUTIN, 2005: 553; pl. 3, figs 5-7. [La Minotière, Challans, Vendée (holotype + paratype 1) and Pierre Aigüe, Loire-Atlantique (paratype 2), western France] (Pliocene) <HT: MNHN-DHT R64088; PT: MNHN-DHT A24830, R64550>.

Genus *Trigonocidaris* A. AGASSIZ, 1869Subgenus *Tuberculocidaris* MARKOV, 1989

Tuberculocidaris MARKOV, 1989: 81. Type-species: *Trigonocidaris (Tuberculocidaris) tuberculata* MARKOV, 1989. [Northern Pacific] (Recent).

Recent

T. (Tuberculocidaris) tuberculata MARKOV, 1989: 81-82; figs 2б, д, з, л, 3б. [“Дмитрий Менделеев”-cruise (“Dmitrig Mendelejev”-cruise) Stat. 1255 (29°46' S, 167°59' E, depth 510 m), Fiji Sea, North Pacific Ocean] (Recent) <HT: IOANSSR, no specimen no. given >.

Subgenus *Lamprechinus* DÖDERLEIN, 1905

In the *Treatise* (1966: U423) this taxon is listed as genus. MARKOV (1989: 83) downgraded it to subgeneric status.

Recent

T. (Lamprechinus) nitidus nascaensis MARKOV, 1989: 82-83; figs 1г, 2в, е, ц, м, 3в. [“Ихтиандр”-6th cruise (“Ikhtiandr”-6th cruise) Stat. 11 (23°25' S, 83°19' W, depth 475 m), 12 (25°36' S, 82°27' W, depth 280 m), “Профессор Штокман”-cruise

(“Professor Shtokman”-cruise) Stat. 1957 (24°56' S, 88°31' W, depth 570 m), 1964 (24°57' S, 88°30' W, depth 570 m), 2013 (25°07' S, 99°41' W, depth 355 m), 2034 (25°04' S, 97°35' W, depth 485 m), Naska Ridge, East Pacific Ocean] (Recent) <HT: IOANSSR, no specimen no. given >.

Genus *Zeuglopleurus* GREGORY, 1889

LEWIS (1986: 61) placed this genus into the Family Zeuglopleuridae LEWIS, 1986.

Turonian

Z. dictyopleuroides SMITH & WRIGHT, 1996: 332-333; text-figs 120C, D, 121B, 122A; pl. 113, figs 4-6; pl. 114, fig. 12. [Middle Chalk, Whitecliff, near Seaton and The Hooken, near Branscomb, Devon, England] (*T. lata* zone, Middle Turonian) <HT: BMNH EE5526; PT: BMNH EE39794, EE5527>.

Family Toxopneustidae TROSCHERL, 1872

Genus *Lytechinus* A. AGASSIZ, 1863

Oligocene

L. baldwini LINDER, DURHAM & ORR, 1988: 950; figs 4.3, 4.4. [Scott Mills Fm., Abiqua Mb., Butte Creek, Marion County, Oregon, USA] (Late Oligocene) <HT: UCMP 38203>.

Genus *Nudechinus* H.L. CLARK, 1912

Recent

N. gravi var. *suezensis* DOLLFUS & ROMAN, 1981: 59-61; figs 12-18; pl. 15, figs 1-8. [Suez, Red Sea] (Quarternary – Recent) <HT: not given>.

Genus *Schizechinus* POMEL, 1869

Miocene

S. pentagonus KIER, 1972a: 89-90; fig. 47; pl. 55, figs 1-6; pl. 56, figs 1-6. [Dam Fm., locality S-126, Saudi Arabia] (Miocene) <HT: USNM 170473; PT: USNM 170474-170475>.

Genus *Tripneustes* L. AGASSIZ, 1841

Recent

T. gratilla elatensis DAFNI, 1983: 2-9; fig. 2A-B, fig. 4. [off Wadi Tweibe, NW Gulf of Elat] (Recent) <HT: HUI EC 1; PT: HUI EC 2-7, BMNH 1981.11.30.13>.

Miocene

[*T. misrai* SRIVASTAVA, 1988]: 151. [India] (*Fasciolites elliptica*-Zone, Middle Eocene) <unknown> {nomen nudum; no description or illustration}.

T. pregratilla MCNAMARA & KENDRICK, 1994: 42; figs 15F-H. [Poivre Fm., "The Ledge" Barrow Island, Western Australia] (Middle Miocene) <HT: WAM 82.326; PT: WAM 82.292.325, 82.328, 32.408, 82.413>.

Family Zeuglopleuridae LEWIS, 1986

Zeuglopleuridae LEWIS, 1986: 61-64. Types-genus: *Zeuglopleurus* GREGORY, 1889. [Europe and North America] (Upper Cretaceous) {other genera included: *Boletechinus* COOKE, 1955; *Glyptocyphus* POMEL, 1883 and probably *Echinocyphus* COTTEAU, 1860; Downgraded to subfamily status within the family Temnopleuridae A. AGASSIZ, 1872 by MARKOV & ENDELMAN, 1991}.

Genus *Sphaeropleurus* MARKOV & ENDELMAN, 1991

Sphaeropleurus MARKOV & ENDELMAN, 1991: 80. Type-species: *Sphaeropleurus geometricus* MARKOV & ENDELMAN, 1991. [Yemen] (Eocene).

Eocene

S. geometricus MARKOV & ENDELMAN, 1991: 80-82; figs B-F; pl. 1, figs 2a-f. [Tselyg Pant-sir, Yemen] (Lower Eocene) <HT: PIN 4387/1>.

Family uncertain

Genus *Gagaria* DUNCAN, 1889

Oligocene

G. crenularis LINDER, DURHAM & ORR, 1988: 953; figs 4.5-4.9. [Scott Mills Fm., Abiqua Mb., Butte Creek, Marion County, Oregon, USA] (Late Oligocene) <HT: UCMP 38205; PT: UCMP 38206>.

Order Echinoida CLAUS, 1876

Family Echinidae GRAY, 1825

Genus *Psammechinus* L. AGASSIZ & DESOR, 1846

Pliocene

P. exoletus McCRADY, in TUOMEY & HOLMES, 1855: 4; pl. 2: fig. 6. [Smith's, Goose Creek, South Carolina, USA] (Pliocene) <not given> {based on an adapical test fragment}.

Oligocene

P. carolinensis KIER, 1997: 5-6; fig. 2; pl. 3, figs 1-4; pl. 4, figs 1-5. [Trent Fm., Pollockville state quarry, North Carolina, USA] (Late Oligocene) <HT: USNM 398321; PT: USNM 398322, 398323, 398474>.

Genus *Sterechinus* KOEHLER, 1901

Recent

S. bernasconiae LARRAIN, 1975: 94-105; figs 109-129, tabs. 9-11, maps 9, 11, 15. [Bahia Tarn, Golfo de Penas and Estrecho de Concepcio, off Chilean coast, between 48°S and 51°S, South-east Pacific] (Recent) <HT: Museo del Departamento de Zoología, Universidad de Concepcion, Chile, no. 7969; PT: as HT, nos. 7970-7976>.

Family Echinometridae GRAY, 1825

Genus *Echinometra* GRAY, 1825

Recent

E. lucunter polypora PAWSON, 1978: 20-22; fig. 9. [Ascension and St. Helena, South Atlantic Ocean] (Recent) <HT: USNM E16206; PT: USNM E16179 to E16187, E16189 to E16191>.

Genus *Heliocidaris* L. AGASSIZ & DESOR, 1846

Recent

H. robertsi LINDLEY, 2004: 124-126; figs 4a-d, 5a-e. [Cape Gazelle, New Britain, East New Britain Province, Papua New Guinea] (Recent) <HT: ANU 60655>.

Genus *Pachycentrotus* H.L. CLARK, 1912

Recent

P. bajulus DARTNALL, 1972: 30-34; figs 1a-b, 2, 3a-e. [Darlington Beach, Maria Island, Tasman Sea off east coast of Tasmania, 42°44' S, 149°05' E and Eaglehawk Neck, SE Tasmania and Adventure Bay, Bruny Island, SE Tasmania] (Recent) <HT: TM. H603; PT: TM H604-H605, AM J6393, J7801, NMV H155>.

Family Strongylocentridae GREGORY, 1900

Genus *Mesocentrotus* TATARENKO & POLTARAUS, 1993

Mesocentrotus TATARENKO & POLTARAUS, 1993: 70. Type-species: *Strongylocentrotus franciscanus* (A. AGASSIZ, 1863). (Recent) {other species included: *Strongylocentrotus nudus* (AGASSIZ, 1863)}.

Family Parasaleniidae MORTENSEN, 1903

Genus *Diplosalenia* MORTENSEN, 1902

Pleistocene

D. faurei DOLLFUS & ROMAN, 1981: 69-71; pl. 36, figs 1-4. [W of Dalol, Afar, Ethiopia] (Pleistocene) <HT: MNHN 1979-7>.

Infraclass Proacroechinoidea VADET, 1999c

Proacroechinoidea VADET, 1999c: 87. {included orders: Plesiocidaroida DUNCAN, 1889; Triadocidaroida VADET, 1999c; Paurocidaroida VADET, 1999c}.

Order Plesiocidaroida DUNCAN, 1889

Family Kieritiaridae VADET, 1999c

Kieritiaridae VADET, 1999c: 88. Type-genus: *Kieritiaris* VADET, 1999c. {other genera included: *Batheritiaris* VADET, 1999a}.

Genus *Kieritiaris* VADET, 1999c

Kieritiaris VADET, 1999c: 88. Type-species: *Kieritiaris thieli* VADET, 1999c. [Southern Europe] (Carnian, Triassic).

Triassic

K. thieli VADET, 1999c: 88-89; fig. 124. [Couches de Saint Cassian, Italy] (Carnian) <HT: Thiel coll. K.29>.

Genus *Batheritiaris* VADET, 1999c

Batheritiaris VADET, 1999c: 89. Type-species: *Hemipedina incipiens* BATHER, 1909. [Southern Europe] (Carnian, Triassic).

Family Serpianotiaridae HAGDORN, 1995

Genus *Zardinitiaris* VADET, 1999c

Zardinitiaris VADET, 1999a: 90. Type-species: *Cidaris flexuosa* MÜNSTER, 1841. [Southern Europe] (Carnian, Triassic).

Order Triadocidaroida VADET, 1999c

Triadocidaroida VADET, 1999c: 93. {included families: Triadocidaridae SMITH, 1994; Braunechinidae VADET, 1999c}.

Family Triadocidaridae SMITH, 1994

Triadocidaridae SMITH, 1994: 186-187. Type-genus: *Triadocidaris* DÖDERLEIN, 1887. {other taxa included: *Zardinechinus* KIER, 1977; *Levicidaris* KIER, 1977; *Megaporocidaris* KIER, 1977; *Mikrocidaris* DÖDERLEIN, 1887; *Parvicidaris* SMITH, 1994; *Paurocidaris* KIER, 1977; *Vinchuscanchaia* SMITH, 1994 and “*Miocidaris*” *adrianae* ZARDINI, 1973}.

Genus *Vinchuscanchaia* SMITH, 1994

Vinchuscanchaia SMITH, 1994: 193. Type-species: *Vinchuscanchaia kieri* SMITH, 1994. [Peru] (Late Triassic/Early Jurassic).

Triassic

V. kieri SMITH, 1994: 193-194; pl. 6, figs 1-7. [Vinchuscancha, Peru] (?Late Triassic or Late Sinemurian) <HT: USNM 398508; PT: USNM 398521, 465263>.

Genus *Parvicidaris* SMITH, 1994

Parvicidaris SMITH, 1994: 194-195. Type-species: *Parvicidaris microapicalis* SMITH, 1994. [Peru] (Late Triassic).

Triassic

P. microapicalis SMITH, 1994: 195-196; text-fig. 6; pl. 2, figs 1-4. [Vinchuscancha, Peru] (?Late Triassic) <HT: USNM 465266>.

Family Braunechinidae VADET, 1999c

Braunechinidae VADET, 1999c: 99. Type-genus: *Braunechinus* VADET, 1999c. {other genera included: *Zardinechinus* KIER, 1977; *Mikrocidaris* DÖDERLEIN, 1887; *Wissmannechinus* VADET, 1999c}.

Genus *Braunechinus* VADET, 1999c

Braunechinus VADET, 1999c: 104. Type-species: *Cidaris dorsata* BRAUN in WISSMANN & MÜNSTER, 1841. [Southern Europe] (Anisian-Carnian) {other species included: *Cidaris waechteri* MÜNSTER, 1841; *Cidaris similis* DESOR, 1855}.

Genus *Wissmannechinus* VADET, 1999c

Wissmannechinus VADET, 1999c: 102-103. Type-species: *Cidaris haussmanni* WISSMANN, 1841. [Europe] (Ladinian-Carnian).

Order Paurocidaroida VADET, 1999c

Paurocidaroida VADET, 1999c: 111. {included families: Paurocidaridae VADET, 1999c}.

Family Paurocidaridae VADET, 1999c

Paurocidaridae VADET, 1999c: 111-112. Type-genus: *Paurocidaris* KIER, 1977. (Anisian-Carnian).

Order Orthopsida MORTENSEN, 1942

Family Orthopsidae DUNCAN, 1889

Genus *Orthopsis* COTTEAU, 1864

Jurassic

O. willei VADET & WILLE, 2002: 11-12; 3 figs [Cesareda, Portugal] (*bimammatum* Zone “Lusitanian”, Late Oxforian) <HT: Wille coll. 1051>.

Order undetermined

Family undetermined

Genus undetermined

Triassic

Genus ? thieli VADET, 1999c: 92-93; fig. 132-133. [Couches de Saint Cassian, Italy] (Carnian) <HT: Thiel coll.> {taxon apparently valid according to ICZN 4th ed., Article 11.9.3; no specimen number given}.

Superorder Gnathostomata ZITTEL, 1879

Order Holoctypoida DUNCAN, 1889

Suborder Holoctypoina DUNCAN, 1889

Family Holoctypoidae LAMBERT, 1899

Genus *Amblypygus* AGASSIZ, 1840

Placed in a separate family (Amblypygidae) and order (Amblypygoidea) by SRIVASTAVA (1988: 150).

Eocene

- A. moriensis* SRIVASTAVA & SINGH, 2001: 28-30; pl. 2, figs 4-7; pl. 3, figs 5-7. [Mori village, district Kachchh, Gujarat, India] (*Asterocyclina alticostata*-zone, Middle Eocene) <HT: DGUL KTE 259; PT: DGUL KTE 260, KTE 263, KTE 264>.
- [*A. moriensis* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <types unknown> {nomen nudum}.
- [*A. ratiparensis* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <types unknown> {nomen nudum}.

Subgenus *Paramblypygus* TESSIER & ROMAN, 1973

Paramblypygus TESSIER & ROMAN, 1973. Type-species: *Amblypygus (Paramblypygus) houghoueti* TESSIER & ROMAN, 1973. [Côte d'Ivoire] (Paleocene).

Paleocene

- A. (P.) houghoueti* TESSIER & ROMAN, 1973: 141-146; figs 1-6; pl. 1, figs 1-3; pl. 3, fig. 1. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Thanetian, Paleocene) <HT: IPM 1972-9Ac>.

Genus *Coenholectypus* POMEL, 1883

SMITH & WRIGHT (1999: 345) place this genus into its own subfamily (Coenholectypinae) within the family Holectypidae.

Aptian

- C. hideshimensis* TANAKA in TANAKA & OBATA, 1982: 122-124; text-figs 2b, 3; pl. 1, figs 3a-c. [Hiraiga Fm., Hideshima, Miyako City, Iwate Prefecture, Honshu, Japan] (Upper Aptian) <HT: GSJ F6165>.

Genus *Coptodiscus* COTTEAU & GAUTIER, 1895

Maastrichtian

- C. magniproctus* SMITH, 1995: 183-185; fig. 44, 45; pl. 19: figs 4-7. [Jebel Rawdah, Oman/United Arabian Emirates Border Region] <HT: BMNH EE3716; PT: BMNH EE3715>.

Genus *Globoholectypus* AZIZ, 1991

Globoholectypus AZIZ, 1991: 18-19. Type-species: *Globoholectypus globus* AZIZ, 1991. [Southern India] (Early Cretaceous).

Early Cretaceous

G. globus AZIZ, 1991: 19-20; figs 2a, pl.2, figs 1-5. [Dalmiapuram Formation, Trichinopoly Subbasin, Southern India] (Late Aptian to Early Albian, Early Cretaceous) <HT: MACSG 1947; PT: MACSG 1950>.

Genus *Holectypus* DESOR, 1842

Early Cretaceous

H. khelilense VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007: 29-30; pl. 15: fig. 1. [falaise, N of Safi, Morocco] (Late Berriasian to Early Valanginian) <Type series: Reboul coll. 216, Vadet coll. V7509 to 75212 [sic!, probably typo for 7521], 7523>.

Jurassic

H. depressoides GERASIMOV, 1955: 21; pl. 5: figs 15a-г. [Border river near Moskva, Gzhel'; Nikitino, Oka River, USSR] (Middle Callovian) <HT: GMM 850>.

H. phelani KIER, 1972a: 59-61; fig. 27; pl. 39, figs 1-5; pl. 40, figs 1-5; pl. 41, figs 1-3. [Upper Dhurma Fm., localities KK9-108-114, KK9-95-97 and L-926, Saudi Arabia] (Callovian) <HT: USNM 170437; PT: USNM 170438-170441>.

H. pictaviense NICOLLEAU & VADET, 1995: 58-59; 2 figs on p. 58; pl. 29: figs 1A-C. [Marnes à spongiares, environs de Niort, and environs de Poitiers, Poitou, France] (*berrense* Subzone, *bimammatum* Zone, Late Oxfordian) <HT: Philippe Nicolleau coll. no. 51a; PT: Philippe Nicolleau coll. no. 558, 2659, 2660>.

Genus *Philolectypus* VADET, 1997

Philolectypus VADET, 1997: 22. Type-species: *Holectypus sarthacensis* COTTEAU in DAVOUST, 1856. (Middle Jurassic).

Subfamily Coenholectypinae SMITH & WRIGHT, 1999

Coenholectypinae SMITH & WRIGHT, 1999: 345. Type-genus: *Coenholectypus* POMEL, 1883. {other genera included: *Lanieria* DUNCAN, 1889; *Coptodiscus* COTTEAU & GAUTHIER, 1895}.

Family Anorthopygidae WAGNER & DURHAM, 1966

Genus *Anorthopygus* COTTEAU, 1869

Cenomanian

A. arabicus ALI, 1990: 111-113; figs 6 (1-5). [Maaddud Fm., Wadi Kated al Qa's, Ras al Kaimah, United Arab Emirates] (Cenomanian) <HT: MGD-UAA > {no specimen no. given}.

Family Discoididae LAMBERT, 1899

JENSEN (1982: 94) placed this family into the order Echinoneoidea JENSEN, 1982.

Genus *Camerogalerus* QUENSTEDT, 1873

SMITH, GALLEMÍ, JEFFERY, ERNST & WARD (1999: 97) placed this genus in the family Holecypoidae Lambert, 1900.

Maastrichtian

C. cantabricus SMITH & GALLEMÍ *in* SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 97; figs 8a-c; pl. 2, figs 9-11. [Santander, Spain] <HT: BMNH EE6132; PT: MGB 37577>.

Genus *Discoides* PARKINSON, 1811

Early Cretaceous

D. cordobai BUITRÓN, 1971: 22-24; pl. 4, figs 6, 8; pl. 5, figs 1-3. [Lágrima Formation, Sierra de la Ranchera, Chihuahua, Mexico] (Middle Albian, Early Cretaceous) <HT: IGMUC 2257; PT: IGMUC 2254-2256>.

Suborder Echinoneina H.L. CLARK, 1925

Family Conulidae LAMBERT, 1911

JENSEN (1982: 94) placed this family into the order Echinoneoidea JENSEN, 1982.

Genus *Adelopneustes* GAUTHIER, 1889

In the *Treatise* WAGNER & DURHAM (1966: U447) considered this genus as synonym of *Galerites* LAMARCK, 1916.

Paleocene

A. ernsti SMITH & GALLEMÍ in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 100; fig. 9a-e; pl. 2, figs 5-8. [Casas de Oraien, Larumbe, Navarra province, Spain] (Upper Thanetian) <HT: BMNH EE6134>.

Genus *Conulus* LESKE, 1778

Late Cretaceous

[*C. akdjarensis* EGOROV, 1972]: 60. [Tadzhikistan] (Maastrichtian, Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

C. angulatus TZANKOV, 1982: 71; pl. 30, figs 2, 2a-b, 3. [Kreta, de Pleven, Bulgaria] <HT: USC CR, 1274>.

[*C. arcus* EGOROV, 1972]: 60. [Tadzhikistan] (Maastrichtian, Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

C. cookei BUITRON, 1974: 11-12; pl. 1, figs 4, 5, 7. [Ocozocuautila, W of Tuxtla Gutierrez, Chiapas, Mexico] (Late Cretaceous) <HT: IGMUC 2544; PT: IGMUC 2545>.

C. djanelidzei GONGADZE, 1972: 231-232; pl. 1: figs 10-12. [Мтавари [Mtavari] Formation, Kutai Region, Georgia] (Coniacian to Santonian, Late Cretaceous) <HT: MGI-ANGSSR 1/2683> {re-described in GONGADZE (1979: 68-70; pl. 5, figs 1a-d)}.

C. kubatliensis MELIKOV, in ALI-ZADE, 1988: 188-189; figs 9-11; pl. 2a, figs 2a-b. [Kubatly, Lower Caucasus, Azerbaijan] (Early Campanian) <HT: AzINEFTEKHIM 201/6>.

C. lamberti BUITRON, 1975: 12-13; pl. 1, fig. 6; pl. 2, figs 1-4. [Ocozocuautila, W of Tuxtla Gutierrez, Chiapas, Mexico] (Late Cretaceous) <HT: IGMUC 2546>.

C. sinensis MU & WU, 1976: 360; pl. 1: figs 1-7. [Zongshan Fm., Duila, Yadong County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27119, 27120 (repository not given; presumably NIGP)>.

Genus *Globator* L. AGASSIZ, 1840

Late Cretaceous

G. minimus MU & WU, 1976: 360; pl. 1: figs 8-11. [Zongshan Fm., Duila, Yadong County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <HT (by monotypy): 27121 (repository not given; presumably NIGP)>.

Genus *Pygopyrina* POMEL, 1883

Oxfordian

P. thibaudi VADET & NICOLLEAU, 1995: 84-88; figs 30-31; pl. 3, figs 3-4. [Oolithe à Dicerat, Dompcevrin, Paris Basin, France] (Middle Oxfordian) <HT: Nicolleau coll. 1257>.

Genus *Pyrina* DESMOULINS, 1835

Santonian

P. sphaerica TZANKOV, 1984: 74; pl. 31, figs 3a-c. [Choumen, Bulgaria] <HT: USC CR₂ 1278>.

Albian

P. azemati DEVRIÈS, 1972: 54-55; pl. 1: figs 11-14. [Sierra de Crevilente, Province Alicante, Spain] (Albian, Late Cretaceous) <ST: coll. J. Azema, Paris, nos. 939, 12, 770>.

Mid-Cretaceous

P. neolaevis SZÖRÉNYI, 1955: 38, 195; pl. 5, figs 18-22. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/168>.

Family Galeritidae GRAY, 1825

JENSEN (1982: 94) placed this family into the order Echinoneoidea JENSEN, 1982.

Genus *Galerites* LAMARCK, 1801

Cretaceous

- G. chovaresmicus* TRAUTSCHOLD, 1859: 309-310; pl. 4, figs 3a-c. [Coast of the Aral Lake, Kazakhstan/Uzbekistan] (Cretaceous) <not given>.
- G. (Galerites) vulgaris minor* SCHULZ, 1985: 50; pl. 5, fig. 1; pl. 10, figs 6; pl. 13, figs 3-8. [Lüneburg and Krons Moor, Northwestern Germany] (middle *lanceolata*-zone to *obtusa*-zone, early Early Maastrichtian) <SNLFB kma358>.

Subgenus *Pironaster* MUNIER-CHALMAS, 1890

Cretaceous

- G. (Pironaster) ernsti* SCHULZ, 1985: 42-43; pl. 1, figs 1-2; pl. 11, figs 5-6. [Lägerdorf bei Itzehoe, Höver and Misburg, Northwestern Germany] (Campanian) <SNLFB kca52>.
- G. (Pironaster) schmidi* SCHULZ, 1985: 45-46; pl. 3, figs 1-2; pl. 11, figs 11-14. [Zeltberg quarry, Lüneburg, Northwestern Germany] (*obtusa*-zone, early Early Maastrichtian) <SNLFB kma358>.

Family Neoglobatoridae Endelman, 1980a

Neoglobatoridae ENDELMAN, 1980a: 96. Type-genus: *Neoglobator* ENDELMAN, 1980. [Russia and Eastern Europe] (Paleogene). {other genera included: ? *Galeraster* COTTEAU, 1890}.

Genus *Neoglobator* ENDELMAN, 1980a

Neoglobator ENDELMAN, 1980a: 96-97. Type-species: *Neoglobator panteleevi* ENDELMAN, 1980a. [Kazakhstan] (Paleocene) {other species included: *Pseudopyrina subovalis* RAVN, 1927; *Pseudopyrina subcircularis* RAVN, 1927; *Globator ravni* NIELSEN in RAVN, 1927; *Pygorhynchus ovalis* SMISER, 1935; *Pyrina houzeau* COTTEAU, 1875; *Pyrina montainvillensis* SORIGNET, 1850; genus placed into the synonymy of *Adelopenustes* GAUTHIER, 1889 by SMITH & JEFFERY (2001: 148)}.

Paleogene

- N. akkajensis* ENDELMAN, 1980b: 99-103; figs 1B; pl. 1, figs 5a-f, 6. [central Crimea, near Bilogursk, Ak-Kaya Mt. Ukraine] (Lower Eocene) <HT: PIN 3501/57>.

- N. danicus* ENDELMAN, 1980b: 93-97; figs 1a; pl. 1, figs 1a-Г, 2a-B, 3a-B. [Sarytasch, North Akatau, Mangyshlak, Kazakhstan] (Danian) <HT: PIN 3500/694>.
- N. insolitus* ENDELMAN, 1980b: 97-99; figs 1б, 2a-г; pl. 1, figs 4a-г. [near Baysarly, Southern Mangyshlak, Kazakhstan] (Lower Paleocene) <HT: PIN 3500/333>.
- N. panteleevi* ENDELMAN, 1980a: 97-103; figs 2a-i; pl. 1, figs 1-6. [near Usak, North Aktau, Mangyshlak, Kazakhstan] (Lower Paleocene) <HT: PIN 3500/621> {*N. panteleevi* ENDELMAN, 1980a was considered a junior synonym of *Adelopneustes montainvillensis* (SORIGNET, 1850) by SMITH & JEFFERY (2001: 149-150)}.

Suborder Conoclypina HAECKEL, 1896

Family Conoclypidae ZITTEL, 1879

Genus *Conoclypus* AGASSIZ, 1839

Oligocene

- C. mittereri* SCHLOSSER, 1923: 258-259. [Häring, Tyrolia, Austria] (Early Oligocene) <not given>.

Family Oligopygidae DUNCAN, 1889

KIER (1967) established a separate order for the oligopygid echinoids.

Genus *Haimea* MICHELIN, 1851

Eocene

- H. bajasurensis* SQUIRES & DEMETRION, 1994: 847-849; figs 2.1-2.9. [Tepetate Fm. and Bataque Fm., Baja California Sur, Mexico] (Middle Lower Eocene) <HT: IGM 5934 (=LACMIP 12121); PT: IGM 5918 (=LACMIP 12120), IGM 5935 – 5937 (=LACMIP 12122 – 12124)>.
- H. rojasi* ŽITŤ, 1993: 30-36; pl. 8, figs 1-3. [Alcalá y Báguanos, Holguín, Cuba] (Eocene) <HT: PIASC 6915>.

Genus *Ovulechinus* LAMBERT, 1918

Late Cretaceous

O. yadongensis MU & WU, 1976: 362; pl. 2: figs 8-14. [Zongshan Fm., Duila, Yadong County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27126, 27127 (repository not given; presumably NIGP)>.

Subgenus *Pseudovulechinus* SZÖRÉNYI, 1955

Pseudovulechinus SZÖRÉNYI, 1955: 79, 217-218. Type-species: *Ovulechinus* (*Pseudovulechinus*) *rotundatus* SZÖRÉNYI, 1955. [Eastern Europe] (Mid-Cretaceous).

Mid-Cretaceous

O. (P.) rotundatus SZÖRÉNYI, 1955: 79-80, 218; pl. 11, figs 7-11. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/399>.

Order Echinoneoidea JENSEN, 1982

Echinoneoidea JENSEN, 1982: 94. {included families: Echinoneidae AGASSIZ & DESOR, 1847, Galeritidae GRAY, 1825, Conulidae LAMBERT, 1911, Discoididae LAMBERT, 1899}.

Order Amblypygoidea SRIVASTAVA, 1988

Amblypygoidea SRIVASTAVA, 1988: 150. Type-family: Amblypygidae SRIVASTAVA, 1988.

Family Amblypygidae SRIVASTAVA, 1988

Amblypygidae SRIVASTAVA, 1988: 150. Type-genus: *Amblypygus* AGASSIZ, 1840.

Family [Kutchypygidae SRIVASTAVA, 1988]

[Kutchypygidae SRIVASTAVA, 1988]: 150. Type-genus: [*Kutchypygus* SRIVASTAVA, 1988]. {Nomen nudum; text refers to an unpublished paper, which apparently was not published subsequently}.

Genus [*Kutchypygus* SRIVASTAVA, 1988]

[*Kutchypygus* SRIVASTAVA, 1988]:150. {Nomen nudum, no type species mentioned, no description provided; text refers to an unpublished paper, which apparently was not published subsequently; on p. 151 a *Kutchypygus bermotensis* SRIVASTAVA is mentioned}.

Eocene

[*K. bermotensis* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <unknown> {nomen nudum}.

Superorder Mesostomata VADET, 1997

Mesostomata VADET, 1997: 136, 138. {included family: Desorellidae VADET, 1997}.

Family Desorellidae LAMBERT, 1911

= Desorellidae VADET, 1997: 136, 138. Type-genus: *Desorella* COTTEAU, 1855. {included genera: *Pachyclypus* DESOR, 1856; indicated as new in VADET (1997: 136), but was established in 1911 by LAMBERT (1911: 30) already}.

Superorder Sarhostomata VADET, 1999b

Sarhostomata VADET, 1999b: 87. {included genera: *Pyrinodia* POMEL, 1883}.

Suborder Uncertain

Genus *Echinogalerus* KÖNIG, 1825

Paleocene

E. raingeardi TESSIER & ROMAN, 1973: 147-149; fig. 7; pl. 1, figs 4-7. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Paleocene, Thanetian) <HT: IPM 1972-9B>.

Superorder Neognathostomata Smith, 1981

Cohort Irregularia LATREILLE, 1825

Order Clypeasteroidea A. AGASSIZ, 1872

Suborder Clypeasterina A. AGASSIZ, 1872

Family Clypeasteridae L. AGASSIZ, 1835

Genus *Clypeaster* LAMBERT, 1801

Recent

- C. isolatus* SERAFY, 1971: 166-168; figs 1a-e, 2a-j, 3a-c. [R.V. Anton Brunn, Cruise 12, Stat. MV-65-IV-37, off San Felix Island, SE Pacific, 26°16' S, 80°06' W, depth 75 m] (Recent) <HT: USNM E11361; PT: USNM, MCZ, BMNH, AM nos. not given (60 specimens)>.
- C. kieri* PAWSON & PHELAN, 1979: 796-800; figs 1A-F. [International Indian Ocean Expedition, Anton Bruun Cruise 4B, Stat. 202A (17°25' N, 71°39' E to 17°21' N, 71°41' E, depth 96-106 m), Stat. 202B (17°41' N, 71°33' E to 17°45' N, 71°32' E, depth 90 m), Stat. 202C (18°27' N, 71°13' E to 18°38' N, 71°09' E, depth: 84-97 m), off Bombay, Indian Ocean] (Recent) <HT: USNM E18188 (Anton Bruun Cruise 4B Stat. 202A); PT: USNM E 18189-18191>.

Pleistocene

- C. chiangchunshanensis* WANG, 1985: 163-167; fig. 3e, 10; pl. 3, fig. 2; pl. 4, fig. 3; pl. 6, figs 1a-b. [Toukoshan Fm., Chiangchunshan, Miaoli-hsien, northern Taiwan] (Late Pleistocene) <HT: NTUG-(E)-0076>.
- C. paishatunensis* WANG, 1985: 167-169; fig. 3d, 11; pl. 2, figs 3a-c. [Toukoshan Fm., Paishatun, Miaoli-hsien, northern Taiwan] (Late Pleistocene) <HT: NTUG-(E)-0095>.

Pliocene

- C. maanliaensis* WANG, 1985: 161-163; fig. 3c, 8; pl. 4, fig. 4; pl. 8, figs 1a-b. [Pitoushan Fm., Maanliao, Taichung-hsien, central Taiwan] (Late Pliocene) <HT: NTUG-(E)-0073>.
- C. maoadentroensis* KIER, 1992: 17-18; pl. 4, figs 5-7; text-figs 3, 4a. [Mao Adentro Limestone, locality NMB 17022, Rio Cana Section, Dominican Republic] (Early Pliocene) <HT: NMB M9746>.
- C. pitouensis* WANG, 1985: 169-174; fig. 3g, 13, 14; pl. 7, figs 1-3; pl. 8, figs 2a-b, 3a-b. [Erhchiu Fm., Pitou-chiao, Taipei-hsien, northern Taiwan] (Early Pliocene) <HT: NTUG-(E)-0095>.

Miocene

- C. butleri* McNAMARA & KENDRICK, 1994: 44-46; figs 17A-D. [Poivre Fm., "The Ledge" Barrow Island, Western Australia] (Middle Miocene) <HT: WAM 75.970; PT: WAM 75.972, 75.973, 82.271, 82.330, 94.428, 94.430, 82.412>.

- C. dondolii* FISCHER, 1985: 201; fig. 5/1; pl. 3, figs 1-2, 8. [Turrúcares Fm., Pendiente, SW del Cerro Candelaria, Costa Rica] (Middle Miocene) <HT: ECG CO-93>.
- C. gombosae* MIHÁLY, 1990a: 238-239; pl. 2: fig. 2; pl. 3; fig. 1; pl. 4: fig. 1. [Sámsonhaza Fm., Gomb Hill, Kemence, Hungary] (Lower Badenian {=Langhian}) <HT: MÁFI Coll. Ech. 340>.
- C. hothiensis* Srivastava. [India] (*Ostrea latimarginata* Zone, Miocene) <types unknown> {nomen nudum; mentioned in SRIVASTAVA (1988: 153)}.
- C. kurangaensis* JAIN, 2002: 123-124; pl. 5, figs 3-4. [Gaj Fm., Kuranga Railway Station, Kathiawar, Gujarat, India] (Early Miocene) <HT: GSI 20764; PT: GSI 20765>.
- C. papilionensis* KOTCHETOFF, KOTCHETOFF & VEIGA FERREIRA, 1975: 75-78; [Penedo cliff, north of Cap d'Espichel, Portugal] (Tortonian, Miocene) <not given>.
- C. tauricus* DESOR var. *elatiore* MARCOPOULOS-DIACANTONI, 1972: 149-150. [Stafidokefala, south-west of Agios Georgios Sitias, Lsithi Prefecture, Crete Island, Greece] ("Helvetian") <ST: 1972/12, 1972/13, repository unknown>.
- C. paraensis* BRITO, 1979: 734; pl. 1, figs 1-3. [Pirabas Fm., Ilha de Fortaleza, baía de Pirabas, Pará, Brazil] <HT: MN 5382-I>.
- C. yeni* WANG, 1985: 155-161; fig. 3f, 5, 6; pl. 4, fig. 2; pl. 5, figs 1a-b. [Kuanyinshan Sandstone, Wu-chi River, near Kantsulin, Kuohsing-hsiang, Nantou-hsien, central Taiwan] (Middle Miocene) <HT: CGS-E-83024>.
- C. yingkoensis* WANG, 1985: 151-155; fig. 3b, 4; pl. 3, figs 1a-c; pl. 4, fig. 1. [Nankang Sandstone, Yingko, Taipei-hsien, northern Taiwan] (Middle Miocene) <HT: NTUG-(E)-0073>.

Family Fossulasteridae PHILIP & FOSTER, 1971

Fossulasteridae PHILIP & FOSTER, 1971: 681-682. Type-genus: *Fossulaster* LAMBERT & THIÉRY, 1925. {included genera: *Scutellinoides* DURHAM, 1955; *Willungaster* PHILIP & FOSTER, 1971}.

Genus *Fossulaster* LAMBERT & THIÉRY, 1925

Miocene

- F. changi* WANG, 1994: 234-235; figs 11A-D; pl. 5, figs 1-2. [Morgan Limestone, Murray Cliffs, South Australia] (Miocene) <HT: YPM 33657; PT: YPM 33646>.
- F. durhami* WANG, 1994: 233-234; figs 8A-B, 9, 10; pl. 4, figs 2a-c. [Morgan Limestone, Murray Cliffs, South Australia] (Miocene) <HT: YPM 33645>.

F. exiguus PHILIP & FOSTER, 1971: 686-687; text-fig. 7; pl. 128: fig.3; pl. 132: figs 2, 3, 6, 8, 9, 11. [Melton Limestone, Tickera, Yorke Peninsula, South Australia] (Longfordian, Lower Miocene) <HT: NMV P27952; PT: NMV P27949, UNE 11759, 12002-05>.

Genus *Orbispala* IRWIN, 1995

Orbispala IRWIN, 1995: 189-191. Type-species: *Orbispala occultoforma* IRWIN, 1995. [Australia] (Miocene).

Miocene

O. occultoforma IRWIN, 1995: 191-194; figs 3A-M. [Zeally Limestone Mb., Puebla Fm.; Cliffs of Jan Juc to Torquay, Victoria, South Australia] (Late Early Miocene) <HT: NMV P139064; PT: NMV P139065-139080>.

Genus *Philipaster* WANG, 1994

Philipaster WANG, 1994: 227. Type-species: *Scutellina morgani* COTTEAU, 1891. [Australia] (Late Oligocene)

Genus *Prowillungaster* WANG, 1994

Prowillungaster WANG, 1994: 229. Type-species: *Prowillungaster major* WANG, 1994. [South Australia] (Oligocene-Miocene).

Oligocene-Miocene

P. major WANG, 1994: 229; figs 5A-D; pl. 2, figs 1-4; pl. 3, figs 1-2. [Mannum Fm., Murray Cliffs, South Australia and Point Addis Limestone, Airey's Inlet, Victoria, Australia] (Oligocene and Early Miocene) <HT: YPM 33662; PT: YPM 33663, 33652, 33648, 33653, 33654>.

P. minor WANG, 1994: 229-231; figs 5E-F; pl. 3, figs 3a-b. [Mount Gambier Limestone, Marte, Mt. Gambier, South Australia] (Miocene) <HT: YPM 33655; PT: YPM 33656>.

Genus *Willungaster* PHILIP & FOSTER, 1971

Willungaster PHILIP & FOSTER, 1971: 687-688. Type-species: *Willungaster scutellaris* PHILIP & FOSTER, 1971. [South Australia] (Oligocene-Miocene).

Oligocene-Miocene

W. scutellaris PHILIP & FOSTER, 1971: 688-689; text-fig. 8; pl. 127: fig. 2; pl. 131: figs 1-5, 7; pl. 133: figs 7, 9. [Port Vincent Limestone, Edithburg, Yorke Peninsula, South Australia] (Janjukian, Upper Oligocene of Longfordian, Lower Miocene) <HT: NMV P27951; PT: UNE 11765-67, 12006-08>.

Family Scutellinoididae IRWIN, 1995

Scutellinoididae IRWIN, 1995: 194. Type-genus: *Scutellinoides* DURHAM, 1955.

Order Laganoida JENSEN, 1982

Laganoida JENSEN, 1982: 95. {included families: Laganidae A. AGASSIZ, 1873, Astri-clypeidae STEFANINI, 1911, Rotulidae GRAY, 1855, Mellitidae STEFANINI, 1911, Dendrasteridae LAMBERT, 1889, Echinarachniidae LAMBERT, 1914, Fibulariidae GRAY, 1855}

Suborder Laganina MORTENSEN, 1948

Family Neolaganidae DURHAM, 1954

Genus *Tetradiella* LIAO & LIN, 1981

Tetradiella LIAO & LIN, 1981: 482. Type-species: *Tetradiella sinica* LIAO & LIN, 1981. [China] (Pliocene).

Pliocene

T. sinica LIAO & LIN, 1981: 482-283; figs 1-2; pl. 1, figs 1-7. [Beibuwan, Guangxi Province, China] (Pliocene) <no holotype defined and no repository given; specimens located at NIGP>.

Family Fibulariidae GRAY, 1855

JENSEN (1982: 95) placed this family into the order Laganoida JENSEN, 1982.

Genus *Cenofibula* GÄSSER, 1994

Cenofibula GÄSSER, 1994: 13-14. Types-species: *Cenofibula castriauxiensis* GÄSSER, 1994.

Eocene

C. castriauxiensis GÄSSER, 1994: 14-15; figs 2a-c, 3. [Castellolí, near Igualada, Barcelona Province, Spain] (Bartonian) <HT: MGSB 55225.a>.

Genus *Echinocyamus* VAN PHELSUM, 1774

Recent

E. insularis MIRONOV & SAGAIDACHNY, 1984: 192-193; fig. 6: 1a.c; pl. 1, fig. 4. [Dimitri Mendeleev Station 582 (27°07' S, 109°26' W, depth 50 m), 589 (26°27' S, 105°29' W, depth 60-80 m); Isle Paschi and Sala-y-Gomez, Southeastern Pacific Ocean] <HT: IOANSSR N XV-65-19>.

E. insularis macroproctus MIRONOV & SAGAIDACHNY, 1984: 193-194; fig. 6: 2a-b. [Dimitri Mendeleev Station 1256 (29°30' S, 167°52' E, depth 50-66 m); Isle Norfolk, Southwestern Pacific Ocean] <HT: IOANSSR N XV-65-20>.

Oligocene

E. nummulicus bernaniensis SRIVASTAVA, 1978: 423-424; pl. 1: figs 6-7. [about 1.6 km N 65° E of Ber Mota (23°27'45" N, 68°38'25" E), Kutch, India] (*Nummulites subclipeus*-zone, Ber Nani Stage, Middle Oligocene) <HT: DGUL K1145> {raised to species level by SRIVASTAVA (1988: 152)}.

E. raoi SRIVASTAVA, 1978: 424-426; pl. 1: figs 8-9. [about 1.6 km N 65° E of Ber Mota (23°27'45" N, 68°38'25" E), Kutch, India] (*Nummulites subclipeus*-zone, Ber Nani Stage, Middle Oligocene) <HT: DGUL K1204>.

E. wilsoni KIER, 1997: 8; figs 4A-E. [Belgrade Fm., Belgrade quarry, North Carolina, USA] (Late Oligocene) <HT: USNM 398476; PT: USNM 398477-398479; 492097-492100>.

Eocene

E. gurnahensis ROMAN & STROUGO, 1994: 34-37; fig. 5; pl. 1, figs 3-5; pl. 3, figs 4-7, 13. [Thebes Fm., Gebel Gurnah, Egypt] (Ypresian) <HT: MNHN R10688>.

E. hungaricus SZÖRÉNYI, 1952: 289-290, 291-292; figs 1-8, 13-14. [Gánt, Zámoly, Hungary] (Early Eocene) <ST: MÁFI Ech-338 according to BODA, 1964>.

- E. jaisalmerensis* SRIVASTAVA & MATHUR, 1996: 54-55; fig. 3; pl. 1, figs 1-9. [Bandah Fm., near Ramgarth, Jaisalmer district, Rajasthan, India] (Middle Eocene) <HT: GSI 20684>.
- E. pannonicus* SZÖRÉNYI, 1952: 290-291, 292-293; figs 9-12. [Gánt, Zámoly, Hungary] (Early Eocene) <ST: MÁFI Ech-337 according to BODA, 1964>.

Subgenus *Lepidocyamus* MIRONOV & SAGAI DACHNY, 1984

- Lepidocyamus* MIRONOV & SAGAI DACHNY, 1984: 194. Type-species: *Echinocyamus crispus* MAZZETTI, 1893. [Indic Ocean and Red Sea] (Recent).

Genus *Fibularia* LAMARCK, 1816

Recent

- F. japonica* SHIGEI, 1982: 11-15; figs 1-48. [off Misaki Marine Biological Station, Sagami Bay (sublittoral); 3 km off Futamachiya (35°06.5' N, 139°35.0' E, depth ca. 70 m), Sagami Bay; east of Iki Island (33°45.1' N, 129°57.5' E, depth 45 m); off Tomioka (Amakusa), W. Kyushu (ca. 30 m depth); off Oki Islands (between Dogo and Dozen, depth ca. 60 m); Suruga Bay (sublittoral zone); off Boso Peninsula (34°51.2' N, 139°55.6' E, 34°51.1' N, 139°55.2' E, depth 100 m); Japan] (Recent) <HT: MMBS Fib. 1 (from Sagami Bay)>.

Pliocene

- F. gassusensis* ELATTAAR, 2001b: 88-91; figs 5A1-3, 7A-F; pl. 2, figs 7-10. [Sharm El Arab Member, Shagra Formation, Wadi Gassus, 10 km SW Safaga, Red Sea coast, Egypt] (Late Pliocene) <HT: AUSGM E (the specimen figured on pl. 2, figs 8-10, no number mentioned); PT: AUSGM E (the specimen figured on pl. 2, fig. 7, no number mentioned)>.
- F. wizrensis* ELATTAAR, 2001b: 86-87; figs 5C1-3; pl. 2, figs 1-3. [Sharm El Arab Member, Shagra Formation, Wadi Wizr, 41 km SW Quseir, Red Sea coast, Egypt] (Late Pliocene) <HT: AUSGM E (the specimen figured on pl. 2, figs 1-3, no number mentioned)> {based on a single specimen}.

Miocene

- F. damensis* KIER, 1972a: 90-91; pl. 59, figs 2-10; pl. 60, fig. 1. [Dam Fm., localities S-137, S-178, S-553, S-568 and S-1392, Saudi Arabia] (Miocene) <HT: USNM 170481; PT: USNM 170482-170484>.

F. depressa JAIN, 2002: 124-126; pl. 5, figs 5-6; pl. 7, fig. 2. [Gaj Fm., 0.5 km east of Lowrali, Kathiawar, Gujarat, India] (Middle Miocene) <HT: GSI 20769; PT: GSI 20767-68>.

F. guavarensis SRIVASTAVA, 1978: 426-427; pl. 1: figs 10-14. [about 2.5 km SW of Guvar (23°38'10" N, 68°32'30" E), Kutch, India] (*Ostrea gajensis*-zone, Lower Miocene) <HT: DGUL K1237> {mis-spelled *Fibularia guvarensis* in SRIVASTAVA (1988: 153)}.

Eocene

F. khargahensis AZAB & ELATTAAR, 1999: 850-853; tab. 7; pl. 1: figs 8-10. [Gebel Um El Ghanayem, ca. 3 km E of Ain Dabadib, and Gebel El Teir, Kharga area, Egypt] (*Linthia (Lutetiaster) cavernosa* Zone, Early Libyan, Early Eocene) <repository and specimen nos. not given>.

Genus *Marginoproctus* BUDIN, 1980

Marginoproctus BUDIN, 1980: 306. Type-species: *Marginoproctus djakonovi* BUDIN, 1980. [Northern Pacific] (Recent).

Recent

M. djakonovi BUDIN, 1980: 306-308; figs 1-8. [“Витязь”-cruise (“Vityaz”-cruise), Stat. 1856 (56°25' N, 143°19' E, depth 80 m), 5640 (44°4' N, 148°57' E, depth 800 m), “Жемчуг”-cruise (“Zhemchug”-cruise) Stat. 33 (54°35' N, 165°34' W, depth 212 m), 251 (51°47' N, 158°7' E, depth 60 m), 1947-1948 (50°03' N, 155°51' E; 50°14' N, 155°54' E; 50°44' N, 156°40' E; 49°52' N, 155°37' E, depth 70-300 m), East Kamchatka, Commander and Kurile Islands and Sea of Okhotsk] (Recent) <HT: IOANSSR XV-69-17 (“Vityaz”-cruise, Stat. 1856)>.

Genus *Mortonia* GRAY, 1852

Miocene

M. lowraliensis JAIN, 2002: 126-127; pl. 5, figs 13-23; pl. 7, figs 3-5. [Gaj Fm., 0.5 km east of Lowrali, Kathiawar, Gujarat, India] (Middle Miocene) <ST: GSI 20770-78>.

Genus *Scutellina* L. AGASSIZ, 1841

Cretaceous

S. supramarginalis ENGEL, 1976: 55-56; figs 1a-f. [Maastricht, The Netherlands] (Cretaceous or Eocene) <HT+PT: Mus. Teyler, Haarlem 16053>.

Infraorder Fibularina SMITH, 1984

Fibularina SMITH, 1984: 171, 173. Type-genus: *Fibularia* LAMARCK, 1816.

Family Laganidae A. AGASSIZ, 1873

JENSEN (1982: 95) placed this family into the order Laganoida JENSEN, 1982.

Genus *Peronella* GRAY, 1855

Pleistocene

- P. mai* WANG, 1982b: 150-154; pl. 3, figs 4-5, 7. [Cholan Fm., Chungkang River near Sanwan, Miaoli County, northern Taiwan] (Late Pliocene to Early Pleistocene) <HT: NTUG-[E]-0112>.
- P. lesueuri augusta* PLEDGE & SADLER, 1990: 103-104; figs 1a-b. [Port Augusta, South Australia] (Late Pleistocene) <HT: SAM P24854>.
- P. miaoliensis* WANG, 1982b: 154-155; pl. 4, figs 4-6, 8. [Toukoshan Fm., between Nansheh and Kungssuliao (Kôshiryô), Miaoli County, northern Taiwan] (Pleistocene) <HT: NTUG-[E]-0114>.
- P. toukoshanensis* WANG, 1982b: 149-150; pl. 3, figs 1-3, 6. [Toukoshan Fm., probably Miaoki County, northern Taiwan] (Pleistocene) <HT: NTUG-[E]-0243>.

Pliocene

- P. ova* McNAMARA, 1996: 195-196; figs 2A-J. [Roe Calcarenite, Roe Plains, Madura district, Western Australia] (Late Pliocene) <HT: WAM 94.854; PT: WAM 82.2103-82.2121, 82.2255, 82.2273-2283, 94.855, 94.856, 82.2095-6, 82.2135-2143, 82.2241-2248, 94.848-854, 82.2151, 85.1876>.

Miocene

- P. changchihkengensis* WANG, 1982b: 148-149; pl. 4, figs 1-3, 7. [Changchihkeng Fm., Changchihkeng (Chôsjiko), Chiayi County, southern Taiwan] (Late Miocene) <HT: NTUG-[E]-0111>.

Eocene

- P. tschimanica* PORETSKAYA, in AKOPJANA, 1974: 361-362; pl. 186, figs 2a-b; pl. 187, figs 4a-b. [Ciman, Armenia] (Late Eocene) <HT: LGU 306/3>.

Family unknown

Genus *Tridium* TANDON & SRIVASTAVA, 1980

Tridium TANDON & SRIVASTAVA, 1980: 1. Type-species: *Tridium kieri* TANDON & SRIVASTAVA, 1980. [India] (Eocene) {TANDON & SRIVASTAVA (1980: 1) placed this genus within the suborder Laganina MORTENSEN, 1948 in a new, unnamed family with affinities to the family Fibulariidae}.

Middle Eocene

T. kieri TANDON & SRIVASTAVA, 1980: 2-3; figs 1-2; pl. 1, figs 1-6. [Near Guvar, Kutch, India] (*Nummulites beaumonti* Zone, Middle Eocene) <HT: DGUL K651; PT: DGUL K652, K653>.

Suborder Scutellina HAECKEL, 1896

Superfamily Scutellidea SMITH, 1984

Scutellidea SMITH, 1984: 171, 173. Type-genus: *Scutella* LAMARCK, 1816.

Family Scutellidae Gray, 1825

Genus *Mennerella* SHMIDT in SHMIDT & SINYELNIKOVA, 1971

Mennerella SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 909 [EN: 72-73]. Type-species: *Mennerella ovata* SHMIDT in SHMIDT & SINYELNIKOVA, 1971. [western Kamchatka] (Miocene).

Miocene

M. ovata SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 909-910 [EN: 73]; pl. 1: figs 2a-6. [Upper Kakertok Suite, Kavran Series, Kavran River, Kavran-Utkholok Bay, western Kamchatka] (Middle to Late Miocene) <HT: CGM Kol. 10255 No. 5>.

Genus *Scutella* Lamarck, 1816

Miocene

S. aegyptiaca ALI, 1998: 543-544; fig. 3A. [Marmarica Fm., Gebel west El Migahhiz, Siwa Oasis, Egypt] (Langhian-Serravalian) <HT: GM-GDEMU, no specimen no.>.

- S. conica* ALI, 1998: 544; figs 3B-C. [Marmarica Fm., Gebel west El Migahhiz, Siwa Oasis, Egypt] (Langhian-Serravalian) <HT: GM-GDEMU, no specimen no.>.
- S. kalksburgensis* WIESBAUR, 1874: 164-165. [Kalksburg near Vienna, Austria] (Badenian {= Langhian-Serravalian}, Middle Miocene) <no specimen no.> {considered a junior subjective synonym of *Parascutella gibbercula* (DE SERRES, 1829) by KROH (2005: 90)}
- S. marchettii* TAVANI, 1939: 37-38; pl. 2 (4), fig. 13. [Sollum Egypt] (Miocene) <no specimen no.>.
- S. muelleri* MIHÁLY, 1985: 241, 260-261; pl. 3, figs 5-6; pl. 4, fig. 1. [Leitha Limestone, Gyakorló út [Street], 10th district, Budapest, Hungary] (Late Badenian (= Early Serravallian), Middle Miocene) <HT: MAFI Ech 358> {considered a junior subjective synonym of *Parascutella gibbercula* (DE SERRES, 1829) by KROH (2005: 91)}
- S. romani* MIHÁLY, 1985: 240-241, 260; pl. 2, figs 4-6. [Leitha Limestone, Kerepesti út [Street], 10th district, Budapest, Hungary] (Late Badenian (= Early Serravallian), Middle Miocene) <HT: MAFI Ech 1; PT: MAFI Ech 2> {considered a junior subjective synonym of *Parascutella gibbercula* (DE SERRES, 1829) by KROH (2005: 91)}.
- S. vindobonensis altus* MIHÁLY, 1990a: 237-238; pl. 1: figs 1-2; pl. 2: fig. 1. [Budapest XIV, Hungary] (Upper Badenian {=Serravalian}) <HT: MÁFI Coll. Ech. 400> {considered a junior subjective synonym of *Parascutella gibbercula* (DE SERRES, 1829) by KROH (2005: 90)}.

Family Protoscutellidae DURHAM, 1955

Genus *Protoscutella* STEFANINI, 1924

Eocene

- P. mississippiensis rosehillensis* KIER, 1980: 38-40; fig. 16; pl. 12, figs 1-5. [Castle Hayne Limestone, Rose Hill locality 11, locality 35, North Carolina, USA] (Middle Eocene) <HT: USNM 264062; PT: USNM 264061, 264063>.

Family Eoscutellidae DURHAM, 1955

Genus *Eoscutella* GRANT & HERTLEIN, 1938

Eocene

- E. mirandae* PARMA, 1985: 37-38; figs 3-4; pl. 1, figs 1-3. [Cerro Blanco, Santa Cruz Province, Argentina] (“Patagoniano”, Eocene?) <HT: CPBA 12902>.

Subgenus *Tigilella* SHMIDT, 1975

Eoscutella (*Tigilella*) SHMIDT, 1975: 30. Type-species: *E. (Tigilella) kamtschatica* SHMIDT, 1975. [Kamchatka] (Eocene).

Eocene

E. (Tigilella) kamtschatica SHMIDT, 1975: 30-31; un-numbered text-fig. on p. 30; pl. 5: figs a-e. [Kavran River, Kavran-Utkholok Bay, western Kamchatka] (Late Eocene) <HT: VNIGRI I/803>.

Family Dendrasteridae LAMBERT, 1889

Genus *Dendraster* L. AGASSIZ, 1847

Pliocene

D. sullivanii DURHAM & MORGAN, 1978: 303-305; figs 4a-g, 5. [UCMP loc. 3399, San Gregorio Beach, California] (Early Pliocene) <HT: UCMP 14450; PT: UCMP 14448, 14451-14456; CAS 58177-58179>.

Genus *Merriamaster* LAMBERT, 1911

Pliocene

M. weaveri DURHAM & MORGAN, 1978: 301-303; figs 3a-g. [UCMP loc. 3399, San Gregorio Beach, California] (Early Pliocene) <HT: UCMP 14259; PT: UCMP 14443-14447, 14449; CAS 58175-58176>.

Family Echinarachniidae LAMBERT, 1914

JENSEN (1982: 95) placed this family into the order Laganoida JENSEN, 1982.

Genus *Astrodapsis* CONRAD, 1856

Miocene

A. bajasurensis SQUIRES & DEMETRION, 1993: 259-262; fig. 2. [Isidro Fm., CSUN locality 1495, near mouth of Arroyo Mezquital, north central Baja California Sur, Mexico] (Middle Miocene) <HT: IGM 5926; PT: IGM 5927-5933, LACMIP 11595-11599>.

- A. ilyinensis* SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 911 [EN: 76]; pl. 1: figs 3a-b. [Il'inskaya [Ilinsk] Suite, Kavran Series, Nepropusk Point, Tochilo section, Grand Otel' area, western Kamchatka] (Early to Middle Miocene) <HT: CGM Kol. 10255 No. 28> {Re-described by SHMIDT (1984) and transferred to *Pseudoastrodaspis*}.

Genus *Echinarachnius* GRAY, 1825

Miocene

- E. astrodapsoides* WAGNER, 1974: 110; pl. 1: figs 5-8. [Unga Fm., Bear Lake, Alaska Peninsula] (Middle to Upper Miocene) <HT: USNM 181139; PT: USNM 181140, 181141, UA 2435, UCMP 14085>.
- E. plafkeri* WAGNER, 1974: 111-112; pl. 3: figs 5-7. [Yakataga Fm., Icy Point, Lituya District, eastern Gulf of Alaska] (Upper Miocene) <HT: USNM 181142; PT: USNM 181143>.
- E. ungaensis* WAGNER, 1974: 112-115; pl. 1: figs 9-11. [Bear Lake Fm., Bear Lake, Alaska Peninsula] (Middle or Upper Miocene) <HT: USNM 181144>.

Genus *Faassia* SHMIDT in SHMIDT & SINYELNIKOVA, 1971

- Faassia* [*Neoscutella*] SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 911 [EN: 73-76]. Type-species: *Faasia globosa* SHMIDT in SHMIDT & SINYELNIKOVA, 1971. [western Kamchatka] (Middle Miocene) {named *Neoscutella* in figure captions and table}.

Miocene

- F. globosa* SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 910 [EN: 76]; pl. 1: figs 1a-b. [Middle Etolon [Etolonsk] Suite, Kavran Series, Nepropusk Point, western Kamchatka] (Late Miocene) <HT: CGM Kol. 10255 No. 6> {re-described by SHMIDT (1984)}.

Genus *Kewia* NISIYAMA, 1935

Pliocene

- K. kehoei* WAGNER, 1974: 117; pl. 3: figs 8-10. [Bear Lake, Alaska Peninsula] (Early Pliocene) <HT: UA 2427>.
- K. tachilniensis* WAGNER, 1974: 119; pl. 2: figs 9-11. [Tachilni Fm., Walrus Peak, Cape Tachilni area, Alaska Peninsula] (Early Pliocene) <HT: USNM 181148; PT: USNM 181149>.

Miocene

- K. elegantula* SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 912 [EN: 75]; pl. 1: figs 6a-6. [Middle Etolon [Etolonsk] Suite, Kavran Series, Nepropusk Point, western Kamchatka] (Late Miocene) <HT: CGM Kol. 10255 No. 12> {mentioned in figure caption only; re-described by SHMIDT (1984)}.
- K. etolonica* SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 911-912 [EN: 76]; pl. 1: figs 4a-6. [Etolon [Etolonsk] Suite and Kakertok Suite, Kavran Series, Nepropusk Point and Kavran River, western Kamchatka] (Middle to Late Miocene) <HT: CGM Kol. 10255 No. 14> {re-described by SHMIDT (1984)}.
- K. kannoi* WAGNER, 1974: 116-117; pl. 2: figs 1-4. [Narrow Cape Fm., Narrow Cape, Kodiak Island, Alaska] (Middle Miocene) <HT: UA 2428; PT: UA 2429-33>.
- K. lituyaensis* WAGNER, 1974: 118-119; pl. 2: figs 12-15. [Yakataga Fm., Icy Point, Lituya Bay area, eastern Gulf of Alaska] (Late Miocene) <HT: USNM 181145; PT: USNM 181146-47, UA 2436, UCMP 14084>.
- K. minima* SHMIDT in SHMIDT & SINYELNIKOVA, 1971: 912 [EN: 75]; pl. 1: fig. 5a. [Upper Kakertok Suite, Kavran Series, Kavran River, Kavran-Utkholok Bay, western Kamchatka] (Middle to Late Miocene) <no type information provided> {mentioned in figure caption only}.

Oligocene

- K. marquamensis* LINDER, DURHAM & ORR, 1988: 953-954; figs 5.1-5.8. [Scott Mills Fm., Abiqua Mb., Butte Creek and Marquam Mb., Wilhoit, Marion County, Oregon, USA] (Late Oligocene) <HT: UCMP 38207; PT: UCMP 38208-A – 38208-F>.

Genus *Remondella* DURHAM, 1955

WAGNER (1974: 120) placed this genus into the family Dendrasteridae LAMBERT, 1889.

Pliocene

- R. waldroni* WAGNER, 1974: 120-121; pl. 3: figs 1-4. [Tachilni Fm., Cape Tachilni, Alaska Peninsula] (Early Pliocene) <HT: UA 2425; PT: USNM 181153-55, UCMP 14086-87>.

Miocene

- R. asiatica* SHMIDT, 1984: 149; pl. 26: figs 6a, б, г. [Etolon [Etolonsk] Suite, Kavran Series, north-eastern part of the Tochilinsky Region [Section], western Kamchatka] (Middle Miocene) <HT: VNIGRI kol. 820, no. 1>.

R. kamtschatica SHMIDT, 1984: 149-150; pl. 27: figs 13a, b. [Etolon [Etolonsk] Suite, Kavran Series, north-eastern part of the Tochilinsky Region [Section], western Kamchatka] (Middle Miocene) <HT: VNIGRI kol. 820, no. 2>.

Genus *Vaquerosella* DURHAM, 1955

Miocene

V. coreyi DURHAM, 1955: 167. (Miocene) <HT: UCMP 31712; PT: UCMP 11027> {new name for *Scutella norrisi* LOEL & COREY, 1932 not KEW, 1920}.

Family Monphorasteridae LAHILLE, 1896

Genus *Amplaster* MARTÍNEZ, 1984

Amplaster MARTÍNEZ, 1984: 506. Type-species: *A. coloniensis* MARTÍNEZ, 1984. [Uruguay] (Late Miocene).

Miocene

A. alatus ROSSI DE GARCÍA & LEVY, 1989: 90-92; fig. 1; pl. 1: figs 1-4. [Monte León Formation, Chubut Province, Patagonia, Argentina] (Late Oligocene to Early Miocene) <HT: SGNP 15.527; PT: SGNP 15.528-31> {placed into *Lunulaster* PARMA & MARTÍNEZ, 1995 by PARMA & MARTÍNEZ (1995: 65)}.

A. coloniensis MARTÍNEZ, 1984: 506-507; figs 1-4. [Camacho Fm. at Barranca de Los Loros, Uruguay] (Late Miocene) <HT: MNA-CPO 3426; PT: MNA-CPO 3425>.

A. ellipticus MOOI, MARTÍNEZ & PARMA, 2000: 269-270; fig. 4.2. [Camacho Fm. at Barranca de Los Loros and El Manzano, Uruguay] (Upper Miocene) <HT: FCDP 2205; PT: MNA-CPO 3425>.

Genus *Lunulaster* PARMA & MARTÍNEZ, 1995

Lunulaster PARMA & MARTÍNEZ, 1995: 65. Type-species: *Amplaster alatus* ROSSI DE GARCÍA & LEVY, 1989. [Monte León Formation, Chubut Province, Patagonia, Argentina] (Late Oligocene to Early Miocene).

Family Mellitidae STEFANINI, 1911

JENSEN (1982: 95) placed this family into the order Laganoida JENSEN, 1982.

Genus *Encope* L. AGASSIZ, 1840

Miocene

- E. gatunensis* TOULA, 1911: 489-491; pl. 30 (I), fig. 2. [upper part of the Gatun Formation, from the base of the second lock of the Panama canal, Gatún, Panama] (Tortonian, Late Miocene) <HT: NHMW 1933/0018/0006 (ex coll. K&K techn. Hochschule)> {species based on a very fragmentary specimen; revised age based on COLLINS et al. (1996)}
- E. michoacanensis* DURHAM, 1994: 113-114; fig. 1; pl. 1, figs 1-3; pl. 2, figs 1, 3. [near La Mira, Michoacán, Mexico] (Late Early Miocene) <IGM 2939; PT: IGM 2940-2943>.

Subgenus *Echinadesma* PHELAN, 1972

Pliocene-Recent

- Encope (Echinadesma)* PHELAN, 1972: 126-128. Type-species: *Encope micropora* L. AGASSIZ, 1841. [North and South America] (Pliocene-Recent). {other species included: *Encope emarginata* (LESKE, 1778); *Encope perspectiva* L. AGASSIZ, 1841; *Encope secoensis* COOKE, 1961; *Encope wetmorei* A.H. CLARK, 1946}.

Genus *Leodia* GRAY, 1851

Pliocene

- L. divinata* MOOI & PETERSON, 2000: 1085-1089; figs 2, 3.1-3.3, 4.1, 4.2. [Mare Fm., FCB-14 (Mencher's locality 292), Caracas, Venezuela] (Lower? Pliocene) <HT: CASG 67946>.

Genus *Mellita* L. AGASSIZ, 1841

Recent

- M. eduardobarrosoi* CASO, 1981: 142-144; figs 1-35; tabs 1-2. [Playa del Revolcadero (Acapulco, Gro.), Playa Encantada (Guerrero, Gro.), and Playa de San Benito (Tapachula, Chiapas, Chis), Pacific Coast, Mexico] (Recent) <ST: UNAM, Colección de Equinodermos del Centro de Ciencias del Mar y Limnología, no specimen nos. given>.
- M. isometra* HAROLD & TELFORD, 1990: 1002-1005; figs 11-13. [East coast of North America, from Nantucket, Massachusetts to Fort Lauderdale, Florida] (Recent)

<HT: USNM E27948; PT: LACM 84-236.1, 84-236.2; MCZ 12050, 12051; USNM E37949 – E37953; ZMUC EE220-2, EE220-3>.

Family *Astriclypeidae* STEFANINI, 1911

JENSEN (1982: 95) placed this family into the order Laganoida JENSEN, 1982.

Genus *Amphiope* L. AGASSIZ, 1840

Miocene

- A. caupianensis* CHAVANON, 1974: Vol.1: 142-143, Vol.2: 98; figs 64-70; pl. 10: figs 5a-b. [Caupian (Saint Médard en Jalles), Aquitaine Basin, Western France] (Early Miocene) <ST: Castex colln, no repository nos given for types; figured specimen: 20.C.14>.
- A. miocenica* ALI, 1998: 545-546; fig. 3D. [Marmarica Fm., Gebel west El Migahhiz, Siwa Oasis, Egypt] (Langhian-Serravalian) <HT: GM-GDEMU, no specimen no.>.

Genus *Astriclypeus* VERRILL, 1867

Pleistocene

- A. miaoliensis* WANG, 1986: 170-175; figs 10, 11; pl. 7, figs 1-5; pl. 8, figs 1-2; pl. 9, figs 7. [Toukoshan Fm., Wuhu, Miaoli-hsien, northern Taiwan] (Late Pleistocene) <HT: NTUG-(E)-0068; PT: NTUG-(E)-0189 – 0191>.

Pliocene

- A. pitouensis* WANG, 1986: 166-170; figs 9a-b; pl. 6, figs 2-5; pl. 9, fig. 3. [Erhchiu Fm., Pitou, Taipei-hsien, northern Taiwan] (Early Pliocene) <HT: CGS-E-83026; PT: CGS-E-83027, 83028>.

Miocene

- A. elegans* WANG, 1986: 158-166; figs 7, 8a-e; pl. 3, figs 1-4; pl. 4, figs 1-6; pl. 5, figs 1-6; pl. 6, fig. 1; pl. 8, fig. 3; pl. 9, fig. 2. [Nankang Sandstone; Shuinantung, Kengtsuliao and Shen-AO; Taipei-hsien, northern Taiwan] (Middle Miocene) <HT: NTUG-E-80017; PT: NTUG-E-80018, 80020 – 80022, 80148, 81053, 81063, 81064, 83029, NTUG-(E)-0167, 0165, CGS-E-83034 – 83037>.
- A. yeliuensis* WANG, 1986: 152-158; figs 5, 6a-e; pl. 1, figs 1-3; pl. 2, figs 1-5. [Taliao Fm., Yeliu Park, Taipei-hsien, northern Taiwan] (Early Miocene) <HT: NTUG-E-81046; PT: NTUG-E-81043 – 81045, 81052>.

Oligocene

- A. waiwulunensis* WANG, 1983: 115-118; pl. 1, figs 1-2; pl. 2, fig. 3. [Wuchihshan Fm., Waiwulun-Ao, Chilung City, northern Taiwan] (Late Oligocene) <HT: NTUG-E-80031; PT: NTUG-E-80033>.

Genus *Echinodiscus* LESKE, 1778

Cenozoic

- E. colchesterensis* SMUTS, 1988: 94-97; figs 2a-d, 3b-c. [Alexandria Fm., Colchester Cliff, Colchester, South Africa] (Late Tertiary) <HT: GSP A8817; PT: GSP A8819-8826, A9000-9001>.

Miocene

- E. yeliuensis* WANG, 1982a: 151-156; pl. 1, figs 1-2; pl. 2, figs 1-4; pl. 3, figs 1-6; pl. 4, fig. 1. [Taliao Fm., Yeliu, Keelung City, Northern Taiwan] (Aquitanian) <HT: NTUG-E-81029; PT: NTUG-E-81030, 81032, 81033, 81049>.

Oligocene

- E. hsianglanensis* WANG, 1986: 176-180; fig. 12; pl. 9, figs 1, 4-6. [Fangchiao Fm., Hsi-aohsianglan, Taipei-hsien, northern Taiwan] (Late Pleistocene) <HT: CGS-E-83031; PT: CGS-E-83032, 83033, 83039>.

Paleocene-Eocene

- E. tiliensis* WANG, 1984a: 107-110; pl. 1, figs 1a-b. [Tachien Sandstone, Tili Village, Hsini-hsiang, Nantou-hsien, Central Taiwan] (Late Paleocene-Early Eocene) <HT: NTUG-E-0236>.

Genus *Kieria* MIHÁLY, 1985

- Kieria* MIHÁLY, 1985: 243, 261. Type-species: *Kieria semseyana* MIHÁLY, 1985 [Hungary] (Miocene) {according to KROH (2005: 97, 100) the genus *Kieria* is based upon juvenile specimens of *Amphiope*}.

Miocene

- K. semseyana* MIHÁLY, 1985: 243, 261; pl. 4, figs 2-6. [Leitha Limestone, Gyakorló út [Street], 10th district, Budapest, Hungary] (Late Badenian (= Early Serravallian), Middle Miocene) <HT: MAFI Ech 355; PT: MAFI Ech 356, 357> {according to KROH (2005: 97, 100) the genus *Kieria* is based upon juvenile specimens of *Amphiope* and this species possibly is a junior synonym of *Amphiope bioculata* (DES MOULINS, 1837)}.

Family Abertellidae DURHAM, 1955

Genus *Abertella* DURHAM, 1955

Miocene

- A. complanata* BRITO, 1981a: 3-5; fig. 1-2. [Pirabas Fm., Capanema, Jazida B-9, Estado do Pará, Brasil] (Lower Miocene) <HT: MNRJ 5460-I>.
- A. gualichensis* MARTÍNEZ, REICHLER & MOOI, 2005: 1230-1232; figs 2.1-2.4; figs 3.1, 3.2. [Gran Bajo del Gualicho Formation, Salina del Gualicho, Río Negro Province, Argentina] (Late Early – Early Middle Miocene) <HT: MACN-Pi 4714; PT: MACN-Pi 4705, 4706, 4709>.

Family Scaphechinidae BEADLE, 1991

Scaphechinidae BEADLE, 1991: 327. Type-genus: *Scaphechinus* A. AGASSIZ, 1863. [Japan, Formosa] (Miocene-Recent) {attributed to MOOI (1987), which is an unpublished Ph.D. thesis}.

Genus *Scaphechinus* A. AGASSIZ, 1863

Recent

- S. brykovi* BUDIN, 1983: 55-56; pl. 3: fig. 3. [Peter the Great Bay, Sea of Japan] (Recent) <HT: IMBFE 3108/5>.

Family Uncertain

Genus *Samlandaster* LAMBERT & THIÉRY, 1914

Bartonian

- S. dineuri* ROMAN, 1989: 300-303; pl. 1, figs 4, 12-13. [Sables roux, à la base de la formation des Sables d'Auvers; Saint-Vaast-lès-Mello, Oise, lie-dit Barisseuse, Paris Basin, France] (Auversien, Early Bartonian) <HT: IPM R09016; PT: IPM R09017>.

Genus *Sinaechinocyamus* LIAO, 1979

Sinaechinocyamus LIAO, 1979: 67, 70-71. Type-species: *Sinaechinocyamus planus* LIAO, 1979. [Yellow Sea] (Recent).

Recent

S. planus LIAO, 1979: 67-69, 71-72; figs 1-3, 4.1-4.4; pl. 1: figs 1-4. [Northern Huang Hai (37°50' N, 124°E, depth 72 m, muddy sand bottom), Yellow Sea] (Recent) <HT: IOAS E00931; PT: IOAS E00932> {according to MOOI (1990b: 143) *Sinaechinocyamus* is a member of the suborder Scutellina related to *Scaphechinus*}.

Superfamily Taiwanasteritida WANG, 1984b

Taiwanasteritida WANG, 1984b: 134. {families included: Fibulariidae GRAY, 1855; Taiwanasteridae WANG, 1984b; MOOI (1990b: 142-143) rejected this superfamily}.

Family Taiwanasteridae WANG, 1984b

Taiwanasteridae WANG, 1984b: 134. Type-genus: *Taiwanaster* WANG, 1984b. (Early Pliocene to Recent) {MOOI (1990b: 142-143) rejected this family}.

Genus *Taiwanaster* WANG, 1984b

Taiwanaster WANG, 1984b: 134-135. Type-species: *Taiwanaster mai* WANG, 1984b. [Taiwan] (Recent) {according to MOOI (1990b: 139) this genus is a junior synonym of *Sinaechinocyamus* LIAO, 1979}.

Recent

T. mai WANG, 1984b: 135-143; figs 4c, 5c, 6a-f, 9a-b; pl. 1, figs 1-3; pl. 2, figs 1-6; pl. 3, figs 1, 3-4; pl. 4, figs 1-4; pl. 5, figs 1-4. [Taiwan Strait near Yunlin-hsien, Taiwan] (Recent) <HT: NTUG-E-81070; PT: NTUG-E-81071, 81072, 81075-81077, 81082-81087>.

Pliocene

T. gutingkengensis WANG, 1984b: 147-151; figs 5a, 8a-c; pl. 8, figs 1-4. [Lower Gutingkeng Fm., Gutingkeng, Kaohsiung-hsien, southern Taiwan] (Latest Pliocene) <HT: NTUG-E-81088; PT: NTUG-E-81089-81093>.

T. pitouensis WANG, 1984b: 143-147; figs 5b, 7a-d; pl. 3, fig. 2; pl. 6, figs 1-6; pl. 7, figs 1-2. [Erhchiu Fm., Pitou-chiao, northern Taiwan] (Early Pliocene) <HT: CGS-E-82001; PT: CGS-E-82002-82004, 82006, 82007, 82026, 82027>.

Superorder Atelostomata ZITTEL, 1879

Order Menopygoida VADET, 1995

Superorder Microstomata SMITH, 1984

Microstomata SMITH, 1984: 171, 173 {includes Cassiduloida and Clypeasteroida}.

Order Cassiduloida CLAUS, 1880

Genus *Pronucleolites* BARRAS, 2006

Pronucleolites BARRAS, 2006: 60-61. Type-species: *Clypeus orbicularis* PHILLIPS, 1829. [Europe] (Jurassic) {regarded as stem-group cassiduloid by BARRAS (2006)}.

Early Jurassic

P. ? crickleyensis BARRAS, 2006: 67-69; text-figs 37a-b, 38; pl. 5: figs 4-6. [Inferio Oolite, Crickley, Gloucestershire, England] (*murchisoniae* zone, Aalenian) <HT: BGS GSM 119711>.

Family Galeropygidae LAMBERT, 1911

Genus *Eogaleropygus* JESIONEK-SZYMAŃSKA, 1978

Eogaleropygus JESIONEK-SZYMAŃSKA, 1978: 191. Type-species: *Pygaster microstoma* LAMBERT, 1933. [Morocco] (Middle Toarcian).

Genus *Galeropygus* COTTEAU, 1856

Jurassic

G. parviphyllus BARRAS, 2006: 58-59; text-figs 32a-b; pl. 5: figs 1-3. [unknown, unknown (? England)] (unknown, possibly Jurassic) <HT: NHM E 42916>.

Genus *Hyboclypeus* L. AGASSIZ, 1839

Middle Jurassic

H.? *pavyi* ROBERT, 1994: 67-68; pl. 2: figs 8-10. [Calcaires siliceux à chailles; Brétignelles, près Druyes, Dept. Yonne, France] (*Antecedens* subzone, *Plicatilis* zone, Middle Oxfordian) <HT: Philippe Robert coll'n No. 604> {based on an internal mould}.

Genus *Pygomalus* POMEL, 1883

Late Jurassic

P. azemati DEVRIÈS, 1972: 45-46; pl. 2: figs 1-6. [Puig Santa Magdalena, north-east of Novelda, Province Alicante, Spain] (Late Kimmeridgian/Tithonian) <HT (by monotypy): coll. J.Azema, Paris [no specimen no. provided]>.

Family Clypeidae LAMBERT, 1898

Genus *Angusticlypeus* VADET, 1997

Angusticlypeus VADET, 1997: 48. Type-species: *Clypeus angustiporus* AGASSIZ in AGASSIZ & DESOR, 1847. [Europe] (Bajocian-Callovian) {other species included: *C. altus* MAC COY, 1848; *C. babeaui* COTTEAU, 1870; *C. davoustianus* COTTEAU in DAVOUST, 1856; *C. mulleri* WRIGHT, 1859; *C. rathieri* COTTEAU, 1849}.

Genus *Bothryopneustes* FOURTAU, 1924

Upper Jurassic

B. inflata KIER, 1972a: 64-66; fig. 31; pl. 34, figs 1-7; pl. 41, fig. 6. [Upper Dhurma Fm., locality KK9-95-112, Saudi Arabia] (Callovian) <HT: USNM 170426; PT: USNM 170427-170429>.

B. kauffmani KIER, 1972a: 63-64; fig. 30; pl. 35, figs 1-7; pl. 36, figs 1-5. [Upper Dhurma Fm., locality KK9-112-113, Saudi Arabia] (Callovian) <HT: USNM 170430; PT: USNM 170431-170432>.

Middle Jurassic

- B. dhrumaensis* KIER, 1972a: 61-62; fig. 28; pl. 33, figs 1-6. [Middle Dhruma Fm., localities KK8-30-44, KK9-35-98, L-921, L-922, S-1045 and S-1154, Saudi Arabia] (Bathonian) <HT: USNM 170421; PT: USNM 170422-170423>.
- B. arabica* KIER, 1972a: 62-63; fig. 29; pl. 32, fig. 5; pl. 33, figs 7-10. [Middle Dhruma Fm., localities KK8-35-46 and L-920, Saudi Arabia] (Bathonian) <HT: USNM 170420; PT: USNM 170424-170425>.

Genus *Clypeus* LESKE, 1778

Middle Jurassic

- C. ederlei* ROBERT, 1994: 68-69; pl. 14: figs 1-3. [Les Rochers du Parc, près de Mailly-la-Villec, Yonne, France] (*Bifurcatus* subzone, *Transversarium* zone, Middle to Late Oxfordian) <HT: Robert Ederlé coll'n, Perrigny (Yonne) No. 514>.

Genus *Colliclypeus* SMITH in SMITH & BENGGTSON, 1991

- Colliclypeus* SMITH in SMITH & BENGGTSON, 1991: 37-38. Type-species: *Conoclypus nettoanus* WHITE, 1887. [Brasil] (Early Cretaceous).

Genus *Kieripygurus* VADET, 1997

- Kieripygurus* VADET, 1997: 59. Type-species: *Pygurus jurensis* MARCOU in AGASSIZ & DESOR, 1847. [Europe] (Callovian-Valanginian) {other species included: *P. geryvilensis* PERON & GAUTHIER in COTTEAU, PERON & GAUTHIER, 1888; *P. rostratus* AGASSIZ, 1839; *P. tenuis* DESOR in AGASSIZ & DESOR, 1847}.

Genus *Mercieripygurus* VADET, 1997

- Mercieripygurus* VADET, 1997: 58. Type-species: *Mepygurus hausmanni* (KOCH & DUNKER, 1837). [Europe] (Oxfordian).

Genus *Pygurus* L. AGASSIZ, 1839

Referred to a new family, Pyguridae, by VADET (2007).

Albian

P. (P.) mendelssohni GREYLING & COOPER, 1993: 13-17; figs 2-7. [Makhatini Flats, SE Jozini Dam, Zululand, South Africa] (Upper Albian) <HT: DNSM-PCZ4649; DNSM-PZ4562-66, 4639, 4660>.

Barremian

P. (P.) posteroexpansus TANAKA, 1984b: 190-192; text-fig. 2; pl. 1, figs 3a-c. [Haidate Grp., SE Nakaosaka, Mie-machi, Ono-gun, Oita Prefecture, Japan] (Upper? Barremian) <HT: GSJ F6076A, B>.

Lower Cretaceous

P. (P.) yamamaensis KIER, 1972a: 72; pl. 46, figs 9-11; pl. 47, figs 1-5. [Yamama Fm., localities KK1-41, L-901, L-902, L-905, L-910, L-908. KK3-45 and S-1262, Saudi Arabia] (Neocomian) <HT: USNM 170456; PT: USNM 170457-170459>.

Late Jurassic

P. (P.) arabicus KIER, 1972a: 66-68; fig. 33; pl. 38, fig. 4-7. [Upper Dhurma Fm., Atash Mb., locality L-926, Saudi Arabia] (Late Jurassic) <HT: USNM 170436>.

P. (P.) andinus LARRAIN & BIRÓ-BAGÓCZKY, 1985: 1410-1413; figs 3, 4.1-4.4. [Baños del Flaco Fm., Río Maitenes, Cajón del Fierro, Chile] (Upper Tithonian) <HT: CPUC RM./85-5; PT: CPUC RM./85-1 – RM./85-3>.

Middle Jurassic

P. beineuensis PORETSKAYA, 1972: 291-292; figs 62-63; pl. 60: figs 12a-b. [Kuvali-Tjube, Tuarkyrsibiyru, western Turkmenistan] (Late Callovian, Middle Jurassic) <HT: LGU 1/296>.

Family Pyguridae VADET, 2007

Pyguridae VADET, 2007: 180-181. Type-genus: *Pygurus* AGASSIZ, 1840. (Jurassic).

Family Nucleolitidae L. AGASSIZ & DESOR, 1847

Genus *Baudhuinipygus* VADET, 1997

Baudhuinipygus VADET, 1997: 76. Type-species: *Echinobrissus haimeii* WRIGHT, 1859. [Europe] (Portlandian) {other species included: *Nucleolites brodiei* (WRIGHT, 1859)}.

Genus *Catopygus* L. Agassiz, 1836

Cretaceous

- C. altus* SZÖRÉNYI, 1955: 69-70, 207-208; pl. 9, figs 5, 7. [calcaire compact à Orbitolines, calcaire gris lamelleux, Alsópere, Olaszfalu-Eperkeshegy, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/293 (from Alsópere)>.
- C. dilatatus* TZANKOV, 1984: 77; pl. 31, figs 7, 7a-b. [Dar-Boaz, Provadia, Bulgaria] (Mastrichtian) <HT: USC CR₂ 1280>.
- C. loevisiformis* TZANKOV, 1984: 76-77; pl. 31, figs 6, 6a-b. [Choumen, Bulgaria] (Mastrichtian) <HT: USC CR₂ 1279>.
- C. neocylindricus* SZÖRÉNYI, 1955: 70-72, 208-210; fig. 24; pl. 9, figs 13-28. [calcaire compact à Orbitolines, calcaire gris lamelleux, Bakonyánána (carrière de la vallée Gaja, côte E du mont Judenberg), Pénezskut-Körisgyörpuszta, and Olaszfalu-Villóhegy, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/265 (from Bakonyánána)>.
- C. subcircularis sulcatus* SAVCHINSKAYA, 1974: 316-317; pl. 98, figs 17-22. [Northern Don Basin, Russian SFSR] (*Belemnitella longei*-zone, Late Campanian) <HT: not given>.

Genus *Hungaresia* SZÖRÉNYI, 1955

Hungaresia SZÖRÉNYI, 1955: 76, 215. Type-species: *Hungaresia hungarica* SZÖRÉNYI, 1955 [Eastern Europe] (Mid-Cretaceous). {other species included: *Hungaresia minor* SZÖRÉNYI, 1955}.

Mid-Cretaceous

- H. minor* SZÖRÉNYI, 1955: 78, 216-217; pl. 11, figs 1-5. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/344>.
- H. hungarica* SZÖRÉNYI, 1955: 77-78, 215-216; fig. 25; pl. 10, figs 14-22. [bane de calcaire marneux jaune et bane de calcaire à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, and Sümeg (grande carrière de Gerine), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/338 (from Sümeg)>.

Genus *Nucleolites* LAMARCK, 1801

(Genus *Echinobrissus* GRAY, 1825; objective junior synonym
of *Nucleolites* LAMARCK, 1801)

Cretaceous

- N. elongatus* CHIPLONKAR & BADVE, 1972: 142-143; pl. 12: figs 17, 22. [Deola-Chirakhan Marl, Chirakhan (22°22'30" N, 75°07' E), India] (Cenomanian) <HT: MACSG CH 401>.
- E. faasi* SHMIDT & SIMAKOV, 1953: 43-44; figs 10a-b; pl. 2: figs 16-18. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Early Turonian) <HT: VNIGRI 38/336>.
- E. gultschensis* SHMIDT & SIMAKOV, 1953: 42-43; figs 9a-r; pl. 2: figs 19-26. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Late Turonian) <HT: VNIGRI 17/336>.
- E. (Nucleolites) pannonicus* SZÖRÉNYI, 1955: 73-74, 211-212; pl. 10, figs 1-5. [bane de marne jaune et grise du groupe de calcaires à Hippurites, Sümeg-Kövesdomb, and Sümeg (grande carrière de Gerine), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/302 (from Sümeg)>.

Late Jurassic

- N. strictiporus* VADET, 2007: 204-205; illustr. on p. 205. [Hesdin l'Abbé, Boulonnais, France] (Oxfordian, Late Jurassic) <HT: Vadet colln V5140; PT: MHNB 662, 1007-1014, Vadet colln V2999-V3006, 3878, 3879, 5377, 5378, 7281, 7282, 7301-7307>.
- E. volgensis* GERASIMOV, 1955: 23-24; pl. 4: figs 6, 7a, б, 8a, б, 9, 10a, б, 11, 12. [Right-hand side of River Volga, near the villages Glebovo, Mostovo and Koprino, Yaroslavskog Oblast, USSR] (Early Volgian (= Middle Tithonian), zone of *Virgatites virgatus*) <HT: GMM 855-4>.

Genus *Oolopygus* D'ORBIGNY, 1857

Late Cretaceous

- O. globulorostratus* SAVCHINSKAYA, 1974: 317-318; pl. 100, figs 1-5. [Northern Don Basin, Russian SFSR] (*Belemnitella longei*-zone, Late Campanian) <HT: not given>.

Genus *Plagiochasma* POMEL, 1883

Maastrichtian

P. lammersmaxi JAGT & VAN DER HAM in JAGT, 2000: 252-254; pl. 17, figs 1-3, 10. [lowermost Meerssen Mb., Maastricht Fm., ENCI-Maastricht Quarry, Maastricht, The Netherlands] <HT: NHMM TL 1985/1>.

Early Cretaceous

P. nadorensis VADET, MARIGNAC, NICOLLEAU & REBOUL, 2007: 32-33; pl. 16: figs 2-3; pl. 17: figs 1-2. [Sidi Bouzid, N of Safi, Morocco] (Late Berriasian to Early Valanginian, Early Cretaceous) <HT: Reboul coll. 308>.

Genus *Primipygus* VADET, 2007

Primipygus VADET, 2007: 192. Type-species: *Nucleolites richardsoni* PARIS in RICHARDSON & PARIS, 1908. [France] (Bajocian, Jurassic). {objective junior synonym of *Notopygus* POMEL, 1883; considered to be a subjective junior synonym of *Pseudosorella* ETALLON, 1859, p. 415 by SMITH (2008 – The Echinoid Directory, accessed 14/07/2008)}.

Genus *Pygaulus* L. AGASSIZ, 1847

Turonian

P. aksuensis SHMIDT & SIMAKOV, 1953: 38; figs 5a-6; pl. 2: figs 4-6. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Early Turonian) <HT: VNIGRI 14/336> {referred to *Pygorrhynchus* by EGOROV (1972: 59)}.

P. faasi SHMIDT & SIMAKOV, 1953: 36-37; figs 4a-b; pl. 1: figs 18-20. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Early Turonian) <HT: VNIGRI 10/336> {referred to *Pygorrhynchus* by EGOROV (1972: 59)}.

Cenomanian

P. richardi NASTAJ, 1985: 19-22; figs 1a-b, 2a-b, 3a-; pl. 1, figs 3a-c, 4a-b. [Brzozówka, near Korzkiew, near Cracow, Poland] (Cenomanian) <HT: Department of the Institute of Geology and Mineral Deposits, University of Mining and Metallurgy, Cracow, no specimen no. given>.

Late Cretaceous

P. cyclotus YANG SHENGQIU, 1991: 116-117; pl. 10; figs 1-10. [Upper Kukebai Fm., 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Late Cretaceous) <HT: NIGP, no holotype defined>.

Genus *Pygopistes* POMEL, 1883

Paleocene

P. parrasae PARMA & CASADIO, 2005: 1076-1078; figs 5.2-5.4, 7, 8. [Roca Formation, Casa de Piedra and Cerros Bayos, La Pampa Province, Argentina; Bajo Hondo, Río Negro Province, Argentina] (NP2-NP3 alcareous nannofossil zone, Early Danian, Paleocene) <HT: PMOZ 4087 (from Casa de Piedra); PT: PMOZ 4088, GHUNLPam 22154 (both from Casa de Piedra), GHUNLPam 22009-22014 (from Cerros Bayos), CPBA 6334 (from Bajo Hondo)> {redescribed by DEL RIO et al. (2007: 260-261)}.

Genus *Pygorhynchus* L. AGASSIZ, 1839

(Genus *Botriopygus* D'ORBIGNY, 1856;
objective junior synonym of *Pygrhynchus* L. AGASSIZ, 1839)

Cretaceous

[*P. acutus* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

B. baconicus SZÖRÉNYI, 1955: 64-65, 202-203; pl. 7, figs 1-4. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/224>.

P. clypeiformis DEVIRIES, 1973: 67-68; pl. 2, figs 7-9. [Kef el Mekene, map sheet Ain Regada, Algeria] (Barremian, Early Cretaceous) <no types defined, material in the colln of A. Devries>.

P. debensis DEVIRIES, 1973: 68-69; pl. 2: figs 10-12. [Kef ed Deb, map sheet Ain Regada, Algeria] (Barremian, Early Cretaceous) <no types defined, material in the colln of A. Devries>.

B. latipetalus SZÖRÉNYI, 1955: 63, 200-201; pl. 7, figs 11-12, 14-15. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/247>.

B. neoataxensis SZÖRÉNYI, 1955: 63-64, 201-202; pl. 7, figs 6, 8, 10, 13. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/244>.

B. ovalis SZÖRÉNYI, 1955: 64, 202; pl. 7, figs 5, 7, 9. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/235>.

B. pappi SZÖRÉNYI, 1955 [BARNABÁS, 1937]: 61, 198; pl. 6, figs 13-15. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <NT: GIH Eb/183> {SZÖRÉNYI (1955) cites BARNABÁS (1937) as author of this

- species, but this paper seems to be an unpublished Ph.D. thesis, unless earlier published descriptions are found, SZÖRENYI (1955) is to be considered as author of this species}.
- B. pappi elongatus* SZÖRENYI, 1955: 62, 199-200; pl. 6, figs 1, 3, 6, 10. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/205>.
- B. pappi kutassyi* SZÖRENYI, 1955 [BARNABÁS, 1937]: 61-62, 199; pl. 6, figs 5, 7-8. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <NT: GIH Eb/187> {SZÖRENYI (1955) cites BARNABÁS (1937) as author of this species, but this paper seems to be an unpublished Ph.D. thesis, unless earlier published descriptions are found, SZÖRENYI (1955) is to be considered as author of this subspecies}.
- B. petalodes planus* SZÖRENYI, 1955: 62-63, 200; pl. 6, figs 2, 4, 9, 11-12. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/250>.
- B. subquadratus* SZÖRENYI, 1955: 65-66, 203-204; pl. 8, figs 1-4, 6. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/219>.
- B. suemegensis* [*sümegensis*] SZÖRENYI, 1955: 66-67, 204-205; pl. 8, figs 8, 11, 13, 15. [groupe de calcaires à *Hippurites*, Sümeg (carrière de Gerine), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/218>.
- B. variabilis* SZÖRENYI, 1955: 67, 205-206; pl. 8, figs 5, 7, 9-10, 12, 14, 16. [bane à *Botriopygus* du groupe de calcaires à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/212>.

Family Echinolampadidae GRAY, 1851

Genus *Arnaudaster* LAMBERT, 1918

Maastrichtian

- A. cylindriformis* SMITH, 1995: 219-222; fig. 73-75; pl. 25: figs 11-12; pl. 29: figs 1-9. [. [Simisima Fm., Jebel Buhays and Jebel Rawdah, Oman/United Arab Emirates Border Region] <HT: BMNH EE4334; PT: BMNH EE4324, EE4331-33, EE3378-81>.

Genus *Echinolampas* GRAY, 1825

Recent

E. alexandri forma *arctambulacrum* DOLLFUS & ROMAN, 1981: 106; pl. 35, figs 3-5. [Station 17, Red Sea] (Quarternary – Recent) <HT: not given>.

Miocene

E. antunesi GONÇALVES, 1971: 308-310; figs 1-2. [Farol des Lagostas, niveau II of Antunes, 1964, région côtière au Nord de Luanda, Angola, Africa] (Middle to Late Miocene) <HT: MMGL no repository no. given> {based on a single, damaged specimen}.

E. (Isolampas) bononii VENZO, 1934: 114-116; pl. 1, figs 9, 9a-b. [Monte Brione, Italy] (“Aquitanian Veneto”, Early Miocene) <HT: ? Instituto Geologico della R. Univ. Pisa> {based on a single specimen}.

E. duncani MCNAMARA, 1987a: 109-110; figs 1a-c. [Bairnsdale Limestone Mb., Gippsland Limestone, Bairnsdale, Victoria, Southern Australia] (Bairnsdalian, Middle Miocene) <HT: BMNH E1107> {name preoccupied; MCNAMARA (1989: 257), proposed *E. laubei* as replacement name}.

E. gregoryi gregoryi MCNAMARA & PHILIP, 1980c: 8; pl. 4, figs 4-6. [Muddy Creek Fm., Clifton Bank, South Australia] (Balcombian, Middle Miocene) <HT: NMV P55477>.

E. gregoryi corrugata MCNAMARA & PHILIP, 1980c: 8; pl. 4, figs 7-9. [Bairnsdale Limestone, South Australia] (Middle Miocene) <HT: NMV P18379>.

E. laubei MCNAMARA, 1989: 257. [Bairnsdale Limestone Mb., Gippsland Limestone, Bairnsdale, Victoria, Southern Australia] (Bairnsdalian, Middle Miocene) <HT: BMNH E1107> {nomen novum for *E. duncani* MCNAMARA, 1987a non COTTEAU, 1891}.

E. (Hypsoclypus) liae TAVANI, 1939: 42; pl. 3 (5), fig. 2. [Umm er Rzem, Cyrenaica, Libya] (Langhian) <no specimen no.>.

E. schultzi KROH, 2005: 119-122; fig. 43.5, 51; pl. 54, figs 2-3; pl. 55, figs 1-3; pl. 56, figs 1-2. [Zogelsdorf Formation, train station Eggenburg, Lower Austria, Austria] (Late Eggenburgian (= Early Late Burdigalian), Early Miocene) <HT: formerly at the Krauhletzt Museum (Eggenburg, Austria), now lost; NT: NMHW 2003z0040/0001> {nomen novum pro *E. laurillardi acuminata* SCHAFFER, 1912, preoccupied by *E. acuminata* ABICH, 1882 (= *E. laurillardi* sensu LAUBE, 1871, non AGASSIZ in AGASSIZ & DESOR, 1847)}.

E. tumulus MCNAMARA & KENDRICK, 1994: 42-44; figs 16A-D. [Poivre Fm.; Eagle’s Nest, Northern Flacourt Bay and Cape Maloet; Barrow Island, Western Australia] (Middle Miocene) <HT: WAM 82.269; PT: WAM 82.207-20, 82222-251, 82.274, 82.259-269, 82.282-288>.

Oligocene

- [*E. chiplonkari* TANDON]. [Kutch, India] (*Lepidocyclina (Eulepidina)* Zone, Oligocene) <types unknown> {nomen nudum; mentioned in SRIVASTAVA (1988: 152); not established in TANDON (1973), which is an abstract only}.
- E. cookei* SRIVASTAVA & SINGH, 1999: 30-31; pl. 5, figs 5-8. [Near Guvar, Kachchh, India] (*Lepidocyclina (Eulepidina)* Zone, Rupelian) <HT: DGUL KTE 241; PT: DGUL KTE 307, KTE 318, KTE 319, KTE 357, 358>.
- E. guvarensis* SRIVASTAVA & SINGH, 1999: 32; pl. 6, figs 7-8; pl. 7, figs 1-3. [Guvar, Kachchh, India] (*Lepidocyclina (Eulepidina)* Zone, Rupelian) <HT: DGUL KTE 328; PT: DGUL KTE 321, KTE 325, KTE 335> {mentioned previously as nomen nudum in SRIVASTAVA (1988: 152)}.
- [*E. kieri* SRIVASTAVA, 1988]: 152. [India] (*Nummulites subclypeus* Zone, Oligocene) <types unknown> {nomen nudum; also mentioned in SRIVASTAVA, MISHRA & SRIVASTAVA (1992: Tab. 1), first mentioned in the unpublished Ph.D. thesis of Dinesh K. SRIVASTAVA (pers. comm. 16.X.2007)}.
- E. posterocrassa curtata* McNAMARA & PHILLIP, 1980: 3; pl. 1, figs 7-9. [Port Vincent Limestone, Adelaide Cement Holdings Quarry, N Wool Bay, Yorke Peninsula, South Australia] (*Guembelitra stavensis* Zone, Oligocene) <HT: NMV P55451; PT: NMV P55452-53>.
- E. tandoni* SRIVASTAVA & SINGH, 1999: 29-30; pl. 4, figs 5-7; pl. 5, figs 1-4. [Jhadwa, Kachchh, India] (*Nummulites fichteli* Zone to *Nummulites subclypeus* Zone, Lattdorfian-Rupelian) <HT: DGUL KTE 279; PT: DGUL KTE 233, KTE 341, KTE 342, KTE 347, KTE 354>.

Eocene

- E. bastai* ELATTAAR & STROUGO, 2001: 70-72; figs 6a-6g. [mouth of Wadi Gharandal, west-central Sinai, Egypt] (Late Eocene) <HT: ASUGM 278E (figs 6d, 6e); PT: ASUGM 278E (figs 6f, 6g)>.
- [*E. cooki* SRIVASTAVA, 1988]: 151. [India] (*Nummulites maculatus* Zone, Middle Eocene) <types unknown> {nomen nudum; mentioned also in SRIVASTAVA, MISHRA & SRIVASTAVA (1992: Tab. 1), first mentioned in the unpublished Ph.D. thesis of Dinesh K. SRIVASTAVA (pers. comm. 16.X.2007); later published as *E. cookei* in SRIVASTAVA & SINGH (1999)}.
- E. daralagezensis* PORETSKAYA, in AKOPJANA, 1974a: 362-363; pl. 187, figs 3a-6. [Rind, Armenia] (Late Eocene) <HT: LGU 306/2>.
- E. garoensis* SRIVASTAVA, SINGH, TIWARI & JAUHRI, 2008: 514; pl. 1: figs 1-9. [Siju Limestone Dhapsagiri village, South Garo Hills, Meghalaya, India] (Late Lutetian to Early Bartonian) <HT: LUGD I/2020; PT: LUGD I/2020_a>.

- [*E. granti* SRIVASTAVA, 1988]: 151. [India] (*Nummulites maculatus* Zone, Middle Eocene) <types unknown> {nomen nudum; mentioned also in SRIVASTAVA, MISHRA & SRIVASTAVA (1992: Tab. 1), first mentioned in the unpublished Ph.D. thesis of Dinesh K. SRIVASTAVA (pers. comm. 16.X.2007)}.
- [*E. jhadwensis* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <types unknown> {nomen nudum; mentioned also in SRIVASTAVA, MISHRA & SRIVASTAVA (1992: Tab. 1), first mentioned in the unpublished Ph.D. thesis of Dinesh K. SRIVASTAVA (pers. comm. 16.X.2007)}.
- E. jigniensis* SRIVASTAVA, MISHRA & SRIVASTAVA, 1992: 99-101; pl. 1, figs 1-4; pl. 2, figs 1-5. [Jigni village, Rajauri District, Jammu & Kashmir, India] (Middle Eocene) <HT: NRF-2/353; PT: NRF.2/354, 2/355, 2/356>.
- E. khariensis* SRIVASTAVA & SINGH, 1999: 26-27; pl. 2, figs 1-5. [Khari, Kachchh, India] (Middle Eocene) <HT: DGUL KTE 283; PT: DGUL KTE 284, KTE 286, KTE 287, KTE 296, KTE 297, KTE 298>.
- E. lakiensis* SAWAR, 1988: 76-80; figs 1a-c, 2a-c, 3a-c. [Laki Hills, Khithar Range, District Dadu, Sindh, Pakistan] (Late Cretaceous & Eocene) <HT: PUPCE No. 64> {based on 3 poorly preserved specimens}.
- E. lipiformis* SRIVASTAVA & SINGH, 1999: 27-28; pl. 2, figs 6-8. [Near Pipur, Kachchh, India] (*Asterocyclina alticostata* Zone, Middle Eocene) <HT: DGUL KTE 677; PT: DGUL KTE 388, KTE 678, KTE 679>.
- [*E. merhi* SRIVASTAVA, 1988]: 151. [India] (*Nummulites maculatus* Zone, Middle Eocene) <types unknown> {nomen nudum; mentioned also in SRIVASTAVA, MISHRA & SRIVASTAVA (1992: Tab. 1), first mentioned in the unpublished Ph.D. thesis of Dinesh K. SRIVASTAVA (pers. comm. 16.X.2007)}.
- E. pipurensis* SRIVASTAVA & SINGH, 1999: 28-29; pl. 3, figs 1-3. [Near Pipur, Kachchh, India] (*Nummulites maculatus* Zone, Middle Eocene) <HT: DGUL KTE 274; PT: DGUL KTE 268, KTE 275>.
- E. qattamiaensis* ALI, 1983: 57-58; pl. 1, figs 1-5. [Cairo-Building stone Fm., Gebel Qattamia, Egypt] (Upper Lutetian) <HT: USNM 341264>.
- E. rotundus* MAMEDOV & MELIKOV, 1976: 164-165; fig. 6; pl. 1, figs 4a, 4b, 4b. [In-Arieh (Tahanabat), Republic Mali] (Middle Lutetian, Eocene) <HT: AzINEFTEK-HIM 14/69>
- E. tachanabatensis* MAMEDOV & MELIKOV, 1976: 161-164; figs 4-5; pl. 1, figs 3a, 3b, 3b. [In-Arieh (Tahanabat), Republic Mali] (Middle Lutetian, Eocene) <HT: AzINEFTEK-HIM 12/69>
- E. tanypetalis* HARPER & SHAAK, 1974: 1166-1168; text-figs 2A-B; pl. 1, figs 1-4. [Ocala Limestone, Suwannee River, Lafayette County, Florida, USA] (Jacksonian, Late Eocene) <HT: BYU 1466; PT: UFS 3378, 3900>.

E. yadongensis MU & WU, 1976: 364; pl. 3: figs 13-15; pl. 4: figs 4-5. [Zongpu Grp., Duila, Yadong County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Paleocene-Eocene) <ST: 27134, 27135 (repository not given; presumably NIGP)>.

Genus *Pseudocatopygus* COTTEAU & GAUTHIER, 1895

In the *Treatise* (KIER, 1966: U508) this taxon is considered as synonymous to *Parapygus* POMEL, 1883

Cretaceous

P. bucharensis SHMIDT & SIMAKOV, 1953: 39-40; figs 6a-b; pl. 2: figs 7-10. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Senonian) <HT: VNIGRI 40/336> {referred to *Catopygus* by EGOROV (1972: 59)}.

Genus *Rostropygus* SZÖRÉNYI, 1955

Rostropygus SZÖRÉNYI, 1955: 67-68, 206. Type-species: *Rostropygus annae* SZÖRÉNYI, 1955. [Hungary] (Mid-Cretaceous) {In the *Treatise* (KIER, 1966: U508) this taxon is considered as synonymous to *Parapygus* POMEL, 1883}.

Cretaceous

R. annae SZÖRÉNYI, 1955: 68, 206-207; pl. 9, figs 1-4, 6. [groupe de calcaires à *Hippurites*, Sümeg (carrière située sur la partie N de la colline Kövesdomb), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/259>.

Family Faujasiidae LAMBERT, 1905

Genus *Domechinus* KIER, 1962

Late Cretaceous

D. sinensis MU & WU, 1976: 365; pl. 4: figs 8-13. [Zongshan Fm., north of Zongshan (= Mt. Zong), Gangba County (south slope of Jidula), Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <HT (by monotypy): 27138 (repository not given; presumably NIGP)>.

D. vialovi MOSKVIN, 1984: 65-67; figs 2a-b; pl. 1: figs 1a-e, 2. [Central Kyzylkumy, Tokhtatau Mt., Uzbekistan] (Maastrichtian, Late Cretaceous) <HT: MGU 242/1>.

Genus *Faujasia* D'ORBIGNY, 1856

Late Cretaceous

[*F. praeacutus* EGOROV, 1972]: 61. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

Genus *Gongrochanus* KIER, 1962

Maastrichtian

G. ariyalurensis SRIVASTAVA, 2003: 60-63; pl. 1, figs 1-8. [Kallankurichchi Fm. and Ottakovil Fm., 3.2 km E of Ariyalur, Tamil Nadu, India] (Maastrichtian) <HT: TCE 151; PT: TCE 155, 158, 160, 161, 163 to 167>.

G. chiplonkari BADVE & AZIZ, 1983: 237; figs 3d, e. [Ottakovil Fm., Ariyalun Grp., Ottakovil Area, South India (Lower to Middle Maastrichtian) <HT: MACSG 1439; PT: MACSG 1440-1447>.

G. stoliczkai BADVE & AZIZ, 1983: 237-238; figs 3f, g-j. [Ottakovil Fm., Ariyalun Grp., Ottakovil Area, South India (Lower to Middle Maastrichtian) <HT: MACSG 1448; PT: MACSG 1449-1456>.

G. kieri BADVE & AZIZ, 1983: 238; figs 3h, i. [Ottakovil Fm., Ariyalun Grp., Ottakovil Area, South India (Lower to Middle Maastrichtian) <HT: MACSG 1457; PT: MACSG 1458-1466>.

G. ottakoviensis BADVE & AZIZ, 1983: 239; figs 4a, b. [Ottakovil Fm., Ariyalun Grp., Ottakovil Area, South India (Lower to Middle Maastrichtian) <HT: MACSG 1467; PT: MACSG 1468-1478>.

G. circularis BADVE & AZIZ, 1983: 239; figs 4c, d, e. [Ottakovil Fm., Ariyalun Grp., Ottakovil Area, South India (Lower to Middle Maastrichtian) <HT: MACSG 1477; PT: MACSG 1478-1486>.

G. tamilnaduensis BADVE & AZIZ, 1983: 240; figs 4 f, g. [Ottakovil Fm., Ariyalun Grp., Ottakovil Area, South India (Lower to Middle Maastrichtian) <HT: MACSG 1487; PT: MACSG 1488-1528>.

Late Cretaceous, undifferentiated

G.? *acutus* MU & WU, 1976: 366; pl. 5: figs 5-8. [Zongshan Fm., Dongripu, Gangba County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <HT (by monotypy): 27141 (repository not given; presumably NIGP)>.

Genus *Hardouinia* HAIME in D'ARCHIAC & HAIME, 1853

Late Cretaceous

H. nuratensis MOSKVIN, 1984: 67-68; figs 3a-b; pl. 1: figs 3a-д, 4. [Western forehills of the southern Nuratau Mts., Uzbekistan] (Maastrichtian) <HT: MGU 242/3>.

Genus *Himalayechinus* MU & WU, 1976

Himalayechinus MU & WU, 1976: 366. Type-species: *Himalayechinus typicus* MU & WU, 1976 [China] (Late Cretaceous) {probably synonymous to *Faujasia* (A. SMITH, pers. comm. Nov. 2007); other species included: *Himalayechinus longus* MU & WU, 1976; *Himalayechinus minimus* MU & WU, 1976}.

Late Cretaceous

H. longus MU & WU, 1976: 367; pl. 5: figs 18-21. [Zongshan Fm., north of Zongshan (= Mt. Zong), Gangba County (south slope of Jidula), Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <HT (by monotypy): 27144 (repository not given; presumably NIGP)>.

H. minimus MU & WU, 1976: 367; pl. 6: figs 1-6. [Zongshan Fm., north of Zongshan (= Mt. Zong), Gangba County (south slope of Jidula), Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27145, 27146 (repository not given; presumably NIGP)>.

H. typicus MU & WU, 1976: 366-367; pl. 5: figs 9-17. [Zongshan Fm., north of Zongshan (= Mt. Zong), Gangba County (south slope of Jidula), Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27142, 27143 (repository not given; presumably NIGP)>.

Genus *Petalobrissus* LAMBERT, 1916

SMITH (1995: 202) placed this genus in the family Cassidulidae AGASSIZ & DESOR, 1847.

Maastrichtian

P. rawdahensis SMITH, 1995: 202-206; fig. 64-65; pl. 24: figs 1-12. [Jebel Rawdah, Oman/ United Arab Emirates Border Region] <HT: BMNH EE3485; PT: BMNH EE3467-84, EE3486-87, EE4321-22> {SMITH & JEFFERY (2000:174) placed this species in the subgenus *Paralampas* DUNCAN & SLADEN, 1882}.

P. (Paralampas) platisternus SMITH & JEFFERY, 2000: 172-174; text-fig 71A-E. [Jebel Rawdah, Oman-United Arab Emirates border region] (*A. fresvillensis* zone, mid Maastrichtian) <HT: BMNH EE4341; PT: BMNH EE3505-10, EE3512-28, EE3530, EE3532, EE3535, EE3537-41>.

Late Cretaceous

[*P. tadjikistanensis* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

Genus *Progongrochanus* AZIZ & BADVE, 2001

Progongrochanus AZIZ & BADVE, 2001: 45. Type-species: *Progongrochanus ariyalurensis* AZIZ & BADVE, 2001. [India] (Albian to Campanian, Cretaceous) {other species included: *Progongrochanus bellus* AZIZ & BADVE, 2001, *Cassidulus crassus* STOLICZKA, 1873, *Cassidulus oldhaminus* STOLICZKA, 1873, *Nucleolites (Pygorhynchus) plantus* FORBES, 1846, *Nucleolites (Pygorhynchus) testudo* FORBES, 1846}.

Maastrichtian

P. ariyalurensis AZIZ & BADVE, 2001: 45-46; pl. 1: figs 1-3. [Kallankurichchi Formation, Ariyalur Group, Kallankurichchi, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Hauericeras rembda*-zone, Early Maastrichtian) <HT: MACSG-1951; PT: MACSG-1952-1969>.

P. bellus AZIZ & BADVE, 2001: 46-47; pl. 1: figs 4-5. [Kallankurichchi Formation, Ariyalur Group, Kallankurichchi, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Hauericeras rembda*-zone, Early Maastrichtian) <HT: MACSG-1970; PT: MACSG-1971-2000>.

Genus *Stigmatopygus* D'ORBIGNY, 1856

SMITH & WRIGHT (2000: 435) placed this genus in the subfamily Stigmatopyginae SMITH & WRIGHT, 2000.

Maastrichtian

S.? *pulchellus* SMITH, 1995: 212-215; fig. 69-70; pl. 27: figs 1-8. [Jebel Rawdah, Oman/United Arab Emirates Border Region] <HT: BMNH EE4314; PT: BMNH EE3324-25, EE3329-30, EE3332-33, EE4312-13>.

Subfamily Stigmatopyginae SMITH & WRIGTH, 2000

Stigmatopyginae SMITH & WRIGTH, 2000: 435. Type-genus: *Stigmatopygus* D'ORBIGNY, 1856. (Middle Jurassic (Bajocian) to Upper Eocene) {other genera included: *Gongrochanus* KIER, 1962; *Hardouinia* HAIME in D'ARCHIAC & HAIME, 1853; *Platypygus* DE LORIO, 1902; *Procassidulus* LAMBERT, 1918; and *Ochetus* POMEL, 1883}.

Family Archiaciidae COTTEAU & TRIGER, 1869

Genus *Archiacia* L. AGASSIZ, in AGASSIZ & DESOR, 1847

Cretaceous

- A. hungarica* SZÖRÉNYI, 1955: 82-87, 221-224; figs 26-33; pl. 11, figs 12-36. [groupes de manes argileuses, Zirc-Tündérmajor, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/348>.
- A. magna* SZÖRÉNYI, 1955: 87-88, 224; pl. 11, figs 37-41. [calcaire à Réquiénies, Zirc-Tündérmajor, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/398>.

Genus *Acriaster* SMITH in SMITH & BENGGTSON, 1991

Acriaster SMITH in SMITH & BENGGTSON, 1991: 46. Type-species: *Acriaster sergipensis* SMITH in SMITH & BENGGTSON, 1991. [Brazil] (Albian).

Albian

- A. sergipensis* SMITH in SMITH & BENGGTSON, 1991: 47-48; figs 38-39; pl. 8, figs H-K. [?Maruim Mb., Riachuelo Fm., Brazil] (?Upper Albian) <HT: USNM 448097>.

Family Cassidulidae L. AGASSIZ & DESOR, 1847

Genus *Cardiopygus* AZIZ & BADVE, 2001

Cardiopygus AZIZ & BADVE, 2001: 49. Type-species: *Cardiopygus cardus* AZIZ & BADVE, 2001. [India] (Maastrichtian).

Maastrichtian

C. cardus AZIZ & BADVE, 2001: 49-50; pl. 2: figs 1-3. [Kallankurichchi Formation, Ariyalur Group, Ariyalur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Hauericeras rembda*-zone, Early Maastrichtian) <HT: MACSG-2001>.

Genus *Cassidulus* LAMARCK, 1801

Eocene

C. santolayae SILLERO in SANTOLAYA & SILLERO, 1994: 21-22; unnumbered fig. on p. 22. [Sierra de Horna, Province Alicante, Spain] (Lutetian, Middle Eocene) <HT: Collection Sillero, Alicante> {re-described in LÓPEZ & SILLERO (2006: 162)}.

Paleocene

C. kieri ADEGOKE, 1977: 58-60; pl. 5: figs 1-14. [Ewekoro Fm., Ewekoro quarry, Ewekoro, 50 km N Lagos, Nigeria] (Paleocene) <HT: UIMG 145; PT: UIMG 164-165, USNM 174760-174762>.

Genus *Hypsopygaster* BAJARUNAS, 1915

Paleocene

H. bajarunasi KADIL'NIKOVA & MOSKVIN, 1984: 37; pl. 6, figs 7. [Sullukapinskaya Fm., Mangyshlak Peninsula, Mt. Aksyrtau, U.S.S.R.] (Upper Paleocene) <HT: PIN 4011/7> {pagination of English translation, Russian original not seen}.

H. rostriformis KADIL'NIKOVA & MOSKVIN, 1984: 36; pl. 6, figs 5-6. [Western Mangyshlak region, Kapam ravine, U.S.S.R.] (Lower Paleocene) <HT: PIN 4011/5> {pagination of English translation, Russian original not seen}.

Genus *Limpasiaster* AZIZ & BADVE, 2001

Limpasiaster AZIZ & BADVE, 2001: 50. Type-species: *Limpasiaster ariyalurensis* AZIZ & BADVE, 2001. [India] (Maastrichtian) {other species included: *Limpasiaster pentagonalis* AZIZ & BADVE, 2001; *Limpasiaster quadralis* AZIZ & BADVE, 2001}.

Late Cretaceous

L. ariyalurensis AZIZ & BADVE, 2001: 50-51; pl. 2: figs 4-6. [Kallankurichchi Formation, Ariyalur Group, Ariyalur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Hauericeras rembda*-zone, Early Maastrichtian) <HT: MACSG-2002; PT: MACSG-2003-2008>.

- L. pentagonalis* AZIZ & BADVE, 2001: 52; pl. 3: figs 4-6. [Kallankurichchi Formation, Ariyalur Group, Ariyalur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Hauericeras rembda*-zone, Early Maastrichtian) <HT: MACSG-2059; PT: MACSG-2060-2098>.
- L. quadralis* AZIZ & BADVE, 2001: 51-52; pl. 3: figs 1-3. [Kallankurichchi Formation, Ariyalur Group, Ariyalur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Hauericeras rembda*-zone, Early Maastrichtian) <HT: MACSG-2009; PT: MACSG-2010-2058>.

Genus *Nucleopygus* L. AGASSIZ, 1840

Paleocene

- N. salgadoi* PARMA, 1989: 94-96; pl. 1: figs 1-6. [Roca Formation, Casa de Piedra, at the border between the Río Negro and La Pampa provinces, Argentina] (Paleocene) <HT: MOZ P 3020>.

Maastrichtian

- N. magnus* SMITH, 1995: 216-219; fig. 71-72; pl. 28: figs 1-7. [Simisima Fm., Jebel Bu-hays, Jebel Faiyah, Jebel Rawdah and Jebel Thanais, Oman/United Arabian Emirates Border Region] <HT: BMNH EE4339; PT: BMNH EE3340, EE3356, EE3358, EE3363, EE3365, EE3367-68, EE4327, EE4335-38>.

Late Cretaceous

- N. bashibulakensis* YANG SHENGQIU, 1991: 117-118; pl. 11, figs 1-13. [Upper Kukebai Fm., Kuzigongsu, 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Late Cretaceous) <HT: NIGP 88371>.
- N. gangbensis* MU & WU, 1976: 368; pl. 7: figs 1-9. [Zongshan Fm., north of Zongshan (= Mt. Zong), Gangba County (south slope of Jidula), Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27151, 27152 (repository not given; presumably NIGP)>.
- N. platypetalus* YANG SHENGQIU, 1991: 118-119; pl. 12, figs 1-9. [Upper Kukebai Fm., Kuzigongsu, 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Late Cretaceous) <HT: NIGP, no holotype defined>.
- N. calcalosus* YANG SHENGQIU, 1991: 118; pl. 13, figs 1-12. [Upper Kukebai Fm., Kuzigongsu, 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Late Cretaceous) <HT: NIGP, no holotype defined>.

Mid-Cretaceous

- N. parvus* SZÖRÉNYI, 1955: 74-75, 213; pl. 10, figs 6-8. [marne glauconieuse, carrière située à côté de la route Bakonynána-Alsóperre, and Péntzeskut-Körisgyörpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/318 (from Bakonynána-Alsóperre)>.
- N. peltitipos* SZÖRÉNYI, 1955: 75-76, 213-214; pl. 10, figs 9-13. [marne glauconieuse, Bakonynána, and Péntzeskut-Körisgyörpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/322 (from Péntzeskut-Körisgyörpuszta)>.

Genus *Ochetus* POMEL, 1883[=*Ochetes* KIER, 1962]

SMITH & WRIGHT (2000: 435) placed this genus in the subfamily Stigmatopyginae SMITH & WRIGHT, 2000, within the family Faujasiidae LAMBERT, 1905.

Cenomanian

- O. pauli* SMITH & WRIGHT, 2000: 439; text-fig. 173; pl. 138, figs 10-12. [White Hart Pit, Wilmington, Devon, England] (*N. carcitanense* zone, Lower Cenomanian) <HT: BMNH EE6377; PT: BMNH EE80644>.

Genus *Paralampas* DUNCAN & SLANDEN, 1882

In the *Treatise* (KIER, 1966: U515) this taxon is considered as synonymous to *Rhynchopygus* D'ORBIGNY, 1856.

Paleocene

- P. rancureli* TESSIER & ROMAN, 1973: 149-151; fig. 8; pl. 1, figs 8-11; pl. 3, fig. 2. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Paleocene, Thanetian) <HT: IPM 1972-9Cc>.

Genus *Rhyncholampas* A. AGASSIZ, 1869

Miocene

- R. candidoi* GARRAFIELO FERNANDES & CARREIRA MORAIS, 1994: 55-57; figs 1-4. [Pirabas Formation, Praia de Atalaia, Município de Salinópolis, Estado do Pará, Brazil] (Early Miocene) <HT: MPEG 1160>.
- R. chipolanus* OYEN & PORTELL, 1996: 62-66; pl. 1, figs 1a-b; pl. 2, figs 2a-b. [Chipola Fm., Chipola River, Calhoun County, Florida] (Early Miocene) <HT: UF 66633>.

Oligocene

R. gouldii newbernensis KIER, 1997: 6; fig. 3; pl. 5, figs 1-7. [Trent Fm., New Bern quarry, North Carolina] (Late Oligocene) <HT: USNM 398324; PT: USNM 398325, 492065-492096>.

Eocene

R. smithi SRIVASTAVA, SINGH, TIWARI & JAUHRI, 2008: 514-515; pl. 2: figs 1-8; pl. 3: figs 5-8. [Siju Limestone Dhapsagiri village, South Garo Hills, Meghalaya, India] (Late Lutetian to Early Bartonian) <HT: LUGD I/2023; PT: LUGD I/2024 to I/2026>.

R. strougoi AZAB & ELATTAAR, 1999: 854; tab. 10; pl. 1: figs 22-26. [El Guss Abu Said, Farafra area, Egypt] (Prenaster alpinus Zone, Early Libyan, Early Eocene) <repository and specimen nos. not given>.

Subgenus *Galerolampas* COTTEAU, 1889

In the *Treatise* (KIER, 1966: U515) this taxon is considered as synonymous to *Rhyncholampas* A. AGASSIZ, 1869.

Paleocene

R. (G.) tinrhertensis AMARD, COLLIGNON & ROMAN, 1983: 93-95; fig. 10; pl. 17, figs 1-6. [Oudeiat Chouikh, Tinrhert occidental, Algeria] (Late Paleocene) <HT: MNHN 1975-15 AB 202-8>.

Genus *Rhynchopygus* D'ORBIGNY, 1856

Eocene

R. ? janchrisorum HOLMES, 2004: 211-213; figs 2A-F, 3A, B, 4A-C, 5, 6A. [Muloowurtie Fm., "Sliding Rocks", Yorke Peninsula, South Australia] (Aldingian, Priabonian, Late Eocene) <HT: NMV P145616; PT: NMV P312113-P312115>.

Upper Cretaceous

R. arumaensis KIER, 1972a: 72-73; pl. 45, figs 1-77. [Aruma Fm., locality KK11, Saudi Arabia] (Campanian) <HT: USNM 170452; PT: USNM 170453>.

Family Clypeolampadidae KIER, 1962

Genus *Vologesia* COTTEAU & GAUTHIER, 1895

Maastrichtian

V. rowdahensis ALI, 1989a: 406; figs 5(1-3). [Simsima Fm., Gebel El Rowdah, United Arab Emirates] (Late Maastrichtian) <MGD-UAA > {no type specimen defined}.

Family Pliolampadidae KIER, 1962

Genus *Calilampas* SQUIRE & DEMETRION, 1996

Calilampas SQUIRES & DEMETRION, 1996: 512-514. Type-species: *Calilampas californiensis* SQUIRE & DEMETRION, 1996. [USA] (Eocene).

Eocene

C. californiensis SQUIRE & DEMETRION, 1996: 514, fig. 5.1-5.7, 6, 7. [Bateque Fm., Baja California Sur, Mexico; Llajas Fm., Simi Valley, Southern California, USA] (Middle Lower Eocene) <HT: IGM 6387; PT: IGM 6388-6391, LACMIP 11532>.

Genus *Eurhodia* HAIME in D'ARCHAIC & HAIME, 1853

MOOI (1990a: 693-695) placed this genus in the family Cassidulidae L. AGASSIZ & DESOR, 1847

Recent

E. relictata MOOI, 1990: 690-693; figs 1-4. [continental shelf off Surinam, South America] <HT: USNM E20480; PT: USNM E12971>.

Eocene

E. baumi KIER, 1980: 23-24; fig. 6; pl. 6, figs 5-8. [Castle Hayne Limestone, Rose Hill locality 11, North Carolina, USA] (Middle Eocene) <HT: USNM 264043>.

E. rugosa idealis KIER, 1980: 26-28; figs 7, 9; pl. 7, figs 7-9. [Castle Hayne Limestone, localities 2, 5, 8, Rose Hill locality 11, Ideal Cement Company quarry localities 12, 26, 28, North Carolina, USA] (Middle Eocene) <HT: USNM 264046>.

E. rugosa depressa KIER, 1980: 28-29; figs 7, 10; pl. 7, figs 10-12. [Castle Hayne Limestone, Maple Hill (East Coast Construction Company quarry) localities 10, 34, North Carolina, USA] (Middle Eocene) <HT: USNM 264047>.

Genus *Gitolampas* GAUTHIER, 1889

Eocene

G. mcnamarae SRIVASTAVA, SINGH, TIWARI & JAUHRI, 2008: 515-516; pl. 3: figs 1-4. [Siju Limestone Dhapsagiri village, South Garo Hills, Meghalaya, India] (Late Lutetian to Early Bartonian) <HT: LUGD I/2027>

Genus *Pseudopygaulus* COQUAND, 1862

Eocene

P. romani AZAB & ELATTAAR, 1990: 216-221; figs 3-5; pl. 1: figs 1-15. [Wadi Abu Gelbana, Gebel Haridi, Wadi Matmar, Wadi El Gabrawi, Tell El-Amarna, and Beni Hassan, Nile Valley, between Minia and Sohag, Egypt] (Late Libyan, Early Eocene) <HT: specimen illustrated in pl. 1: figs 1-7 (from Wadi Abu Gelbana), no repository given>.

Paleocene

P. malavoyi TESSIER & ROMAN, 1973: 152-155; figs 9-14; pl. 1, figs 15-18; pl. 2, figs 1-3; pl. 3, fig. 3. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Paleocene, Thanetian) <HT: IPM 1972-9Df>.

Genus *Studeria* DUNCAN, 1891

Paleocene

S. ? frescoensis TESSIER & ROMAN, 1973: 155-156; pl. 1, figs 12-14; pl. 3, fig. 4. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Paleocene, Thanetian) <HT: IPM 1972-9Ea>.

Genus *Tamililampus* AZIZ & BADVE, 2001

Tamililampus AZIZ & BADVE, 2001: 52-54. Type-species: *Tamililampus tumidus* AZIZ & BADVE, 2001. [India] (Maastrichtian).

Maastrichtian

T. tumidus AZIZ & BADVE, 2001: 54; pl. 3: figs 7-8. [Kallankurichchi Formation, Ariyalur Group, Ariyalur, Tamil Nadu region, Trichinopoly Basin, Southern India] (Hauericeras rembda-zone, Early Maastrichtian) <HT: MACSG-2009 (probably type error for 2099 as no. 2009 is mentioned as holotype of *Limpasiaster quadralis* AZIZ & BADVE, 2001 also)>.

Family Apatopygidae KIER, 1962

Genus *Apatopygus* HAWKINS, 1920

Miocene

A. mannumensis HOLMES, 1999: 57-60; figs 3B, 3E, 4A-E, 7B, 7E. [Mannum Fm., Murray River, South Australia] (Longfordian, Early Miocene) <HT: NMV P148496; PT: NMV P148385-148395>.

Oligocene

A. gaudensis GATT, 2005: 104-109; figs 1, 1a, 2, 2a, 3, 3a, 4, 5. [Lower Coralline Limestone Fm. and Lower Globigerina Limestone Formation, Qammieh, northwestern Malta; numerous sites along the southern and western coast of Gozo, Maltese Islands, Central Mediterranean] (Late Chattian, Oligocene to Aquitanian, Early Miocene*) <HT: Michael Gatt coll. (Rabat, Malta) E.574; PT: Michael Gatt coll. E.1695, E.1701, E.1705> {*According to the revised age dates FORESI et al. (2008) the Lower Globigerina Limestone Formation is of Late Chattian age, thus restricting the occurrence of *A. gaudensis* GATT, 2005 to the Late Oligocene}.

Eocene

A. vincentinus (TATE, 1891): 280. [reported as coming from “Aldinga”, but according to HOLMES (1999: 55) they probably come from the Tortachilla Limestone, north of Blanche Point, Maslin Bay, South Australia] (Johannian – Aldingan, Middle to early Late Eocene) <LT: SAM T266H; PLT: SAM T266A-G, I, J> {originally referred to the genus *Echinobrissus*; not mentioned in LAMBERT & THIÉRY (1909-25); revised by HOLMES (1999), who designated the lectotype}.

Genus *Porterpygus* BAKER, 1983

Porterpygus BAKER, 1983: 164. Type-species: *Porterpygus kieri* BAKER, 1983. [New Zealand] (Recent).

Recent

P. kieri BAKER, 1983: 164-168; figs 3, 10-28. [Three Kings Is., New Zealand] (Recent) <HT: NMNZ Ech. 3707; PT: NMNZ Ech. 3708-10>.

Miocene

P. devlinensis HOLMES, 1999: 60-64; figs 3C, 6A-G, 7C, 7F. [Mannum Fm., Murray River, South Australia] (Longfordian, Early Miocene) <HT: NMV P148399; PT: NMV P148396-148398, 148400 and 148401>.

Family Claviasteridae ALI, 1992a

Claviasteridae ALI, 1992: 72. Type-genus: *Claviaster* D'ORBIGNY, 1855. [Northern Africa and France] (Late Cretaceous).

Family Plesiolampadidae SMITH & JEFFERY, 2000

Plesiolampadidae SMITH & JEFFERY, 2000: 246-247. Type-genus: *Plesiolampas* DUNCAN & SLADEN, 1882. {other genera included: *Oriolampas* MUNIER-CHALMAS, 1882; *Ter-mieria* LAMBERT, 1931; and *Pseudopygaulus* COQUAND, 1862}.

Family Clitopygidae VADET, 2007

Clitopygidae VADET, 2007: 211-212. Type-genus: *Clitopygus* POMEL, 1883. [Europe] (Jurassic).

Family Infraclypeidae SAUCÈDE, MOOI & DAVID, 2007

Infraclypeidae SAUCÈDE, MOOI & DAVID, 2007: 352-353. Type-genus: *Infraclypeus* GAUTHIER, 1875 in COTTEAU, PERON & GAUTHIER, 1873-1891. {other genera included: *Pyri-nodia* POMEL, 1883; *Desorella* COTTEAU, 1855; *Pachyclypus* DESOR, 1858}.

Family Uncertain

Genus *Echinanthus* LESKE, 1778

AMARD, COLLIGNON & ROMAN (1983: 110) placed this genus into the family Pliolampadidae.

Eocene

E. reguanti REGUANT, ROMAN & VILLATTE, 1972: 903-905; pl. 33, figs 9a-d, 10a-b, 11a-b. [Niveau m1 of REGUANT (1967) = "marnes de Banyoles", 4 km N of Sau, Region of Vic, Barcelona, Spain] (Biarritzien inférieur, Middle Eocene) <HT: MNHN 1970-6 (AA03-1)>.

Paleocene

E. schweinfurthi devriési AMARD, COLLIGNON & ROMAN, 1983: 110-112; fig. 11-12; pl. 17, figs 7-16. [Ed Dahna, Tademait oriental, Algeria] (Early Paleocene) <HT: MNHN 1975-15 AB 709-21>.

Order Holasteroidea DURHAM & MELVILLE, 1957

Genus *Smithiaster* BARRAS, 2007

Smithiaster BARRAS, 2007: 158; figs 5a, b; pl. 2: figs 8a-c. Type-species: *Dysaster loryi* GRAS, 1852. [France & North Africa] (Oxfordian to Kimmeridgian, Late Jurassic) {stem group holasteroid}.

Family Collyritidae D'ORBIGNY, 1853

Genus *Collyrites* DESMOULINS, 1835

Middle Jurassic

C. analis dimota VIALOV, 1945: 468-469; figs a-e. [Balkhan Mt., Western Turkmenistan] (Callovian, Jurassic) <HT: coll'n Vasiljevski, current repository unknown>.

C. cheveti VADET, NICOLLEAU & PINEAU, 1998: 55-56; 1 fig., pl. 17. [l'Assise de Carreaux, environs de Mortagne au Perche, Sarthe, France] (Callovian) <HT: V 5124; Type series: V 3830-31, 5123, 5126-5132>.

Genus *Collyropsis* GAUTHIER, 1896

Albian

C. ovoides DEVRIÈS, 1972: 46-47; pl. 2: figs 7-12. [Sierra de Crevilente, Province Alicante, Spain] (Albian, Late Cretaceous) <ST: coll. J. Azema, Paris [no specimen no. provided]>.

Berriasian

C. ultimus SOLOVJEV, 1971: 63-64; fig. 28a-b; pl. 10, figs 2a-b, 3a-b. [Kutsky, SE Crimea, Ukraine] (Berriasian, Early Cretaceous) <HT: PIN 2282/1>.

Genus *Orbignyana* EBRAY, 1860

Late Jurassic

O. cordiformis DEVRIÈS, 1972: 43-45; pl. 1: figs 1-5; pl. 3: figs 1-8. [northern slope of Puig Santa Magdalena, north-east of Novelda, Province Alicante, Spain] (Late Kimmeridgian/Tithonian) <HT (by monotypy): coll. J. Azema, Paris, no. 750>.

Genus *Pyrgorhytis* POMEL, 1883

Jurassic

P. klairi VADET & RÉMY, 2004: 17-19; 8 unnumbered figs on pp. 18-19. [Calcaire gréseux à oolithes ferrugineuses, environs de Sedan, Ardennes, France] (*sauzei* zone, Early Bajocian) <HT. Rémy coll. 972; PT: Rémy coll. 817-818, 922-924, 971, 1006-1009; Vadet coll. 6262-6274, 6278-6284>.

P. magnus SOLOVJEV, 1971: 47; pl. 2, figs 3a-b, 4a-f. [Janysharsky Horizon, Kara-Dag (Kordonnaya Balka), SE Crimea, Ukraine] (Middle Callovian) <HT: PIN 2275/5>.

Genus *Tetraromania* SOLOVJEV, 1971

Tetraromania SOLOVJEV, 1971: 64. Type-species: *Dysaster ovulum* DESOR, 1842. [Ukraine] (Berriasian to Barremian) {other species included: *Collyrites jaccardi* DESOR, 1869}.

Subfamily Collyropsinae SOLOVJEV, 1966

Collyropsinae SOLOVJEV, 1966: 45-46. Type-genus: *Collyropsis* GAUTHIER, 1896.

Family Disasteridae GRAS, 1848

Genus *Metaporinus* L. AGASSIZ, 1844

Middle Jurassic

M. drogiacus ROBERT, 1991: 22; pl. 5. [Calcaires à chailles; Brétignelles, près Druyes, Dept. Yonne, France] (*Antecedens* subzone, *Plicatilis* zone, Middle Oxfordian) <HT: Philippe Robert coll'n No. 604> {based on an internal mould; re-described in ROBERT (1994: 54; pl. 6: figs 2, 10); original paper not seen, based on data in ROBERT (1994)}

Genus *Proacrolusia* PORETSKAYA, 1974b

Proacrolusia PORETSKAYA, 1974b: 86-87. Type-species: *P. kelatensis* PORETSKAYA, 1974b. [Turkmenistan] (Late Jurassic).

Late Jurassic

P. kelatensis PORETSKAYA, 1974b: 87; figs a-e; pl. 1: figs 1-4. [Kopet Dag, Turkmenistan] (Late Jurassic) <HT: LGU 327/1; PT: LGU 327/2 to 327/17>.

Family Holasteridae PICTET, 1857

Genus *Cardiaster* FORBES, 1850

Cretaceous

C. bolschechetensis SHMIDT, 1976: 154-155; pl. 1: figs 3a-6, 5. [Borehole 4, Large Kheta, side brance of Jounig River, Russian SFSR] (Early Santonian, Late Cretaceous) <HT: /8846 [sic! – number of specimen appears to be incomplete, repository not given]>.

C. cotteauanus latohumilis SAVCHINSKAYA, 1974: 319-320; pl. 102, figs 1-5. [Northern Don Basin, Russian SFSR] (*Belemnitella longei*-zone, Late Campanian) <HT: not given>.

C. pseudoplanus [*protoplanus*] SMITH & WRIGHT, 2003: 488-489; text-figs 196A, 196B, 202, 203; pl. 153: figs 5-9. [“*Rhynchonella curvieri* Bed” [pars], Tingley’s Pit, Wouldham and Lydden zigzag, East of Dover, Kent, England] (I. labiatus Zone, Early Turonian and T. lata Zone, Middle Turonian, Late Cretaceous) <HT: BMNH E16899 (from Wouldham); PT: BMNH E9799> {in the abstract this species is called *Cardiaster protoplanus*}.

Genus *Crassiholaster* SMITH & WRIGHT, 2003

Crassiholaster SMITH & WRIGHT, 2003: 556. Type-species: *Spatangus subglobosus* LESKE, 1778. [Europe] (Cenomanian, Cretaceous) {other species included: *Holaster bischoffi* RENEVIER, 1868; *Echinocorys sphaericus* SCHLUETER, 1869; *Crassiholaster sulciproctus* SMITH & WRIGHT, 2003}.

Cenomanian

C. subglobosus forma *norfolkensis* SMITH & WRIGHT, 2003: 559-561; text-figs 234, 235; pl. 177: figs 13-16. [Lower Inoceramus Bed, Hunstanton, Norfolk, and Welton Parish Pit, near Alford, Lincolnshire, England] (M. dixoni Zone, Early Cenomanian, Late

Cretaceous) <HT: none defined; ST: BMNH E42595, E42620, E42622, E42651-52, E42654-56, E75478, E81606-608, E81610>.

Crassiholaster sulciproctus SMITH & WRIGHT, 2003: 563-564; text-figs 237A, B; pl. 179: figs: 12-16. [Grey Chalk facies, Dover, Kent, England] (Middle or Late Cenomanian, Late Cretaceous) <HT: BMNH E38901> {based on a single specimen}.

Genus *Echinocorys* LESKE, 1778

Paleocene

E. ancileformis MOSKVIN & SHIMANSKAYA, 1981: 135-136; fig. 2a-r; pl. 1, fig. 3; pl. 2, figs 1a-6. [west Ustyurt, Kazakhstan] (Upper Paleocene) <HT: MSU 226/7>.

E. australis FOSTER & PHILIP, 1978: 815-816; text-fig. 5; pl. 92: figs 1, 3, 6; pl. 93: figs 2, 4. [Wadera Calcarenite, Toothawarra Creek, Carnarvon Basin, Western Australia] (Middle to Late Paleocene) <CPC F4818>.

E. kongieli MOSKVIN & SHIMANSKAYA, 1993: 56; figs 2з; 3H; 4: 1a-b. [Michurino, Crimea, Ukraine] (Danian) <HT: MSU 267>.

E. schafferi KÜHN, 1930: 551; pl. 1: fig. 3. [Bruderndorf Formation, Haidhof near Ernstbrunn, Lower Austria, Europe] (Upper Danian) <HT: NHMW 1930V11> {placed into the synonymy of *Echinocorys scutata* forma *ovata* (LESKE, 1778) by KROH (2001: 398-399)}.

E. schwetzeri MOSKVIN & SHIMANSKAYA, 1981: 134-135; fig. 2e, з; pl. 1, figs 2a-6. [Urta Mt., Western Georgia] (Upper Paleocene) <HT: MSU 226/4>.

E. sulcata cognatus MOSKVIN & SHIMANSKAYA, 1981: 129-132; fig. 1a-д; pl. 1, figs 1a-b. [Usak, Kazakhstan] (Lower Paleocene) <HT: MSU 226/1>.

E. sulcata orbiculata MOSKVIN & SHIMANSKAYA, 1993: 54; figs 5: 2a-b. [Tus-Bair, Mangyshlak, Kazakhstan] (Danian) <HT: MSU 268>.

E. turanica MOSKVIN & SHIMANSKAYA, 1981: 136-137; fig. 2д-ж; pl. 2, figs 1a-b, 3. [Ustyurt, Kazakhstan] (Upper Paleocene) <HT: MSU 226/10>.

Maastrichtian

E. balcanicus TZANKOV, 1984: 90-91; pl. 38, figs 2, 2a-b; pl. 50, figs 2, 2a-b. [Dermantzi, de Vratza, Bulgaria] <HT: USC CR₂ 1292>.

E. elongatus TZANKOV, 1984: 92; pl. 39, figs 2, 2a-b. [Drumevo, de Provadia, Bulgaria] <HT: USC CR₂ 1294>.

E. kharagoulensis GONGADZE, 1976: 39-41; pl. 1: figs 1a-c. [southwestern part of the Kharagouli [Charagoulski] Syncline, southern border of the Dzirula Massif, Georgia]

(Late Maastrichtian, Cretaceous) <HT: TGU Gongadze Coll'n 4/26> {re-described in GONGADZE (1979: 80-81; pl. 12, figs 1a-b)}.

E. stomias McNAMARA, 1987b: 421-425; figs 1-3. [Miria Fm., Giralia Range, Western Australia] (Late Maastrichtian) <HT: WAM 84.442; PT: WAM 82.3088, 84.420, 84.441, 84.443, 86.1388, NMV P102120, P102398, RUCA 20152>.

E. terminata MOSKVIN & SHIMANSKAYA, 1997: 70-71; figs 1: 1a-б, 2, 3, 4a-б; figs 2a-b. [Mangyshlak, Kazhakstan] (Late Maastrichtian, Late Cretaceous) <HT: MSU 269(1)>.

Campanian

E. zejszneri MACZYŃSKA, 1984: 454-455; pl. 206, fig. 1a-d; pl. 207, fig. 1a-d. [Poskwitów koło Krakowa, Miechów Trough, Outer Carpathians, Poland] (Campanian) <HT: MZ VIII/Ee-854> {see also MACZYŃSKA in MALINOWSKA (1989: 312)}.

Cenomanian

E. euxinus SHIMANSKAYA, 1974: 151-152; figs 1a-д, 2a-г. [Mount Sel'Bulkhra, SW Crimea] (Cenomanian, Late Cretaceous) <HT: MSU 1/38>.

Genus *Eoholaster* SOLOVJEV, 1989

Eoholaster SOLOVJEV, 1989: 150. Type-species: *Eoholaster poslavskae* SOLOVJEV, 1989. [Ukraine] (Berriasian) {other species included: *Toxaster laffittei* DEVRIES, 1960}.

Berriasian

E. poslavskae SOLOVJEV, 1989: 150-153; fig. 1a-д; unnumbered pl., figs 1a-г, 2a-г, 3a-b. [south-western Crimea, Ukraine] (Berriasian) <HT: PIN 2276/7>.

Genus *Galeola* QUENSTEDT, 1874

In the *Treatise*, WAGNER & DURHAM (1966: U528) considered *Galeola* a subjective synonym of *Echinocorys* LESKE, 1778.

Campanian

G. papillosa basiplana ERNST, 1971: 219-222; Text-fig. 20a, 21, 23 (fig. 2-6). [Teutonia quarry, Misburg, near Hannover, northern Germany] (Early Campanian) <HT: Coll. Ernst; presumably in the MBE collection now (2005)>.

Genus *Garumnaster* LAMBERT, 1906

Paleocene

G. lamberti KÜHN, 1930: 552; pl. 1: figs 6-7. [Bruderndorf Formation, Bruderndorf, Lower Austria, Europe] (Upper Danian) <HT: NHMW 1930V9> {placed into the synonymy of *Echinocorys scutata* forma *pyrenaica* SEUNES, 1888 by KROH (2001: 400)}

Genus *Giraliaster* FOSTER & PHILIP, 1978

Giraliaster FOSTER & PHILIP, 1978: 804-805. Type-species: *Giraliaster jubileensis* FOSTER & PHILIP, 1978. [Australia & New Zealand] (Palaeogene).

Oligocene

G. bellissae FOSTER & PHILIP, 1978: 813-814; pl. 91: figs 4-6. [Prydes Gully Mb., Otekaike Limestone, North Otago, New Zealand] (Duntroonian-Waitakian, Late Oligocene) <HT: NZGS EC849>.

Paleocene

G. jubileensis FOSTER & PHILIP, 1978: 805-808; text-figs 3, 4a; pl. 89: figs 1-6. [Cardabia Grp., Jubilee Bore, Carnarvon Basin, Western Australia] (Middle or Late Paleocene) <HT: WAM 73.362; PT: WAM 73.363-64>.

Genus *Guettaria* GAUTHIER, 1888

Maastrichtian

G. fecunda SCHULZ, 1983: 718; fig. 2d; pl. 1, figs 2-3. [Moos-Graben, Siegsdorf, Bavaria, Germany] (Late Early Maastrichtian) <HT: BSPG 1977 XXI/22>.

Campanian

G. schamchorensis MELIKOV, in ALI-ZADE, 1988: 198; figs 35-37; pl. 66, figs 2a-r. [Schamchor, Lower Caucasus, Azerbaijan] (Late Campanian) <HT: AzINEFTEK-HIM 201/2>.

Genus *Hagenowia* DUNCAN, 1889

Santonian

H. blackmorei anterior ERNST & SCHULZ, 1971: 140-141; figs 8: 1a-c, 2a-c; pl. 14, figs 2-3. [Breitenburg-Schinkel pit, Lägerdorf, Northern Germany] (Base of the *rogalae*/

westfalicagranulata-zone, Middle Santonian) <HT: GPIH 121; PT: coll. Diersche and coll. Schulz>.

Genus *Hemipneustes* AGASSIZ, 1836

Maastrichtian

H. arabicus ALI, 1989a: 408; figs 6(1-3), 7. [Simsima Fm., Gebel El Rowdah, United Arab Emirates] (Late Maastrichtian) <MGD-UAA > {no type specimen defined}.

Late Cretaceous

H. duncani SAWAR, 1989: 376-380; figs 1a-c, 2a-c. [Laki Hills, District Dadu, and Nathiagali, District Abbottabad, Khithar Range, Sind, N.W.F.P., Pakistan] (Late Cretaceous or Eocene) <HT: PUPCE No. 63 (from the Laki Hills)> {based on 2 poorly preserved specimens}.

H. indicus AZIZ & BADVE, 1990: 326-330; figs 2a-d. [Kallankurichchi Formation, Ariyalur Group, Ariyalur (11°09' N, 79°07'30" E), Tiruchirapalli District, Tamil Nadu region, Trichinopoly Basin, Southern India] (Maastrichtian, Late Cretaceous) <HT: MACSG-2104; PT: MACSG-2105-2107>.

Genus *Holaster* L. AGASSIZ, 1836

Late Cretaceous

[*H. sanglakensis* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

H. tanamensis SHMIDT, 1976: 153-154; pl. 1: figs 1a-b, 2a-b, 4a-b. [Krasnojarskiy Kr., Tanama Valley, near mouth of Yennisey [Enisei] River, Russian SFSR] (Early Santonian, Late Cretaceous) <HT: I/8846 [repository not given]>.

Mid-Cretaceous

H. hungaricus SZÖRÉNYI, 1955: 93-95, 229-230; fig. 37; pl. 12, figs 10-12; pl. 13, figs 8-9. [marne à *Turrilites*, Olaszfalu, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/440>.

H. pseudonodulosus SZÖRÉNYI, 1955: 95, 230-231; pl. 13, figs 1, 3-7. [marne à *Turrilites*, Jásd, and Lókút, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/412 (from Lókút)>.

H. subquadratus SZÖRÉNYI, 1955: 95-96, 231; pl. 13, figs 10-13. [marne glauconieuse e marne à *Turrilites*, Pénezskut-Körisyörpuszta, Lókút, and Kisgyón-

Rákoshegy, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/434 (from Pénzeskut-Körisgyörpuszta)>.

Genus *Infulaster* WRIGHT & WRIGHT, 1948

Late Cretaceous

I. praecursor SMITH & WRIGHT, 2003: 513-514; text-figs 210, 211; pl. 157: figs 1-5. [Kiplingcotes, pit 1, and Drymere pit, near Swaffham, Norfolk, England] (*S. plana* Zone, Late Turonian, Late Cretaceous) <HT: BMNH EE6771 (from Kiplingcotes); PT: BMNH EE6762, EE6763, EE6766, EE6768>.

Genus *Lampadocorys* POMEL, 1883

Late Cretaceous

L. ennisi SMITH & WRIGHT, 2003: 566-567; text-fig. 238; pl. 182: figs 1-5. [Lower Chalk, Speeton, Yorkshire, England] (Late Cenomanian, Late Cretaceous) <HT: BMNH EE6952; PT: EE6730>.

L.? *estermanni* KROH & JAGT, 2004: 553-556; text-figs 3b, c, e,f; pl. 1: figs 1-6; pl. 2, figs 1-6. [Gschiefe Graben tectonic window, near Gmunden, Upper Austria] (standard nannoplankton zones CC18–CC23, Middle to Late Campanian) <HT: NHMW 2003z0067/0001; PT: 2003z0067/0002>.

Genus *Medjesia* JEFFERY, 1997b

Medjesia JEFFERY, 1997b: 244-247. Types-species: *Enallopneustes meslei* GAUTHIER, 1892. [Tunisia] (Cenomanian).

Genus *Offaster* DESOR, 1858

Maastrichtian

O. granulatus KUTSCHER, 1978a: 630-631; pl. 3, figs 1-4. [Saßnitz, Rügen, Germany] (Late Early Maastrichtian) <HT: SGWG 59> {renamed *O. rugius* REICH, VILLIER & KUTSCHER, 2004 because of primary homonymy with *Offaster granulatus* LAMBERT, 1931}.

O. rugius REICH, VILLIER & KUTSCHER, 2004: 499. [Complex V, north of Saßnitz, Jasmund Peninsula, Rügen, Germany] (Late Early Maastrichtian) <HT: SGWG 59> {nomen novum for *Offaster granulatus* KUTSCHER, 1978a, p. 630 non LAMBERT, 1931}.

Santonian

O. nuciformis ERNST, 1971: 209-214; Text-fig. 17a-f, 18 (figs 1a-d, 2a-e). [Bülten-Adenstedt quarry, near Peine, northern Germany] (Late Mid-Santonian) <HT: GPIH type catalogue no. 122>.

Cretaceous

O. granulosis LAMBERT, 1931: M3-M4; pl. 1, figs 14-15. [Road from Zougouldak to Devrek and Marnes de Bartine near Guv-Tepessi, région d'Héraclée, Anatolia, Turkey] (Cretaceous) <HT: not given>.

Genus *Plesiocorys* POMEL, 1883

Late Cretaceous

P. (P.) basiprocta SMITH & WRIGHT, 2003: 512; text-fig. 209; pl. 161: fig. 9. [Bulford, Wiltshire, England] (Micraster coranguinum Zone, Early Santonian, Late Cretaceous) <HT: BMNH EE6762>.

P. (P.) transiens SMITH & WRIGHT, 2003: 509-510; text-figs 202, 203, 207; pl. 162: figs 1-8. [Cuxton (Kent), Dover (Kent), Sussex, Guildford (Surrey), and Louth (Lincolnshire), England] (*S. plana* Zone, Late Turonian to basal Coniacian, Late Cretaceous) <HT: BMNH E16896 (from Cuxton, Kent); PT: BMNH 4740a-b, 30143, 75922, E9788-89, E41792>.

Genus *Plesiohemipneustes* SMITH & WRIGHT, 2003

Plesiohemipneustes SMITH & WRIGHT, 2003: 483. Type-species: *Holaster revestensis* LAMBERT in LAMBERT & THIÉRY, 1924 [England and France] (Cenomanian, Cretaceous) {assigned to the family Hemipneustidae by SMITH & WRIGHT (2003)}.

Genus *Protocardiaster* SMITH & WRIGHT, 2003

Protocardiaster SMITH & WRIGHT, 2003: 476. Type-species: *Spatangus truncatus* GOLDFUSS, 1829. [Western Europe] (Cenomanian to Early Coniacian, Late Cretaceous) {other species included: *Cardiaster cotteauanus* D'ORBIGNY, 1855}.

Genus *Pseudanachys* POMEL, 1883

Campanian

P. tumida TANAKA, 1984b: 192-193; text-fig. 3; pl. 1, figs 4a-c; pl. 2, figs 1a-c. [Chinomigawa Fm., Upper Yezo Grp., Higashimachi, Urakawa-cho, Urakawa-gun, Hokkaido, Japan] (Middle Campanian) <HT: YCM U611-1; PT: YCM U611-2>.

Cenomanian

P. alpina DE VILLOUTREYS & BIDAR, 1973: 1977-1979; fig. 1; pl. 1, figs 1-2. [Ravin de Font de la Poule, la Mure, Alpes-de-Haute-Provence, France] (Middle Cenomanian, base of *Rotomagense* zone) <HT: Muséum d'Histoire Naturelle de Nice, coll. Thomel no. 17182>.

Genus *Pseudholaster* POMEL, 1883

Late Cretaceous

P. mangyschlakensis SHMIDT in SHMIDT, GORBATOV & ZHELEZKO, 1979: 106; figs 2: 3a-b. [Besakty, Mangyshlak, Kazhakstan] (Early Cenomanian) <ST: CGM 3/II569, 4/II569> {as *Pseudoholaster* [sic!] *mangyschlakensis*}.

Mid-Cretaceous

P. baconicus SZÖRÉNYI, 1955: 97-100, 232-234; figs 38-40; pl. 13, figs 14-15; pl. 14, figs 1-16. [marne glauconieuse et calcaire gris lamellé, Bakonyháza (carrière de la vallée Gaja), Bakonyháza (pente Est du mont Judenberg), Bakonyháza (carrière située à l'Est du ruisseau Gaja), Bakonyháza (carrière près de la route de Felsőpere), Szápármalom à l'Est de Csörpuszta, Pénzeskút-Körisgyörpuszta, Jásd, Olaszfalu-Villóhegy, Olaszfalu-Eperkeshegy, and carrière à l'est de la nouvelle route Csörpuszta à Inotapuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/415 (from Bakonyháza, carrière de la vallée Gaja)>.

P. blackdownensis SMITH & WRIGHT, 2003: 469; pl. 148: figs 1-5. [Blackdown Greensand, Blackdown, Devon, England] (*P. inflatum* Zone, Late Albian, Early Cretaceous) <HT: BMNH E1334; PT: BMNH EE8218>.

P. caseyi SMITH & WRIGHT, 2003: 453-455; text-figs 177, 178; pl. 140: figs 4-7. [Red Bed, Sandling Junction, Hythe, Kent, England] (*H. jacobi* Zone, Late Albian, Early Cretaceous) <HT: BGS GSM Zm 32>.

P. depressus SMITH & WRIGHT, 2003: 468-469; text-fig. 185; pl. 148: figs 6-7. [Blackdown Greensand, Blackdown, Devon, England] (*P. inflatum* Zone, Late Albian, Early Cretaceous) <HT: BGS GSM 110500>.

Genus *Rispolia* LAMBERT, 1917

Cretaceous

R. hungarica SZÖRÉNYI, 1955: 100-101, 235-236; pl. 15, figs 1-4, 6. [marne glauconieuse, Bakonyhána (carrière de la vallée Gaja), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/428>.

Genus *Stegaster* POMEL, 1883

SMITH, GALLEMÍ, JEFFERY, ERNST & WARD (1999: 110) place this genus within the family Stegasteridae LAMBERT, 1913.

Paleocene

S. palaeocenicus SMITH & GALLEMÍ in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 113-114; figs 25a-b; pl. 5, figs 4-6. [Aristregui & Larumbe, Navarra, Spain] (Late Danian) <MGB 37351>.

Genus *Seunaster* LAMBERT in BLAYAC, 1912

In the *Treatise*, WAGNER & DURHAM (1966: U533) considered *Seunaster* a subjective synonym of *Stegaster* POMEL, 1883.

Coniacian

S. (Seunaster) schmidti EGOYAN, 1955: 165-167; pl. 3, figs 1a-г. [Vedi River, between Aziz-Kend and Dashnov, Vedin Region, Armenia] (Coniacian) <HT: repository not given>.

Senonian

S. bulgaricus TZANKOV, 1984: 101; pl. 45, figs 2, 2a-c. [Kondel, Dragoman, de Sofia, Bulgaria] (Lower Senonian) <HT: USC CR₂ 1302>.

Genus *Tholaster* SEUNES, 1890

Campanian

T. carvalhoi GREYLING & COOPER, 1995: 68-71; figs 6-9. [Praia Egito, near Quimbala, 70 km N of Lobito, Angola] (Marroti zone, Middle Campanian) <HT: SAfM-PCA2240>.

Cretaceous

T. fourmarieri LAMBERT, 1931: M8 ; pl. 1, figs 5-6. [Marnes de Bartine near Guv-Tepessi, région d'Héraclée, Anatolia, Turkey] (Cretaceous) <HT: not given>.

Subfamily Cardiotaxinae SMITH & JEFFERY, 2000

Cardiotaxinae SMITH & JEFFERY, 2000: 287. Type-genus: *Cardiotaxis* LAMBERT, 1917. {other genera included: *Infulaster* WRIGHT & WRIGHT, 1948; *Hagenowia* DUNCAN, 1889}.

Subfamily Pseudholasterinae SMITH & JEFFERY, 2000

Pseudholasterinae SMITH & JEFFERY, 2000: 289. Type-genus: *Pseudholaster* POMEL, 1883. {other genera included: *Hemipneustes* L. AGASSIZ, 1836; *Giraliaster* FOSTER & PHILIP, 1978}.

Family Corystidae FOSTER & PHILIP, 1978

Corystidae FOSTER & PHILIP, 1978: 792-793. Type-genus: *Corystus* POMEL, 1883. [Australia, New Zealand] (Cenozoic) {emended to Corystusidae to remove homonymy with a crustacean family (BOYKO, 2008; ICZN Opinion 2238)}.

Genus *Cardabia* FOSTER & PHILIP, 1978

Cardabia FOSTER & PHILIP, 1978: 798. Type-species: *Cardabia bullarensis* FOSTER & PHILIP, 1978. [Australia] (Paleocene).

Paleocene

C. bullarensis FOSTER & PHILIP, 1978: 798-799; pl. 90: figs 3-4; pl. 91: figs 1-3. [Cardabia Grp., Giralia Anticline, Carnarvon Basin, North West Division, Western Australia] (Middle or Late Paleocene) <HT: WAM 73.361; PT: WAM 73.365-66>.

Genus *Huttonechinus* FOSTER & PHILIP, 1978

Huttonechinus FOSTER & PHILIP, 1978: 799. Type-species: *Macropneustes spatangiformis* HUTTON, 1873. [Australia] (Oligocene).

Late Cretaceous

H. antarctica NÉRAUDEAU, CRAME & KOOSER, 2000: 462-463; fig. 4.1-2, 9. [Lachman Crags Mb., Santa Marta Fm., Marambio Grp., northern James Ross Island, Antarctica] (Late Santonian-Early-Campanian) <HT: BAS D.8615.55>.

Family Plexechinidae MOOI & DAVID, 1996

Plexechinidae MOOI & DAVID, 1996: 946-947. Type-genus: *Plexechinus* A. AGASSIZ, 1896. [circum-Antarctic and Southern Pacific] (Recent).

Genus *Plexechinus* A. AGASSIZ, 1896

Recent

P. sulcatus DAVID & MOOI, 2000: 167-171; figs 1a-c, 2a-d, 3, 4a-g. [R/V “Marion Dufresne”, MD03-station 17, northwest of Kerguelen Islands; 47°24,9' S, 66°04' E] (Recent) <HT: MNHN EcEs 9343; PT: MNHN EcEs 9344>.

Family Urechinidae DUNCAN, 1889

Genus *Antrechinus* MOOI & DAVID, 1996

Antrechinus MOOI & DAVID, 1996: 946. Type-species: *Urechinus mortenseni* DAVID & MOOI, 1990. [circum-Antarctic and West Pacific] (Recent) {other species included: *Urechinus drygalskii* MORTENSEN, 1905 and *Plexechinus nordenskjoldi* MORTENSEN, 1905}.

Genus *Pilematechinus* A. Agassiz, 1904

Recent

P. belyaevi MIRONOV, 1975: 208-210; figs 1a-д, 2.4, 2.7-2.9. [R/V Akademik Kurchatov 14th cruise Stat. 1259 (19°04' N, 80°30' W, depth 5800-6500 m), 1267 (19°38' N, 76°37' W, depth 6740-6780 m), Caribbean Sea] (Recent) <HT: IOANSSR XV-69-10 (from R/V Akademik Kurchatov 14th cruise Stat. 1259)>.

Genus *Urechinus* A. AGASSIZ, 1879

Recent

- U. antipodeanus* MCKNIGHT, 1974: 33-35; figs 4a-b. [NZOI Stat. G701 (46°20' S, 171°30' E, 1400 m depth), G704 (46°17' S, 172°37' E, 1600 m depth), G705 (46°04' S, 172°28.5' E, 1500 m depth), G706 (45°49' S, 172°30' E, 1550 m depth), New Zealand] (Recent) <HT: NZOI 184 (Stn G705); PT: NZOI P242 (Stn G705)>.
- U. aoteanus* MCKNIGHT, 1974: 35-37; figs 5a-b. [NZOI Stat. E903a (37°33' S, 172°05' E, 964-962 m depth), New Zealand] (Recent) <HT: NZOI 185; PT: NZOI P243>.
- U. drygalskii perfidus* MIRONOV, 1976: 146-147; figs 1б-ж; pl. 2, figs 1-2; pl. 4, figs 2, 4, 5, 7-9; pl. 5, figs 1-3. [R/V Vitayaz Stat. 3359 (51°30' N, 172°04' E, depth 5020 m), 5620 (44°48' N, 156°33' E, depth 5005-5045 m), 6088 (53°58'5 N, 157°36' W, depth 5740 m), 6142 (53°13'5 N, 163°43' W, depth 4990-5000 m), Northern Pacific Ocean, Bering Sea] (Recent) <HT: IOANSSR XV-69-11 (from R/V Vitayaz Stat. 5620)>.
- U. mortenseni* DAVID & MOOI, 1990: 76-81; figs 2-16. [“Eltanin” station 410, southwest of Elephant Island, South Shetland Islands, Antarctic Ocean] (Recent) <HT: USNM E40039; PT: USNM E11016> {placed into the genus *Antrechinus* MOOI & DAVID, 1996 by MOOI & DAVID, 1996: 946}.
- U. parvus* MIRONOV, 1978a: 219-220; figs 4б-в; pl. 3, figs 1, 3, 6-7. [16th Cruise R/V “Dimitri Mendeleev” Stat. CT 1388, off South-west Australia; 2320-2360 m depth] (Recent) <HT: IOANSSR XV-69-15>.
- U. planus* MIRONOV, 1978a: 216-219; tab. 4; figs 2ж,3, figs 3a-в; pl. 1, figs 7-8; pl. 2, figs 1-2, 5-6. [16th Cruise R/V “Dimitri Mendeleev” Stat. CT 1347, South of Tasmania; 1800-1820 m depth] (Recent) <HT: IOANSSR XV-69-14>.

Family Carnarechininae MIRONOV, 1993

Carnarechininae MIRONOV, 1993: 221. Type-genus: *Carnarechinus* MIRONOV, 1978a. (Recent).

Genus *Carnarechinus* MIRONOV, 1978a

Carnarechinus MIRONOV, 1978a: 209-211. Type-species: *Cystechinus clypeatus* A. AGASSIZ, 1879. [Southern Pacific Ocean] (Recent).

Family Pourtalesiidae A. AGASSIZ, 1881

Genus *Ceratophysa* POMEL, 1883

Recent

C. ceratopyga valvaecristata MIRONOV, 1976: 149-150; figs 26-3; pl. 2, fig. 3; pl. 5, fig. 10. [R/V Vitayaz Stat. 3162 (43°15' N, 157°48' E, depth 5502 m), 3166 ? (44°43' N, 153°49' E, depth 5057 m), 3575 (38°02' N, 146°33' E, depth 5495 m), 4120 (53°37' N, 159°40' W, depth 6296-6328 m), 4213 ? (34°54' N, 123°56' W, depth 4200-4231 m), 5605 (46°10' N, 153°07' E, depth 4915-4985 m), 5624 (45°26' N, 154°12' E, depth 5200 m), 6088 (53°58' N, 157°36' W, depth 5740 m), 6142 (53°13' N, 163°43' W, depth 4990-5000 m), 6143 (51°40' N, 163°00' W, depth 4860 m), North Pacific Ocean] (Recent) <HT: IOANSSR XV-69-12 (from R/V Vitayaz Stat. 3575)>.

Genus *Echinocrepis* A. AGASSIZ, 1879

Recent

E. rostrata MIRONOV, 1973: 240-243; pl. 1, figs 1-6. [R/V Vitayaz Stat. 3359 (51°30' N, 172°04' E, depth 5020 m), 4147 (49°35' N, 133°57' W, depth 3470 m), 4158 (46°57' N, 143°59' W, depth 4661-4665 m), 4213 (34°54' N, 123°56' W, depth 4200-4231 m), 4265 (24°58' N, 113°25' W, depth 3315-3340 m), 5605 (46°10' N, 153°07' E, depth 4915-4985 m), 5634 (44°17' N, 149°33' E, depth 4690-4720 m), 6107 (57°38' N, 143°12' W, depth 3800 m), 6109 (56°13' N, 139°43' W, depth 3460 m), North Pacific Ocean] (Recent) <HT: IOANSSR no specimen no. given (from R/V Vitayaz Stat. 4213)>.

Genus *Pourtalesia* A. AGASSIZ, 1869

Recent

P. heptneri MIRONOV, 1978b: 721-726; figs 1: 1-2, 4; figs 2-4. [“Витязь”-cruise (“Vityaz”-cruise) Stat. 6785 (5°34' S, 131°08' E, depth 7130 m), 7271 (5°37' S, 131°07.5' E, depth 7340-7335 m), Banda Trench, West Pacific] (Recent) <HT: IOANSSR XV-69-16>.

P. jeffreysi gibbosa MIRONOV, 1995a: 70-74; figs 1: 1-2, 2: 1-2. [Expedition of the Plavoochego Institute of Marine Research 1923-1924 Stat. 97 (79°50' N, 43°30' E, depth 334 m), 186 (77°44' N, 38°35' E, depth 230 m), 189 (78°10' N, 31°23' E, depth 225 m), “Седов”-cruise (“Sedob”-cruise) Stat. 25 (78°33' N, 63°10' E, depth 363 m), “Ломоносов”-cruise (“Lomonosob”-cruise) Stat. 7 (78°54' N, 70°14' E, depth 500 m), 12 (78°03' N, 79°47' E, depth 426 m), “Садко”-cruise 1935 (“Sadko”-cruise) Stat. 33 (80°55' N, 72°29' E, depth 520 m), 35 (81°11' N, 66°53' E, depth 520 m), 39 (80°43' N, 68°08' E, depth 542 m), “Садко”-cruise 1936 (“Sadko”-cruise) Stat. 10

- (80°50'4" N, 71°47' E), 11 (80°23'8" N, 70°09'5" E, depth 582 m), “Литке”-cruise (“Litke”-cruise) Stat. 13 (82°11' N, 60°37' E, depth 923 m), 29 (81°18' N, 9°36' E, depth 1301 m), “Лена”-cruise (“Sena”-cruise) Stat. 2 (81°32' N, 5°15' E, depth 760 m), “Севастополь”-cruise (“Sevastopol”-cruise) Stat. 1380 (67°54' N, 14°18' E, depth 1270 m), Arctic Ocean] (Recent) <HT: ZI 1/22940 (from “Lomonosob”-cruise Stat. 12)>.
- P. jeffreysi lata* MIRONOV, 1995a: 74-76; figs 2: 3, 4. [“Литке”-cruise (“Litkye”-cruise) Stat. 37 (82°39' N, 33°30' E, depth 2899 m), “Объ”-cruise (“Ob”-cruise) Stat. 45 (79°47' N, 1°41' W, depth 2800 m), “Polarstern”-cruise Stat. 32 (78°43' N, 132°33' E, depth 3011-3028 m), 54 (79°12' N, 119°56' E, depth 3076-3081 m), Arctic Ocean] (Recent) <HT: ZI 22941 (from “Polarstern”-cruise Stat. 54)>.
- P. thomsoni* MIRONOV, 1976: 152; figs 3г-д; pl. 3, figs 1-7; pl. 5, figs 3-6, 8, 11, 13. [R/V Vitayaz Stat. 4213 (34°54' N, 123°56' W, depth 4200-4231 m), 4265 (24°58' N, 113°25' W, depth 3315-3340 m), 6117 (56°12' N, 139°12' W, depth 3350-3370 m), North Pacific Ocean, West American and Alaskan Coast] (Recent) <HT: IOANSSR XV-69-13 (from R/V Vitayaz Stat. 4265)>.
- P. vinogradovae* MIRONOV, 1995b: 67-68; figs 6: 1A-D. [“Академик Курчатов”-cruise (“Akademik Kurchatov”-cruise) Stat. 896 (56°52' S, 24°53' W, depth 5530-5651 m), 909 (60°13' S, 44°00' W, depth 5450-5480 m), 914 (56°21' S, 50°48' W, depth 5670-6070 m), 916 (56°29' S, 50°51' W, depth 4664-5631 m), “Дмитрий Менделеев”-cruise (“Dmitrig Mendelejev”-cruise) Stat. 1290 (54°33' S, 159°24' E, depth 5450-5410 m), 1292 (54°48.7' S, 159°10.4' E, depth 5400 m), 1306 (59°12' S, 158°32' E, depth 6210-6100 m), ? 4086 (60°50' S, 41°07' W, depth 6120-6290 m, tentative record based on fragments), ? 4090 (60°52' S, 40°56' W, depth 6145-5550 m, tentative record based on fragments), Antarctic, Sout-West Atlantic Ocean] (Recent) <HT: IOANSSR, no specimen no. given (from “Akademik Kurchatov”-cruise Stat. 914)>.

Genus *Echinisigra* MORTENSEN, 1907

Subgenus *Echinisigra* MORTENSEN, 1907

Recent

- E. amphora* MIRONOV, 1974a: 245-249; figs 1a, б, г-е, и, л, м, 2a-г, ж, з, к; pl. 1, figs в-е; pl. 2, figs а-з. [see respective ssp., North west Pacific] (Recent) <HT: IOANSSR XV-60-3 (from R/V Vityaz Stat. 3114)>.
- E. amphora amphora* MIRONOV, 1974a: 246-247; figs 1a, г, д, м, 2г, к; pl. 1, figs г-е. [R/V Vitayaz Stat. 2074 (42°32' N, 150°41' E, depth 5140 m), 2119 (46°08' N, 155°16' E, depth 5070-5090 m), 3114 (48°51' N, 160°01' E, depth 5511, 5670-5680 m), 3156 (39°57' N, 165°08' E, depth 5535 m), 3166 (44°43' N, 153°49' E, depth 5057 m),

3225 (37°51' N, 144°13' E, depth 5290-5390 m), 3363 (48°15' N, 169°39' E, depth 6272-6282 m), 3575 (38°02' N, 146°33' E, depth 5495 m), 3886 (32°11' N, 143°10' E, depth 5680-5690 m), 5621 (45°18' N, 156°00' E, depth 5035-5210 m), 5624 (45°26' N, 154°12' E, depth 5200 m), North-west Pacific Ocean] (Recent) <HT: IOANSSR XV-60-3 (from R/V Vityaz Stat. 3114)>.

E. amphora antarctica MIRONOV, 1974a: 248; figs 2ж, з. [R/V Ob Stat. 57 (64°03' S, 161°59' E, depth 2937 m), Pacific Antarctic Ridge, Ross Sea] (Recent) <HT: IOANSSR XV-69-6 (from R/V Ob Stat. 57)> {MIRONOV (1997: 185) raised this ssp. to species level}.

E. amphora fabrefacta MIRONOV, 1974a: 247-248; figs 2а-в; pl. 1, fig. в; pl. 2, figs а, б. [R/V Vityaz Stat. 6097 (57°00' N, 148°18' W, depth 4740 m), 6106 (58°15' N, 142°36' W, depth 3610 m), 6107 (57°38' N, 143°12' W, depth 3800 m), 6143 (51°40' N, 163°00' W, depth 4860 m), Northern Pacific, Alaskan Coast] (Recent) <HT: IOANSSR XV-69-4 (from R/V Vityaz Stat. 6107)> {MIRONOV (1997: 183) raised this ssp. to species level}.

E. amphora indica MIRONOV, 1974a: 248; figs 1е, л; pl. 2, figs в-е. [R/V Vityaz Stat. 4535 (9°58' S, 107°56' E, depth 6820-6850 m), 5168 (8°42' S, 105°31' E, depth 6433 m), Indian Ocean] (Recent) <HT: IOANSSR XV-69-5 (from R/V Vityaz Stat. 4535)>.

E. amphora valvaedentata MIRONOV, 1974a: 249; figs 1б, и; pl. 2, figs ж, з. [R/V Akademik Kurchatov Stat. 896 (56°52' S, 24°59' W, depth 5530-5651 m), 909 (60°13' S, 44°00' W, depth 5450-5480 m), 916 (56°29' S, 50°51' W, depth 4664-5631 m), Falkland Islands, South Atlantic] (Recent) <HT: IOANSSR XV-69-7 (from R/V Akademik Kurchatov Stat. 916)> {MIRONOV (1997: 183) raised this ssp. to species level}.

E. mortenseni MIRONOV, 1974b: 1804-1805; pl. 1, figs 5-6, 13; pl. 2, fig. з. [“Витязь”-cruise (Vityaz-cruise), Stat. 3364 (48°21.2' N, 169°54.1' E; depth 2915-3015 m), Northern Pacific] (Recent) <HT: IOANSSR XV-69-8>.

E. (E.) phiale partita MIRONOV, 1997: 176-177, figs 3A-D, H. [R/V Dmitrig Myendelev Stat. 1276 (48°5' S, 171°42' E, depth 1100-1200 m), 1281 (53°23' S, 167°07.8' E, depth 1026 m), 1347 (44°06.6' S, 145°56' E, depth 1800 m), Subantarctic] (Recent) <HT: IOANSSR XV-69-30 (from R/V Dmitrig Myendelev Stat. 1276)>.

E. porrecta MIRONOV, 1974b: 1805; pl. 1, figs 3-4, 7, 11-12; pl. 2, fig. и. [“Дмитрий Менделеев”-cruise (“Dmitrig Mendelejev”-cruise), Stat. 525; “Витязь”-cruise (“Vityaz”-cruise), Stat. 4868 (11°16.9' N, 70°59' E, depth 2623 m), 5153; “Академик Курчатов”-cruise (“Akademik Kurchatov”-cruise), Stat. 22₁ (12°21.9' N, 61°12.4' E, depth 3614-3800 m), 22₂ (12°22.1' N, 61°15.9' E, depth 2280-2920 m), 28-29 (5°25.9' S, 68°34.8' E, depth 4800-5100 m); Cocos-Keeling Basin, Chagos Archipelago; depth: 2623-5200 m] (Recent) <HT: IOANSSR XV-69-9 (from “Akademik Kurchatov”-cruise, Stat. 28-29)>.

E. (E.) vityazi MIRONOV, 1997: 178-180, figs 6A-C, E, K. [R/V Vityaz Stat. 4954 (9°34.9' N, 90°54.4' E, depth 3485 m), 7325 (1°51.5' S, 144°40.8' E, depth 2550-2580 m), R/V Dmitrig Myendeleev Stat. 1235 (11°30.5' S, 152°11.7' E, depth 3070-3080 m), 1253 (29°28.5' S, 164°55.1' E, depth 3400 m), 1254 (30°00.0' S, 169°02.6' E, depth 2970-3020 m, Indo-Pacific Ocean] (Recent) <HT: IOANSSR XV-69-31 (from R/V Vityaz Stat. 4954)>.

Subgenus *Echinogutta* MIRONOV, 1997

Echinogutta MIRONOV, 1997: 180. Type-species *Echinosisgra amphora amphora* MIRONOV, 1974a. (Recent) {*Echinocutta* [sic!] in the heading of the paragraph, but *Echinogutta* elsewhere in the text; other species included: *Echinosisgra amphora indica* MIRONOV, 1974a, *Echinosisgra fabrefacta* MIRONOV, 1974a, *Echinosisgra valvaedentata* MIRONOV, 1974a, *Echinosisgra antarctica* MIRONOV, 1974a}.

Genus *Rictocystis* MIRONOV, 1996

Rictocystis MIRONOV, 1996: 1113-1114. Type-species: *Rictocystis jensenae* MIRONOV, 1996. [Indian Ocean and Western Pacific] (Recent):

Recent

R. jensenae MIRONOV, 1996: 1114-1117; figs 3-5. [“Galathea” cruise Stat. 474 (9°49' S, 114°13' E, depth 3920-3940 m), “Академик Курчатов”-cruise (“Академик Курчатов”-cruise) Stat. 25 (4°32.5' S, 63°19.6' E, depth 4430-4440 m), “Витязь”-cruise (“Vityaz”-cruise) Stat. 6775 (5°54.7' N, 128°30' E, depth 4920 m), “Дмитрий Менделеев”-cruise (“Dmitrig Mendeleyev”-cruise) Stat. 1235 (11°30.5' S, 152°11.7' E, depth 3070-3080 m); Indian Ocean and Western Pacific] (Recent) <HT: ZMUC ECH-1 (from “Galathea” cruise Stat. 474)>.

Genus *Solenocystis* MIRONOV, 2008

Solenocystis MIRONOV, 2008: 13-16. Type-species: *Solenocystis imitans* MIRONOV, 2008. [North Atlantic] (Recent).

Recent

S. imitans MIRONOV, 2008: 16-18; figs 2A-C, 7A-D, 8A-I, 9C. [RV G.O. Sars, MAR-ECO expedition, St. 72/386 (27 July 2004, 53°16' N, 35°31' W, 2555-2517 m), Northern Atlantic Ocean] (Recent) <HT: MZUB MAR-ECO 008141> {based upon a single specimen}.

Family Somaliasteridae WAGNER & DURHAM, 1966

A cladistic analysis of this group carried out by JEFFERY (1999:1038-1039) shows that this family should be placed into the Spatangoida instead of Holasteroida.

Genus *Iranianster* COTTEAU & GAUTHIER, 1895

Upper Cretaceous

- I. affinidouvillei* KIER, 1972a: 74-77; figs 37D, 39D, 44; pl. 52, figs 4-7; pl. 53, figs 1-6. [Lower Aruma Fm., locality KK11, Saudi Arabia] (Campanian) <HT: USNM 170467; PT: USNM 170466, 170468-9, 170504, 170507>.
- I. affinimorgani* KIER, 1972a: 77-81; figs 38A, 39E, 43, 44; pl. 52, figs 1-3. [Lower Aruma Fm., locality KK11, Saudi Arabia] (Campanian) <HT: USNM 170462; PT: USNM 170463-170465, 170505>.
- I. bowersi* KIER, 1972a: 81-84; figs 38B, 40; pl. 54, figs 1-6. [Lower Aruma Fm., locality S-289, S-290, S-291, S-1234 and S-1419, Saudi Arabia] (Campanian) <HT: USNM 170470; PT: USNM 170471, 170506>.
- I. omanensis* JEFFERY, 1999: 1032-1034; text-fig. 4a-c; pl. 1, figs 6-8. [Simsima Fm., Jebel Lahjan, Yanqul, Oman] (Maastrichtian) <HT: BMNH E82606; PT: BMNH E82603-E82605, E82607>.

Family Uncertain

Genus *Coraster* COTTEAU, 1886

JEFFERY in SMITH & JEFFERY (2000: 354) placed the genus into the family Corasteridae LAMBERT in LAMBERT & THIÉRY, 1924 in the order Spatangoida CLAUS, 1876.

Paleocene

- C. urmaensis* MOSKVIN, 1982: 106; figs 1j-l; pl. 10, figs 4a-e, 5a-e. [Urma, Dagestan] (Upper Paleocene) <HT: PIN 3939/4> {JEFFERY in SMITH & JEFFERY (2000: 355-356) placed this species into the synonymy of *C. vilanovae* COTTEAU, 1886}.

Genus *Cottreaucorys* LAMBERT, 1920

Paleocene

C. kollmanni KROH, 2004: 313-317; fig. 2: 1a-d; pl. 1, figs 1a-d. [Bruderndorf Formation, Haidhof, near Ernstbrunn, Lower Austria, Austria] (Late Danian, Paleocene) <HT: NHMW 2004z0075/0004> {based on a single specimen}.

Genus *Nordenskoeldaster* LAMBERT, 1910

Campanian

N. ? australis NÉRAUDEAU, CRAME & KOOSER, 2000: 460-462; fig. 4.7-8. [Rabot Point Mb., Santa Marta Fm., Marambio Grp., Rabot Point, James Ross Island, Antarctica] (Lower to Mid-Campanian) <HT: BAS DJ.662.6>.

Superfamily Orthosternata SMITH & JEFFERY, 2000

Orthosternata SMITH & JEFFERY, 2000: 264. {taxa included: *Corystidae* FOSTER & PHILIP, 1978; *Basseaster* LAMBERT, 1936; *Galeaster* SEUNES, 1889; and *Garumnaster* LAMBERT, 1936}.

Order Collyritoida VADET, 2007

Collyritoida VADET, 2007: 217. Type-family: Collyritidae D'ORBIGNY, 1853. (Jurassic).

Order Spatangoida CLAUS, 1876

Suborder Toxasterina FISCHER, 1966

Family Toxasteridae LAMBERT, 1920

Genus *Douvillaster* LAMBERT, 1917

Cretaceous

D. subtrigonalis SZÖRÉNYI, 1955: 118-119, 253-254; fig. 44; pl. 21, figs 1-5. [marne glauconieuse, Bakonyháza (carrière de la vallée Gaja, pente de l'Est du mon Judenberg, carrières situées à l'Est du ruisseau Gaja et à côté de la route de Felsöper), and Bakonyháza-Csigahegy, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/531 (from Bakonyháza)>.

Genus *Epiaster* D'ORBIGNY, 1854

This genus was considered as junior synonym of *Heteraster* D'ORBIGNY, 1853 in the Treatise (FISCHER, 1966: U553)

Cenomanian

E. brevipetalus SMITH & WRIGHT, 2008: 633-635; text-figs 249, 275, 276A, B; pl. 207: fig. 2; pl. 209: figs 1-2. [Grey Chalk, Dover, Kent, and Isle of Wight, UK] (*S. dispar* Zone, Late Albian and *C. subglobosus* Zone, Middle Cenomanian) <HT: BMNH E1170; PT: BGS GSM 119609, BMNH E1092>.

E. romani COLLIGNON, 1983: 270; pl. 6, fig. 8. [coupe de Gavbasht, Kazhdumi, Iran] (Cenomanian) <HT: CFP 8005-175>.

Aptian

E. zonarius TANAKA in TANAKA & OBATA, 1982: 132-135; text-figs 8-9; pl. 2, figs 5; pl. 3, figs 1a-e, 2a-b, 3. [Hiraiga Fm., Hideshima, Miyako City, Iwate Prefecture, Honshu, Japan] (Upper Aptian) <HT: GSJ F6138; PT: GSJ F6139>.

Albian

E. miyakoanus TANAKA in TANAKA & OBATA, 1982: 129-132; text-figs 6-7; pl. 2, figs 1a-e, 2a-c, 3a-e, 4a-b. [Aketo Fm., Raga, Tanohatamura, Shimohei-gun, Iwate Prefecture, Honshu, Japan] (Lower Albian – middle (?) Upper Aptian) <HT: NSM PA12129; PT: GSJ F6142, NSM PA12130a>.

Mid-Cretaceous, undifferentiated

E. angulosus SZÖRÉNYI, 1955: 106-107, 241-242; pl. 16, figs 2, 5-6, 9, 14. [marne à *Turrilites*, Olaszfalu-Eperkeshegy, Jásd-Doboshegy, Bakonynána (carrière de la vallée Gaja), and Bakonynána-Heuberg, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/450 (from Olaszfalu, pente Est de l'Eperkeshegy)>.

E. baconicus SZÖRÉNYI, 1955: 112-113, 248; pl. 20, figs 15, 19. [marne à *Turrilites*, Péntzeskút, and Lókút, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/484 (from Péntzeskút)>.

E. hemiasteriformis SZÖRÉNYI, 1955: 111-112, 247; pl. 19, figs 8-10; pl. 20, figs 1-2. [marne à *Turrilites* et marne glauconieuse, Olaszfalu-Eperkeshegy, Bakonynána-Judenberg, carrières à l'Est du ruisseau Gaja, and Jásd-Doboshegy, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/457 (from Olaszfalu-Eperkeshegy)>.

E. hungaricus SZÖRÉNYI, 1955: 107-108, 242-243; pl. 16, fig. 12; pl. 17, figs 1-4. [marne à *Turrilites*, Olaszfalu (à droite de la route d'Eplény), Jásd (route de Csösz), Jásd-Doboshegy, Jásd-Szentkút, and Bakonynána-Csigahegy, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/459 (from Olaszfalu, à droite de la route d'Eplény)>.

- E. hungaricus carinatus* SZÖRÉNYI, 1955: 108-109, 244; pl. 18, figs 1-2, 4-6. [marne à *Turrilites*, Bakonynána-Judenberg (carrière située à l'Est du ruisseau Gaja), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/473>.
- E. hungaricus intermedius* SZÖRÉNYI, 1955: 108, 243-244; pl. 17, figs 5-9. [marne à *Turrilites*, Jásd (route de Csösz), Jásd-Doboshegy, and Bakonynána, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/469 (from Jásd-Doboshegy)>.
- E. hungaricus rotundatus* SZÖRÉNYI, 1955: 109, 244-245; pl. 18, figs 3, 7-9. [marne à *Turrilites*, Jásd (route de Csösz), Jásd-Doboshegy, and Várpalota-Csöszpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/462 (from Jásd, route de Csösz)>.
- E. pseudodistinctus* SZÖRÉNYI, 1955: 110, 245-246; pl. 19, figs 1-4. [marne à *Turrilites*, Olaszfalu (à côté de la route de Veszprém), Olaszfalu-Eperkeshegy, Jásd (route de Csösz), Jásd-Doboshegy, and Bakonynána, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/479 (from Olaszfalu, à côté de la route de Veszprém)>.
- E. pseudodistinctus oblongus* SZÖRÉNYI, 1955: 111, 246; pl. 19, figs 5-6. [marne à *Turrilites* et marne glauconieuse, Olaszfalu-Eperkeshegy, Jásd (route de Csösz), Jásd-Doboshegy, and Bakonynána (ravine du ruisseau Gaja), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/479 (from Olaszfalu-Eperkeshegy)>.
- E. pseudodistinctus rotundiformis* SZÖRÉNYI, 1955: 111, 246-247; pl. 19, figs 7, 11-12. [marne à *Turrilites*, Jásd (ravine au bord du chemin creux menant vers l'usine d'électricité), Olaszfalu-Eperkeshegy, and Olaszfalu (côté Est du mont Eperkeshegy, à côté de la route), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/481 (from Jásd)>.

Genus *Heteraster* D'ORBIGNY, 1853

Mid-Cretaceous

- H. zircensis* SZÖRÉNYI, 1955: 113-118, 249-252; figs 41-43; pl. 20, figs 3-14, 16-18, 20; pl. 21, figs 6-10. [groupe de marnes argileuses, Zirc-Tündérmajor, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/485>.

Barremian

- H. bungoensis* TANAKA & NODA in TANAKA et al., 1984: 448-450; text-figs 2-3; pl. 82, figs 3-6; pl. 83, fig. 4. [Haidateyama Group, Honjomura, Minami-mabe-gun, Oita Prefecture, Kyushu, Japan] (Lower Barremian) <HT: GSJ F6013; PT: GSJ F6014, F6036, F6033A, B, F6039A, B>.
- H. debensis* DEVIRIES, 1973: 69-71; pl. 1: figs 26-28; pl. 3: figs 1-4. [Kef ed Deb, map sheet Ain Regada, Algeria] (Barremian, Early Cretaceous) <no types defined, material in the coll'n of A. Devries>.

- H. heckeri* MELIKOV, 1989: 157-160; figs a-b; pl. 1, figs 1-6. [Azerbaijan] (Barremian) <HT: AZIOC 775/15φ/4 >.
- H. magnus* PORETSKAYA, 1961: 104; pl. 20, figs 1a-д, 3, 4a, б, 5. (Barremian) <unknown> {cited and illustrated in MELIKOV (in ALI-ZADE, 1988: 187, pl. 1, figs 6a-b), but reference missing from list}.
- H. renngarteni* PORETSKAYA, 1961: 171; pl. 19, figs 4a-д, 5, 6a-b. (Barremian-Aptian) <unknown > {cited and illustrated in MELIKOV (in ALI-ZADE, 1988: 186, pl. 1, figs 3a-b), but reference missing from list}.

Subgenus *Enallaster* D'ORBIGNY, 1853

In the Treatise (FISCHER 1966: U553), *Enallaster* is considered a junior synonym of *Heteraster*.

Early Cretaceous

- H. (E.) hemiheterus* SMITH & WRIGHT, 2008: 575-576; text-figs 241B, D; pl. 184: figs 4-5. [Perna Beds, Redhill, Surrey, UK] (*P. fissicostatus* Zone, Early Aptian) <HT: BMNH E8585; PT: BMNH E8586 to E8589>.

Genus *Proisaster* AZIZ & BADVE, 2001

Proisaster AZIZ & BADVE, 2001: 57. Type-species: *Proisaster coramandeli* AZIZ & BADVE, 2001. [India] (Campanian).

Campanian

- P. coramandeli* AZIZ & BADVE, 2001: 57-58; pl. 5: figs 4-6. [Sillakudi Formation, Ariyalur Group, Mallur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Karapadites karapadense*-zone, Campanian) <HT: MACSG-2142>.

Genus *Toxaster* D'ORBIGNY, 1853

Campanian

- T. cauveriae* AZIZ & BADVE, 2001: 55; pl. 4: figs 4-6. [Sillakudi Formation, Ariyalur Group, Mallur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Karapadites karapadense*-zone, Campanian) <HT: MACSG-2128; PT: MACSG-2129-2131>.
- T. compressa* [*compresa*] AZIZ & BADVE, 2001: 55-57; pl. 4: figs 7-9. [Sillakudi Formation, Ariyalur Group, Mallur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Karapadites karapadense*-zone, Campanian) <HT: MACSG-2132; PT: MACSG-

2133-2137> {spelled “*compresa*” in heading, but “*compressa*” elsewhere in the paper (plate descriptions, differential diagnoses)}.

T. jugamis AZIZ & BADVE, 2001: 57; pl. 5: figs 1-3. [Sillakudi Formation, Ariyalur Group, Mallur, Tamil Nadu region, Trichinopoly Basin, Southern India] (*Karapadites karapadense*-zone, Campanian) <HT: MACSG-2138; PT: MACSG-2139-2141>.

Berriasian

T. bajarunasi LOBATSHEVA in PORETSKAYA & LOBATSHEVA, 1988: 172-173; pl. 36: figs 2a-b, 3a, 6, 4. [Sorbulak, West Karatau; Dzhaprakty mountain, East Karatau; Doshchan; Dzharmyui village; and Karakuduk, Karasyaz-Taspaskaya Anticline, Mangys-hlak, Kazhakstan] (Berriasian) <HT: CNIGR 27/11107>.

T. granosus var. *kouensis* LOBATSHEVA, 1961: 157; pl. 1: fig. 5. [Sorbulak, West Karatau and Dzhaprakty mountain, East Karatau, Karasyaz-Taspaskaya Anticline, Mangys-hlak, Kazhakstan] (Berriasian) <unknown> {elevated to species rank and re-described by PORETSKAYA & LOBATSHEVA (1988: 172)}.

T. priscus TANAKA, 1984b: 193-195; text-figs 4a-c, 5a-d; pl. 2, figs 2a-d, 3a-b. [Koyamada Fm., Soma Grp., Yamashita, Kashima-machi, Soma-gun, Fukushima Prefecture, Japan] (Berriasian) <HT: GSJ F6185A, B; PT: GSJ F6134>.

Hauterivian

T. crassisulcatus CLAVEL, 1989: 176-178; figs 11a-h; pl. 1, figs 2a-c. [Le Landeron; Cressier; Valangin; Menthnières, Hauteville (Ain), Censeau (Jura); Mont-de-Musièges and Salève (Haut-Savoie), France] (Base de la Zone à Radiatus, Hauterive) <HT: FSL 196109; PT: FSL 196114, 196115, 196116, 196124, 196125>.

T. remanei CLAVEL, 1989: 174-176; figs 10a-f; pl. 1, figs 1a-c. [Montlebon, Doubs; Le Landeron; Cressier; Valangin; Villier-le-Lac; Mont-de-Musièges and Mont-Clergeon, France] (Zone à Radiatus, Hauterive) <HT: FSL 196106; PT: FSL 196107, 196108, 196122, 196123>.

Genus *Pliotoxaster* FOURTAU, 1907

In the Treatise (FISCHER, 1966: U551) *Pliotoxaster* is considered a junior synonym of *Toxaster*.

Albian

P. angustisulcus SMITH & WRIGHT, 2008: 590-592; text-figs 249, 251, 252; pl. 190: fig. 5; pl. 191: figs 1-3. [Blackdown Greensand, Blackdown, Devon, UK] (*M. inflatum* Zone, Late Albian) <HT: BMNH E4941; PT: BMNH E2536, E4942, BGS GSM 110505, 110509>.

Suborder Hemiasterina FISCHER, 1966

Family Hemiasteridae CLARK, 1917

Genus *Hemiaster* AGASSIZ in AGASSIZ & DESOR, 1847

Late Cretaceous

- [*H. aktaschensis* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- H. amudariensis* SHMIDT & SIMAKOV, 1953: 65-66; figs 25a-r. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Senonian) <HT: VNIGRI 115/336> {referred to *Parahemiaster* [nomen nudum] by EGOROV (1972: 60)}.
- H. anisopetalus* SMITH & WRIGHT, 2008: 616-617; text-figs 266A, B, 267; pl. 203: fig. 2a-c. [Upper Chalk, Dover, Kent, UK] (*Micraster cortestudinarium* Zone, Coniacian, Late Cretaceous) <HT: BMNH E37783>.
- H. arachnius* DEVIRIES, 1973: 71-73; pl. 2: figs 13-15. [Envir. De Lambèse, Algeria] (Cenomanian, Late Cretaceous) <HT: Colln Heintz 402 (repository unknown)> {DEVIRIES (1973: 71) attributes this name to GAUTHIER (in coll.), but being a manuscript name this name should be attributed to DEVIRIES (1973) according to the ICZN rules}.
- H. arcticus* SHMIDT, 1976: 155-156; pl. 1: figs 7a-b. [Krasnojarskiy Kr., Tanama Valley, near mouth of Yennisey [Enisei] River, Russian SFSR] (Early Santonian, Late Cretaceous) <ST: 8/8846, plus further specimens without number [repository not given]>.
- [*H. bedakensis* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- H. cearensis* BRITO, 1981b: 406-407; pl. 2: figs 1-2. [Jandaira Formation, Praia do Retiro Grande, Aracati, Ceará, Brazil] (Zone of *Hoplitoides* and *Mammites*, Early Turonian) <HT: IGUFP 591 (two casts of that specimens are housed at IGUFRJ 330)>.
- [*H. costatus* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- [*H. curukensis* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

- H. gissarensis* SHMIDT & SIMAKOV, 1953: 68-69; fig. 27; pl. 7: figs 4-6. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Late Turonian) <HT: VNIGRI 125/336>.
- H. himalayensis* MU & WU, 1976: 372; pl. 10: figs 3-10. [Gangba Grp., 7 km east of Gangba, Gangba County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27165, 27166 (repository not given; presumably NIGP)>.
- H. javanicus* SHMIDT & SIMAKOV, 1953: 69-71; figs 28a-b; pl. 3: figs 6-7; pl. 4: fig. 1; pl. 8: figs 5-7. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Late Turonian) <HT: VNIGRI 128/336>.
- [*H. kafirniganensis* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- H. kunlunensis* YANG SHENGQIU, 1991: 121-122; pl. 16, figs 1-11. [Upper Kukebai Fm., Kuzigongsu, 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Late Cretaceous) <HT: NIGP 88413>.
- [*H. pecten* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- [*H. planus* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- [*H. rochatensis* EGOROV, 1972]: 59. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.
- H. rostratus* DEVIRIES, 1973: 73-74; pl. 2: figs 16-18. [Envir. De Batna (Abattoir), Algeria] (Turonian, Late Cretaceous) <HT: Coll'n Heintz 405 (repository unknown)>.
- H. sibiricus* SHMIDT, 1976: pl. 1: fig. 6. [Tscheljabiusk Basin, western Ural, Russian SFSR] (Maastrichtian, Late Cretaceous) <HT: possibly 9/8846, the figured specimen [repository not given]> {only mentioned in plate descripton, but not in the main text}.
- H. simakovi* SHMIDT & SIMAKOV, 1953: 63-65; figs 24a-ж; pl. 4: figs 2-6. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Senonian) <HT: VNIGRI 014a/336>.
- H. subameliae* DEVIRIES, 1973: 74-75; pl. 3: figs 5-10. [Ouled Braham, N of Ain Rafa, Algeria] (Cenomanian, Late Cretaceous) <type-material: Coll'n V. Gauthier (repository unknown, no number given)>.
- H. subbibansensis* DEVIRIES, 1973: 82-84; pl. 5: figs 7-12. [Djebel Bel Kfife and Wadi Cheria, map sheet Ain Beida, Algeria] (Late Cretaceous) <type-material: Coll'n Université de Bâle (no numbers given)>.

- H. subconicus* DEVIRIES, 1973: 75-76; pl. 3: figs 11-14. [Ahmar Khaddou, Bon Milane 4, Algeria] (Senonian, Late Cretaceous) <type-material: Coll'n V. Gauthier (repository unknown, no number given)>.
- H. submirabilis* DEVIRIES, 1973: 76-77; pl. 5: figs 4-6. [El Kantara, Algeria] (Senonian, Late Cretaceous) <HT: Coll'n V. Gauthier 374 (repository unknown)>.
- H. turkestanensis* SHMIDT & SIMAKOV, 1953: 66-68; figs 26a-r; pl. 7: figs 1-3. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Turonian) <HT: VNI-GRI 121/336>.

Mid-Cretaceous

- H. baconicus* SZÖRÉNYI, 1955: 124-125, 259-260; fig. 45, pl. 22, figs 8, 12, 14, 17, 19, 22, 24-25, 27. [calcaire gris lamellé et marne glauconieuse, Bakonyháza (carrière de la vallée Gaja), Bakonyháza (carrière près de la route de Felsőpöre), Olaszfalu-Villóhegy, Olaszfalu-Eperkeshegy, Csöszpuszta, Jásd, and Pénezskút-Körösgyörpuszta, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/545 (from Bakonyháza)>.
- H. estepi* LUCAS, 2000: 101-105; figs 4A-N. [Old Hachita Member, U-Bar Formation, NMMNH locality L-3413 (UTM Zone 13, NAD27, 745525E, 3493350N), L-3414 (UTM 745871E, 3492987N), western flank of U-Bar Ridge, Big Hatchet Mountains, T32S, R16W, Hidalgo County, New Mexico, USA] (*Douvilleiceras mammillatum*-zone, Early Albian, Mid-Cretaceous) <HT: NMMNH P-26515 (from loc. L-34-14); PT: NMMNH P-26501, P-26508 (from loc. L-3413), 26517 (from loc. L-3414)>.
- H. pulcher* SZÖRÉNYI, 1955: 125-126, 260-261; pl. 22, figs 9-10, 13, 18, 23. [groupe de calcaire à *Hippurites*, Sümeg (grande carrière de Gerine), Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/556>.
- H. wayensis* LARRAIN, 1985b: 1402-1405; figs 3.1-3.3, 4.1, 4.2, 5.1, 5.2. [El Way Fm., Antofagasta, Chile] (Aptian) <HT: MZUC 10811 W-32; PT: IIGA-6 W-33, IIGA-392 W-41, IIGA-394 W-42>.

Unknown age

- H. rollandi* var. *oblongus* DEVIRIES, 1973: 86. [Algeria] (unknown) <no types defined, repository unknown>.

Subgenus *Bolbaster* POMEL, 1869

JEFFERY in SMITH & JEFFERY (2000: 320) synonymized the *Bolbaster* POMEL, 1869 with *Hemiasiter* AGASSIZ in AGASSIZ & DESOR, 1847 because they considered the characters used to differentiate these two taxa are not substantive and commonly size related.

Miocene

- H. (B.) callidus* McNAMARA, 1987c: 346-347; text-figs 3, 4, 5B, 6C, 8, 10, 12B; pl. 47: figs 4-8; pl. 48: fig. 7. [Port Campbell Limestone, Sherbrooke River, Port Campbell, Victoria, Australia] (Bairnsdalian, Middle Miocene) <HT: NMV P100503 PT: NMV P100504-05, P100508-09>.
- H. (B.) verecundus* McNAMARA, 1987c: 340-342; text-figs 4, 8; pl. 45: figs 4-5; pl. 46: figs 1-2. [Puebla Fm., Fisherman's Steps, Torquay, Victoria, Australia] (Longfordian, Early Miocene) <HT: NMV P18578; PT: NMV P18761, P20145, P78458>.

Oligocene

- H. (B.) dolosus* McNAMARA, 1987c: 338-340; text-figs 4, 8; pl. 45: figs 1-3, Port Willunga Fm., Maslin's Beach, Aldinga, South Australia] (Willungan, Early Oligocene) <HT: NMV P53172; PT: SAM P26557-59, WAM 86.1206-1209>.

Eocene

- H. (B.) subidus* McNAMARA, 1987c: 336-338; text-figs 3, 4, 5A, 6A, 8, 10, 11A, 12A; pl. 44: figs 1-5; pl. 48: fig. 8. [Tortachilla Limestone, Maslin Beach-Port Willunga district, south of Adelaide, South Australia] (Aldingan, Late Eocene) <HT: SAM P26554; PT: SAM P26555-56, NMV P20484, P53211>.

Paleocene

- B. argentinensis* DEL RÍO, MARTÍNEZ, STILWELL & CONCHEYRO, 2007: 261-264; figs 5E, F, 6D-F. [Jagüel Formation, Malargue Group, Cerros Bayos section (37°40' S, 67°30' W), southern hills of the Salitral Agua de La Perra, 22 km NNE of Colonia 25 de Mayo, Departamento Puelen, La Pampa Province, Argentina] (Danian, NP1 Zone, Early Paleocene) <HT: GHUNL Pam-22890; PT: GHUNL-Pam 22891, 22892>.

Late Cretaceous

- B. compressus* MU & WU, 1976: 372-373; pl. 10: figs 19-23. [Zongshan Fm., Zongshan (= Mt. Zong), Gangba County, Jo-mo glang-ma (Mount Jolmo Lungma Region), Everest Massif, China] (Late Cretaceous) <ST: 27169-27170 (repository not given; presumably NIGP)>.
- H. (B.) hattaensis* ALI, 1989a: 409-410; figs 6(5-8). [Simsima Fm., Gebel El Rowdah, United Arab Emirates] (Late Maastrichtian) < MGD-UAA > {no type specimen defined}.

Subgenus *Leymeriaster* LAMBERT & THIÉRY, 1924

JEFFERY in SMITH & JEFFERY (2000:328) elevated *Leymeriaster* LAMBERT & THIÉRY, 1924 to genus rank.

Maastrichtian

H. (L.) eluvialis VAN DER HAM, 1995: 156-159; text-fig. 3, 4a; pl. 1, figs 1-5; pl. 2, figs 1-6; pl. 3, figs 1-5. [Gulpen Fm. and Maastricht Fm., Maastricht Area, SE Netherlands, NE Belgium] (Late Maastrichtian) <HT: NHMM 1993050; PT: 6 specimens in private collections>.

Campanian

H. (L.) polygonalis TANAKA, 1984a: 435-437; text-figs 7-8; pl. 81, figs 1-2. [Shichi Shale Mb., Anaga Fm., Lower Subgroup of the Izumi Group, Shichiminami, Seidan-cho, Mihara-gun, Hyogo Prefecture, Japan] (Campanian) <HT: GSJ F6063A, B; PT: GSJ F6072>.

Subgenus *Malwaster* CHIPLONKAR & BADVE, 1974

Late Cretaceous

H. (M.) CHIPLONKAR & BADVE, 1974: 52. Type-species: *Opissater subsimilis* FOURTAU, 1918. [India] (Albian to Maastrichtian).

Subgenus *Mecaster* POMEL, 1883

Paleocene

H. (M.) majungensis TANAKA in TANAKA, KANIE & OBATA, 1979: 36-37; fig. 9; pl. 1, figs 6a-d; pl. 2, fig. 1. [C¹⁰ Fm., Majunga area, Northwestern Madagascar] (Danian) <HT: NSM.PA 11994; PT: NSM.PA 11995>.

Santonian

M. arnonensis NEUMANN, 1999: 178-182; fig. 3A-E, 4; pl. 1, figs E-L. [Khurayj Limestone Fm., Wadi al Mujib, south of Dhiban, Jordan] (Santonian) <HT: MBE 2883; PT: MBE 2884>.

Cenomanian

H. (M.) mikasaensis TANAKA, 1984a: 196-198; text-figs 6a-c; pl. 3, figs 1a-e, 2a-c, 3a-b. [Mikasa Fm., Middle Yezo Grp., Ohashi bridge, Ikushumbetsu River, Mikasa City, Central Hokkaido, Japan] (Lower Cenomanian) <HT: NSM-PA5392; PT: NSM-PA5393; GSJ F6144A, B>.

H. (M.) rarus CHIPLONKAR & BADVE, 1972: 146-147; pl. 12: figs 2, 6. [Deola-Chirakhan Marl, Sitapuri (22°20' N, 75°05'30" E), India] (Cenomanian, Late Cretaceous) <HT: MACSG Si 238/69>.

Genus *Aliaster* VALDINUCCI, 1974

Aliaster VALDINUCCI, 1974: 454-462. Type-species: *Opissaster lovisatoi* COTTEAU, 1895. [Mediterranean] (Neogene) {other species included: *Hemiaster cotteaudi* WRIGHT, 1855; *Opissaster jourdyi* PERON & GAUTHIER, 1891; *Opissaster cotteri* de LORIOLE 1896; *Opissaster almerai* LAMBERT, 1906; *Trachyaster aichinoi* CHECCHIA RISPOLI, 1927}.

Genus *Ditremaster* MUNIER-CHALMAS, 1855

Oligocene

D. aslaniani PORETSKAYA, in AKOPJANA, 1974a: 363-364; pl. 186, figs 4a-г. [Armenia] (Early to Middle Oligocene) <HT: LGU 306/8>.

Eocene

D. mereirensis AZAB, 1989: 180-182; pl. 1: figs 1-12. [Midawara Fm., El Mereir Plateau, 14 km south-east of El Sheik Fadl village, Central Egypt (holotype); Gebel Qarara; North Wadi El Sheik; Wadi El Fagirah; Wadi El Rokham; Minqar El Rayan; El Midawara; El Mishgiga; all in Central Egypt] (Earl Mokattamian, Middle Eocene) <HT: repository and number not mentioned (figured on pl. 1: figs 1-4)>.

Genus *Kupeia* MCKNIGHT, 1974

Kupeia MCKNIGHT, 1974: 37-38. Type-species: *Kupeia toi* MCKNIGHT, 1974. [off New Zealand] (Recent).

Recent

K. toi MCKNIGHT, 1974: 38-40; figs 6a-b. [NZOI Stat. D221 (40°06' S, 171°16' E, 688 m depth), D222 (40°38' S, 170°46' E, 651 m depth), D226 (39°54' S, 168°40' E, 823 m depth), D245 (39°54' S, 172°00' E, 532 m depth), E883 (36°00' S, 172°52' E, 999-1046 m depth), New Zealand] (Recent) <HT: NZOI 186 (Stn E883); PT: NZOI P244 (Stn E883), P245 (Stn D221)>.

Genus *Opissaster* POMEL, 1883

HENDERSON (1975: 19-20) placed this genus into the family Schizasteridae.

Miocene

O. tainui HENDERSON, 1975: 22-23; pl. 3, figs 6-9. [Table Mt., Tahora, North Taranaki; Gum Farm, Gower River, North Canterbury; and near Ethleton School, Ethleton,

North Canterbury, New Zealand] (Tongaporutuan, Late? Miocene) <HT: NZGS EC.400; PT: NZGS EC.700-703>.

Genus [*Parahemiaster* EGOROV, 1972]

Parahemiaster EGOROV, 1972: 60. Type-species: none specified. (Late Cretaceous) {Nomen nudum according to ICZN Art. 13.3 (type species fixation missing); included species: *Hemiaster stella* MORTON; *Hemiaster nucleus* DESOR; *Parahemiaster akkaptshigensis* (SHMIDT); *Parahemiaster bobkovae* (SHMIDT); *Parahemiaster soluni* (SHMIDT); *Parahemiaster djabarovi* EGOROV, 1972; *Parahemiaster djaililovi* EGOROV, 1972; *Parahemiaster iljakensis* EGOROV, 1972; *Parahemiaster lucundus* EGOROV, 1972; *Parahemiaster moskvini* EGOROV, 1972; *Parahemiaster schmidtae* EGOROV, 1972; *Parahemiaster subrotundus* EGOROV, 1972}.

Late Cretaceous

[*P. djabarovi* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. djaililovi* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. iljakensis* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. lucundus* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. moskvini* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. schmidtae* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. subrotundus* EGOROV, 1972]: 60. [Tadzhikistan] (Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

Genus *Psephoaster* McNAMARA, 1987c

Psephoaster McNAMARA, 1987c: 347. Type species: *Psephoaster klydonos* McNAMARA, 1987c. [Australia] (Eocene-Miocene).

Miocene

P. klydonos McNAMARA, 1987c: 350; text-figs 7C, 10, 13C; pl. 48: figs 1-3. [Mannum Fm., Murray River, South Australia] (Longfordian, Early Miocene) <HT: SAM P24631; PT: SAM P565, P8933, WAM 86.296, SAM P22017, NMV P13167>.

Oligocene

P. apokryphos McNAMARA, 1987c: 348-50; text-figs 7B, 13BA; pl. 47: figs 9, 11-12; pl. 48: figs 4-6. [Jan Juc Fm., Torquay, Victoria, Australia] (Janjukian, Late Oligocene) <HT: NMV P100506; PT: NMV P100507>.

Eocene

P. lissos McNAMARA, 1987c: 347-48; text-figs 7A, 10, 13A; pl. 47: figs 3, 10. [Kingscote Limestone, Kingscote, Kangaroo Island, South Australia] (Aldingian, Late Eocene) <HT: SAM P26560; PT: SAM P26561>.

Genus *Pseudowashitaster* TANAKA, 1982

Pseudowashitaster TANAKA in TANAKA & KOZAI, 1982: 346-348. Type-species: *Pseudowashitaster mysticus* TANAKA in TANAKA & KOZAI, 1982. [Japan] (Barremian) {other species included: *Washitaster japonicus* TANAKA & OKUBO, 1954}.

Lower Cretaceous

P. mysticus TANAKA in TANAKA & KOZAI, 1982: 348-351; text-figs 3-4; pl. 55, figs 5-7; pl. 56, figs 1-2. [Yunoki Fm., Yunoki, Kahoku-cho, Kami-gun, Kochi Prefecture, Japan] (Upper Barremian) <HT: GSJ 6017; PT: GSJ 6018A, B, GSJ 6019>.

Family Palaeostomatidae LOVÉN, 1867

Genus *Homoeaster* Pomel, 1883

Danian

H. conicus ILIEVA, 1998: 138; pl.1, figs 1a-g. [Byala, Varna District, Bulgaria] (Early Danian) <HT: MGU 360>.

Genus *Leiostomaster* LAMBERT, 1920

Campanian

L. angolanus GREYLING & COOPER, 1995: 63-68; figs 2-5. [Praia Egito, near Quimbala, 70 km N of Lobito, Angola] (Marroti zone, Middle Campanian) <HT: SAfM-PCA2309; PT: SAM-PCA2369, 2392, 2393, 2284>.

Genus *Orthaster* MOSKVIN, 1982

Orthaster MOSKVIN, 1982: 102. Type-species: *Orthaster dagestanensis* MOSKVIN, 1982. [Europe, Asia] (Campanian-Paleocene) {JEFFERY in SMITH & JEFFERY (2000: 357) placed the genus into the family Corasteridae LAMBERT in LAMBERT & THIÉRY, 1924}.

Paleocene

O. dagestanensis MOSKVIN, 1982: 104; figs 1a-c; pl. 10, figs 1a-e. [Dzhinabichay River, Dagestan] (Lower Paleocene) <HT: PIN 3939/1>.

O. okhliensis MOSKVIN, 1982: 105; figs 1g-i; pl. 10, figs 3a-e. [Urma, Dagestan] (Upper Paleocene) <HT: PIN 3939/3> {JEFFERY in SMITH & JEFFERY (2000: 357) placed this species into the synonymy of *O. dagestanensis* MOSKVIN, 1982}.

Campanian

O. alievi MOSKVIN, 1982: 105; figs 1d-f; pl. 10, figs 2a-c. [Alikuliushagy, Bazarchay River, Lesser Caucasus] (Upper Campanian) <HT: PIN 3939/2>.

Genus *Palaeostoma* LOVÉN in A. AGASSIZ, 1872

Pliocene

P. kairukuensis LINDLEY, 2003: 156-158; figs 1d-i, 2. [Kairuku Fm., NW of Aru're village, east coast of Yule Island, Central Province, Papua New Guinea] (Lower Pliocene) <HT: UPNG F1186; PT: UPNG F1185>.

Family Pericosmidae LAMBERT, 1905

Genus *Pericosmus* L. AGASSIZ, 1847

Recent

P. porphyrocardius McNAMARA, 1984: 89-95; figs 1-3. [NNW Port Hedland, Northwestern Australia and ENE of Raine Island, Queensland, Australia] (Recent) <HT: WAM 729.83; PT: WAM 730.83, 731.83, 732.83, 733.83, 734.83; AM J17014; BMNH 1983.3.4.1>.

Miocene

P. borraeus HENDERSON, 1975: 58-59; pl. 17, figs 4-6. [Pakaurangi Point, Kaipara Harbour, New Zealand] (Altonian, Early Miocene) <HT: AU E.315>.

P. celsus McNAMARA & PHILIP, 1984: 329-331; figs 7-9. [Mannum Fm., Mannum, South Australia] (Longfordian, Early Miocene) <HT: SAM P23823; PT: NMV P18354>.

P. (P.) hsui WANG, 1984c: 251-254; pl.: figs 1-9; pl. 2: figs 1-6. [Nankang Sandstone, Shuinantung, Taipei-hsien, northern Taiwan] (Middle Miocene) <HT: CGST E-83003; PT: CGST E-83004 to E-83008>.

P. quasimodo McNAMARA & PHILIP, 1984: 332-335; figs 10-13. [Rutledge Marl Mb., Port Campbell Limestone, Ingles Creek, Port Campbell, Victoria, Australia] (Bairnsdalian, Middle Miocene) <HT: NMV P55512; PT: NMV P55503, 55504>.

P. scaevus HENDERSON, 1975: 59-60; pl. 18, figs 1-3. [Mt. Brown Lst., Weka Pass, New Zealand] (Awamoan, Early Miocene) <HT: CM zfe.33>.

P. torus McNAMARA & PHILIP, 1984: 336-338; figs 14-16. [Batesford Limestone, Batesford, Victoria, Australia] (Batesfordian, Early to Middle Miocene) <HT: NMV P20072; PT: NMV P20071, 20073, 20074, 55502, 55505>.

Eocene

P. annosus HENDERSON, 1975: 60; pl. 16, figs 1-2. [Waiareka Tuff, South Aorere Point, New Zealand] (Kaiatan, Late Eocene) <HT: OU 4760>.

P. farresi CARRASCO, 2003: 25-27; figs 2 a-f. [Gurb de la Plana, near Vic, 65 km N of Barcelona, Spain] (Eocene) <HT: MGSB 67676; PT: MGSB 67276>.

Subgenus *Lambertona* SÁNCHEZ ROIG, 1953

HENDERSON (1975: 24) restored this taxon to genus rank and referred it to the family Schizasteridae.

Miocene

L. perplexa HENDERSON, 1975: 26-27; text-fig. 5b; pl. 6, figs 1-2; pl. 7, figs 1-3. [Blackhead, Dunedin; west side of Burnt Hill, Canterbury; Bushy Park, Palmerston; and Caversham Sdst., Matanaka, New Zealand] (Otaian or Altonian to Waiauian, Miocene) <HT: OU 8845; PT: AU E.311a-c; OU 6835, 8550>.

Oligocene

L. perditia HENDERSON, 1975: 27-28; text-fig. 5a; pl. 8, figs 1-2. [Canterbury, New Zealand] (? Oligocene to Early Miocene) <HT: CM zfe.288>.

Family Paleopneustidae A. AGASSIZ, 1904

Genus *Eopericosmus* MARKOV & SOLOVJEV, 2001

Eopericosmus MARKOV & SOLOVJEV, 2001: 16, 81-82. Type-species: *Eopericosmus typicus* MARKOV & SOLOVJEV, 2001. [Kazhakstan] (Paleocene) {other species included: *Brissopsis alta* HUTTON, 1873}.

Paleocene

E. sveshnikovi MARKOV & SOLOVJEV, 2001: 22-26, 83-84; fig. 4a-б; pl. 1, figs 1a-б, 2a-b. [Mangyshlak, Kazhakstan] (Montian, Early Paleocene) <HT: PIN 4772/49>.

E. typicus MARKOV & SOLOVJEV, 2001: 17-22, 82-83; fig. 3a-д; pl. 1, figs 3a-г, 4. [Mangyshlak, Kazhakstan] (Montian, Early Paleocene) <HT: PIN 4772/2>.

Superfamily Paleopneustoidea MARKOV & SOLOVJEV, 2001

Paleopneustoidea MARKOV & SOLOVJEV, 2001: 80. {included families: Paleopeustidae, Prenasteridae, Schizasteridae}.

Family Schizasteridae LAMBERT, 1905

Genus *Abatus* TROSCHER, 1851

Eocene

A. kieri MCKINNEY, MCNAMARA & WIEDMAN, 1988: 502-503; figs 3, 1-2. [Eastern Seymour Island, Antarctic Peninsula] (Late Eocene) <HT: USNM 416181; PT: USNM 416182>.

Genus *Agassizia* L. AGASSIZ & DESOR, 1847

Miocene

- A. algarbiensis* FERREIRA, 1962: 293-295; pl. 1, figs 1-6. [Forte de S. João de Ferragudo, Algarve, Portugal] (Middle Miocene) <ST: Museu dos Serviços Geológicos de Portugal, repository no. not given>.
- A. eugeniae* BRITO & RAMIRES, 1974: 270-272; pl. 3, figs 1-5; pl. 4, figs 1-3. [Pirabas Fm., procedentes de Capanema, Pará, Brazil] (Lower Miocene) <HT: MN 5236-I; PT: MN 5237-I>.
- A. (A.) powersi* KIER, 1972a: 97-98; pl. 64, figs 1-8; pl. 65, figs 1-2. [Dam Fm., localities S-357, S-360, S-361 and S-568, Saudi Arabia] <HT: USNM 170497; PT: USNM 170498-499>.

Subgenus *Anisaster* POMEL, 1886

Eocene-Oligocene

- Ag. (An.) arabica* KIER, 1972a: 96-97; pl. 65, figs 3-8; pl. 66, figs 1-3. [localities S-761 and S1603, Saudi Arabia] (Eocene-Oligocene?) <HT: USNM 170501; PT: USNM 170500>.
- Ag. (An.) wilmingtunica inflata* KIER, 1980: 44; fig. 19; pl. 15, figs 8-10. [Santee Limestone, Georgetown localities 37, 44, South Carolina, USA] (Middle Eocene) <HT: USNM 264073>.

Genus *Aguayoaster* SÁNCHEZ ROIG, 1952

Eocene

- A. schickleri* DONOVAN & ROWE, 2000: 656-658; fig. 2. [Swanswick Fm., Pimento Hill, Beecher Town, Jamaica] (mid Middle to low Upper Eocene) <HT: BMNH EE 6340>.

Genus *Amphipneustes* KOEHLER, 1900

Recent

- A. davidi* MADON-SENEZ, 2002: 52-55; figs 1a-d, 2a-b, 3a-e. [Belgian & Dutch Antarctic Expedition 1959-196, Iris Mission (70°19' S, 24°12' E to 70°17' S, 24°06' E), and Belgian & Dutch Antarctic Expedition 1964-1965, Glacier Bay, Stat. 219 (70°18'5" S, 23°58'0" E), Stat. 223 (70°13'2" S, 23°55'1" E), Antarctic Ocean, 207 and 216 m

depth] (Recent) <HT: MNHN EcEh 9362 (female); PT: MNHN EcEh 9363-9369 (3 males, 4 females)>.

- A. *mironovi* MARKOV, 1991: 154-155; figs A-C. [“Академик Курчатов”-cruise (“Академик Курчатов”-cruise), Stat. 870 (55°08' S, 25°03' W, depth 4689-4704 m), South Sandwich Islands, South East Atlantic Ocean] (Recent) <HT: IOAN, no specimen no. given> {re-described in MARKOV (1994: 86-88; figs 25a)}.

Genus *Brachysternaster* LARRAIN, 1985a

Brachysternaster LARRAIN, 1985a: 121. Type-species: *Brachysternaster chesheri* LARRAIN, 1985a. [South Shetland Islands] (Recent).

Recent

- B. *chesheri* LARRAIN, 1985a: 121-123; figs 1-8. [USNS Eltanin Stat. 410, South Shetland Islands (61°18'-61°20' S, 56°09'-56°10' W, depth 220-240 m), Stat. 437 (62°50'-62°51' S, 60°40'-60°35' W, depth 267-311 m); RV Eastwind Stat. EW-66-009 (62°43.1' S, 62°17.1' W, depth 305 fathoms), Stat. EW-66-009A (62°43.1' S, 62°17.5' W, depth 560 m), Stat. EW-66-038 (61°14.8' S, 54°48' W, depth 105 fathoms)] (Recent) <HT: USNM E11025 (male spec.; Eltanin Stat. 410); PT: USNM E11254 (female spec.; RV Eastwind Stat. EW-66-038), E11136 (juvenile; Eltanin Stat. 410)>.

Genus *Brisaster* GRAY, 1855

Recent

- B. *tasmanicus* MCKNIGHT, 1974: 40; figs 7a-b. [NZOI Stat. E399 (46°10' S, 171°33' E, 1222 m depth), E788 (44°00' S, 168°11' E, 1184-1193 m depth)] (Recent) <HT: NZOI 187 (Stat. E788); PT: NZOI P246 (Stat. E399)>.

Paleocene

- B. *latus* MARKOV, 1994: 23-24, 34, **72-73**; figs 19e, 19з; pl. 3, figs 6a-б. [Усак (Usak), Mangyshlak Peninsula, Kazakhstan] (Mons, Paleocene) <HT: PIN N 4351/336>.
- B. *micropetalus* MARKOV, 1994: 23, 33-34, **69-72**; figs 19а-д, ж; pl. 3, figs 2а-г, 3а-б, 4. [Усак (Usak), Mangyshlak Peninsula, Kazakhstan] (Danian, Paleocene) <HT: PIN N 4351/295>.
- B. *moskvini* MARKOV, 1994: 68-69; figs 19а; pl. 3, figs 5а-б. [щелье Шахбагата, Mangyshlak Peninsula, Kazakhstan] (Danian, Paleocene) <HT: PIN N 4351/286>.

Genus *Calzadaster* CARRASCO, 2005

Calzadaster CARRASCO, 2005: 50-51. Type-species: *Calzadaster friasi* CARRASCO, 2005. [South-western Europe] (Eocene).

Eocene

C. friasi CARRASCO, 2005: 51-54; figs 2, 3a-d, 4; pl. 1: figs A-E. [Sant Julià de Vilatorra, 60 km N Barcelona, Spain] (Late Lutetian, Middle Eocene) <HT: MGSB 67754a; PT: MGSB 67754b>.

Genus *Caribbaster* KIER, 1984b

Caribbaster KIER, 1984b: 67-68. Type-species: *Prenaster loveni* COTTEAU, 1875. [Cuba and Jamaica] (Eocene) {other species included: *Schizaster dyscritus* ARNOLD & CLARK, 1927}.

Genus *Diploporaster* MORTENSEN, 1950

Miocene

D. flemingi HENDERSON, 1975: 18-19; text-figs 2a-b; pl. 2, figs 3-6. [Waikuku Lst., North Cape and Pakaurangi Fm., Northland, New Zealand] (Otaian – Tongaporutuan, Miocene) <HT: AU E.332; PT: NZGS EC.470, AU E.313>.

Genus *Hemifaorina* JEANNET & MARTIN, 1937

Paleocene

H.? rex McNAMARA, 1993a: 333-335; figs 1, 2A. [Kings Park Fm., Caisson N6 L3, Swan River, Perth, Western Australia] (Lower Paleocene; planktonic foraminiferal zone P4) <HT: WAM 71.1518; PT: WAM 71.1502-71.1507>.

Genus *Kina* HENDERSON, 1975

Kina HENDERSON, 1975: 28. Type-species: *Kina gracilis* HENDERSON, 1975. [New Zealand] (Eocene-Oligocene).

Paleogene

K. gracilis HENDERSON, 1975: 28-29; text-fig. 7; pl. 8, fig. 7; pl. 9, figs 1-3. [Culverden; Kahurangi Lighthouse; Bullock Creek; and Waihao Lst., Eaihao River, New Zealand]

(Kaiatan to Waitakian, Late Eocene to Oligocene) <HT: CM zfe.276; PT: NZGS EC.423, 454, 495, 570, 574>.

Genus *Linthia* DESOR, 1853

Oligocene

L.? *summesbergeri* KROH, 2005: 161-163; fig. 71; pl. 68, figs 2-7. [Ebelsberg Formation, Weikerlsee, in the eastern part of Linz, Upper Austria, Austria] (Egerian (Chattian to Aquitanian), Late Oligocene to Early Miocene) <HT: NHMW 2003z0026/1187a (cast of the aboral side on a slab together with 3 other specimens, two of which are designated as paratypes); PT: NHMW 2003z0026/1178, 1187b (cast of the aboral side), 1187c (cast of the oral side), 1190 (cast of the aboral side), 1212 (cast of the oral side; 1213 is the corresponding mould), 1226 (cast of the oral side)>.

Eocene

L. africana MAMEDOV & MELIKOV, 1976: 157-158; pl. 1, figs 1a, 1б, 1в. [In-Arieh, Republic Mali] (Middle Eocene) <HT: AzINEFTEKHIM 2/69>

L. (L.) harmatuki KIER, 1980: 45-46; figs 20-21; pl. 16, figs 1-2; pl. 17, figs 1-2. [Castle Hayne Limestone, Maple Hill localities 10, 34, North Carolina; Santee Limestone, Georgetown locality 37, South Carolina; USA] (Middle Eocene) <HT: USNM 264076; PT: USNM 264074, 264075>.

L. maliensis MAMEDOV & MELIKOV, 1976: 159-161; figs 1-3; pl. 1, figs 2a, 2б, 2в. [In-Arieh (Tahanabat), Republic Mali] (Middle Lutetian, Eocene) <HT: AzINEFTEKHIM 10/69>

L. monteroae KIER, 1984b: 76-77; pl. 40, figs 1-4. [Palmer loc. 1085; E of Arroyo Blanco, 150 m, in road of Majagua, Camagüey Province, Cuba] (Eocene) <HT: ANSP 16656>.

L. pulchra McNAMARA, 1985a: 162-164; fig. 1. [Tortachilla Limestone, southern Maslin Beach, South Australia] (Late Eocene) <HT: NMV P20455; PT: GSWA F5828; WAM 66.637, 85710, 85.711>.

Paleocene

L. bajsarensis BAJARUNAS in MARKOV, 1994: 58-61; figs 15д, 16а-д; pl. 1, figs 3а-г; pl. 2, figs 1а-г, 2-3. [Байсарлы (Bagsarly), Mangyshlak Peninsula, Kazakhstan] (Mons, Paleocene) <HT: PIN N 4351/84>.

L. brevipetala MARKOV, 1994: 61-62; figs 15в-г; pl. 1, figs 2. [Кеньырли (Kyendyrli), Mangyshlak Peninsula, Kazakhstan] (Mons, Paleocene) <HT: PIN N 4351/76>.

L. ghiroboensis TESSIER & ROMAN, 1973: 157-158; pl. 2, figs 4-7. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Paleocene, Thanetian) <HT: IPM 1972-9F>.

Maastrichtian

- L. (L.) madagascariensis* TANAKA in TANAKA, KANIE & OBATA, 1979: 41-43; fig. 12a-c; pl. 3, figs 2a-d, 3, 4. [C⁹ Fm., Majunga area, Northwestern Madagascar] (Maastrichtian) <HT: NSM.PA 11996; PT: NSM.PA 11997, 11998>.
- L. romani* BRITO, 1981c: 576; pl. 3, figs 1-6. [Gramame Fm., Pedreira da Fábrica de Cimento Poty, Paulista, Permanbuco, Brazil] (Maastrichtian) <HT: DGM do DNPM 5569; PT: DGM do DNPM 5570>.

Genus *Neoproraster* MARKOV, 1994

Neoproraster MARKOV, 1994: 82-83. Type-species: *Neoproraster bajarunasi* MARKOV, 1994. [Kazakhstan] (Paleocene-Eocene).

Palaeogene

- N. bajarunasi* MARKOV, 1994: 24-25, 34, 85-86; figs 24д-з; pl. 4, figs 3а-г. [Санды (Sandyk), Mangyshlak Peninsula, Kazakhstan] (Early Eocene) <HT: PIN N 4351/388>.
- N. usakensis* MARKOV, 1994: 83-84; figs 24а-г; pl. 4, figs 2а-б. [Усак (Usak), Mangyshlak Peninsula, Kazakhstan] (Mons, Paleocene) <HT: PIN N 4351/385>.

Genus *Prenaster* DESOR, 1853

Eocene

- P. desori caucasica* PORETSKAYA, in AKOPIANA, 1974: 364-365; pl. 185, figs 2а-в, 3. [Armenia] (Late Eocene) <HT: LGU 306/20>.
- P. synapticus* HENDERSON, 1975: 20-21; text-figs 3; pl. 2, figs 7, 9-12. [Waiareka Volcanics, Lorne Railway Station, New Zealand] (Kaiatan, Late Eocene) <HT: NZGS EC.456; PT: NZGS EC.699>.

Genus *Proraster* LAMBERT, 1895

Paleocene

- P. oedumi* STOKES, 1975: 24. [Kjbenhavens Havn, Knippelsbro, Copenhagen, Denmark] (Late Danian) <HT: Ødum coll. 7647> {replacement name for *Micraster desori* ØDUM, 1926 non HÉBERT, 1856}.

Late Cretaceous

P. granti KIER, 1972a: 84-87; fig. 45; pl. 47, figs 6-7; pl. 48, figs 1-6. [Aruma Fm., locality S-748, Saudi Arabia] (Campanian) <HT: USNM 170461; PT: USNM 170460>.

P. magnus MARKOV, 1994: 66-67; figs 18a-б; pl. 3, figs 1a-б. [Камысты (Kamysty), Mangyshlak Peninsula, Kazakhstan] (Maastrichtian) <HT: PIN N 4351/547>.

Genus *Protenaster* POMEL, 1883

FISCHER (1966: U576) regarded *Protenaster* as subgenus of *Prenaster* DESOR, 1853; McNAMARA (1985b:312-313) rejects this.

Oligocene

P. philipi McNAMARA, 1985b: 322-324; text-fig. 7; pl. 34: figs 6-7. [Waurm Ponds Limestone, Waurm Ponds, Victoria, Australia] (Janjukian, Late Oligocene) <HT: MUGD 1690; PT:NMV P4776, P19990, P63040, P63059, P63064>.

Eocene

P. preaustralis McNAMARA, 1985b: 321; text-figs 5-6; pl. 33: figs 6-8; pl. 34: figs 1-5. [Tortachilla Limestone, Maslin Beach – Port Willunga district, South Australia] (Late Eocene) <HT: SAM P24519; PT: SAM P24520-24523, NMV P71348, P71349>.

Genus *Pseudolinthia* MARKOV, 1994

Pseudolinthia MARKOV, 1994: 80. Type-species: *Pseudolinthia triporata* MARKOV, 1994. [Kazakhstan] (Paleocene).

Paleocene

P. triporata MARKOV, 1994: 25-26, 34-35, 80-81; figs 23a-b; pl. 4, figs 1a-б. [Капам (Kapam), Mangyshlak Peninsula, Kazakhstan] (Danian, Paleocene) <HT: PIN N 4351/362>.

Genus *Schizaster* AGASSIZ, 1836Subgenus *Schizaster* AGASSIZ, 1836

Pliocene

S. (S.) alphonsei LINDLEY, 2003: 154-156; figs 1a-c. [Kairuku Fm., NW of Aru're village, east coast of Yule Island, Central Province, Papua New Guinea] (Lower Pliocene) <HT: UPNG F1179>.

Miocene

S. scherzeri GABB, 1881: 348; pl. 45: figs 28, 28a-b. [above Paquare, on the Reventazon River and at Sapote, Costa Rica] (Miocene) <repository and specimen nos. not given>.

Oligocene

S. (S.) halli McNAMARA & PHILIP, 1980a: 52-53; figs 3 A-D. [Table Cap Beds, Table Cap; Cape Grim Beds, Cape Grim, Tasmania] (Janjukian (Late Oligocene) to Longfordian (Early Miocene)) <HT: NMV P19036; PT: NMV P53219, P53220>.

Eocene

S. archiaci complanatus CHAVANON, 1974: Vol.1: 237; pl. 16: figs 2a-d. [Saint-Palais and Couquèques, Aquitaine Basin, western France] (Middle to Late Eocene) <ST: Castex coll'n, Degrange-Touzin coll'n, Fabre coll'n; no repository nos given for types; figured specimen: 28.C.7>.

S. (S.) caddoensis ZACHOS in ZACHOS & MOULINEUX, 2003: 499; figs 6.4-6.7. [Reklaw and Weches Formations, Claiborne Group, Localities 81 and 82, Cherokee and Milam Counties, and Locality 54, Nacogdoches County, Texas, USA] (Middle Eocene) <HT: TMM 2000TX1 (from Locality 54); PT: BEG 17749 (from Locality 81)>.

S. formelli KIER, 1984b: 43-44; pl. 24, figs 1-7. [Palmer loc. 687a, 24 km E of Camagüey on Maraguan road from Guanabanito River, Camagüey Province, Cuba] (Eocene) <HT: ANSP 16675a; PT: ANSP 16675b>.

S. haleifiensis ROMAN & STROUGO, 1988: 148. [Bir Haleifia, Sinai, Egypt] (Priabonian). {replacement name for *S. humei* LAMBERT, 1932 non FOURTAU, 1909}.

S. rindensis PORETSKAYA, in AKOPIANA, 1974a: 366; pl. 188, figs 1a-6. [Rind, Armenia] (Late Eocene) <HT: LGU 306/39>.

S. (S.) stenzeli ZACHOS in ZACHOS & MOULINEUX, 2003: 499-501; figs 6.9, 6.13. [Weches Formation, Claiborne Group, Localities 32, 33, 34, 36, Leon County, Texas, USA] (Middle Eocene) <HT: TMM 1988TX1 (from Locality 34); PT: TMM 1988TX2, 1988TX3 (from Locality 34), 2006TX1 (from Locality 36)>.

Palaeogene

S. tarimensis YANG SHENGQIU, 1991: 123; pl. 17, figs 1-8; pl. 20, figs 1-4. [Wulagen Fm. Bashibulake, Wuqia County, Tarim Basin, China] (Early Tertiary) <HT: NIGP 88424>.

Subgenus *Dipneustes* ARNAUD, 1891

Miocene

S. (D.) fosteri McNAMARA & PHILIP, 1980a: 58-59; figs 7 A-B. [?Mannum Fm., Murray River at Swan Reach, South Australia] (?Longfordian (Early Miocene)) <HT: NMV P55472>.

Subgenus *Ova* GRAY, 1825

Recent

S. (O.) myorensis McNAMARA & PHILIP, 1980b: 133-140; figs 5A-E, 7A-c, 8A-C, 9A. [Myora, Moreton Bay, Queensland, Australia] (Recent) <HT: QM G12063; PT: QM G3870-1, 11826-8, 12064-6, 12094, 12428-48, 3809>.

S. (O.) portjacksonensis McNAMARA & PHILIP, 1980b: 141-143; figs 9B, 10A-Gport Jackson, Australia] (Recent) <HT: AM J1732; PT: AM J1731>.

Subgenus *Paraster* POMEL, 1869

Recent

P. doederleini CHESHER, 1972: 44866; figs 1-7, 9, tabs. 1-2. [Pillsbury Stat. 402 (8°51' N, 77°02' W, off northern Columbia, depth 73 m), Pillsbury Stat. 399 (9°01' N, 76°39' W, off northern Columbia, depth 147 m); White Shoal, Dry Tortugas, Florida; RHCK 45, off Aligator Reef, Florida Keys; Gerda Stat. 683 (25°52' N, 77°53,5' W, Northwest Providence Channel, Bahama Islands, depth 250 m)] (Recent) <ST: MCZ 8396-8397, 9398, USNM E11376-E11378>.

S. (P.) ovatus LINDLEY, 2004: 131-132; figs 6a-d. [Cape Gazelle, New Britain, East New Britain Province, Papua New Guinea] (Recent) <HT: ANU 60653>.

Pliocene

S. (P.) ashrafi ALI, 1985: 289; figs 7B-E. [Wadi Gabir, Western Egypt] (Early Pliocene) <HT: USNM 339745; PT: USNM 339746>.

Miocene

[*S. (P.) jhadwensis* TANDON & SRIVASTAVA, 1988]: 153. [India] (*Spiroclypeus ranjanae* Zone, Miocene) <unknown> {nomen nudum, name mentioned only}.

Eocene

- S. (P.) incorrectus* ELATTAAR & STROUGO, 2001: 74-76; figs 8i-8l, 9c. [Bir Haleifiya, west-central Sinai, Egypt] (Late Eocene) <n/a> {nomen novum pro *Schizaster humei* LAMBERT, 1932, non FOURTAU, 1909}.
- S. (P.) susana* ZACHOS in ZACHOS & MOULINEUX, 2003: 501-503; figs 6.10–6.12, 6.14, 6.15. [Caddell Formation and Moodys Branch Formation, Jackson Group, Localities 1, 2 and 3; Angelina and San Augustine Counties, Texas; and Montgomery Landing, Grant County, Louisiana, USA] (Late Eocene) <HT: TMM 886TX1; PT: TMM 886TX2, 886TX3 (all from Locality 2)>.
- S. (P.) tatei* McNAMARA & PHILIP, 1980a: 51-52; figs 2 A-E. [Tortachilla Limestone, Maslin Beach – Port Willunga district, South Australia] (Late Eocene) <HT: NMV P55467; PT: NMV P55468, P55470>.

Paleocene

- S. (P.) carinatus* McNAMARA & PHILIP, 1980a: 50-51; figs 1 A-C. [Pirie Calcarenite, Giaralia Range, Carnarvon Basin, Western Australia] (Middle to Late Paleocene) <HT: CPC 4822; PT: CPC 4823, 4824> {JEFFERY in SMITH & JEFFERY (2000: 336) suggested that this species might be a synonym of *Paraster chargensis* (WANNER, 1902)}.
- S. (P.) rougeriei* TESSIER & ROMAN, 1973: 158-160; pl. 2, figs 8-16. [Falaises à l'ouest de l'ancien de Fresco, Côte d'Ivoire] (Paleocene, Thanetian) <HT: IPM 1972-9BGc>.

Palaeogene

- S. (P.) extumidus* YANG SHENGQIU, 1991: 124; pl. 19, figs 6-7. [Upper Qimugen Fm., Kuzigongsu, 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Early Tertiary) <HT: NIGP, no holotype defined>.

Genus *Schizocosmus* MARKOV, 1990

Schizocosmus MARKOV, 1990: 55. Type-species: *Pericosmus abatoides* CLARK, 1925. [South Atlantic Ocean] (Recent).

Genus *Tripylus* PHILIPPI, 1845

Recent

- T. beatriceae* LARRAIN, 1986: 116-118; figs 2a-f, 3a-c. [RV "Eltanin" Sta. 126 and 353, south east of Cape Horn] (Recent) <HT: USNM E11008; PT: USNM E11018, E14618>.

Genus *Waurnia* McNAMARA & PHILIP, 1984

Waurnia McNAMARA & PHILIP, 1984: 341-342. Type-species: *Pericosmus nelsoni* McCoy, 1882. [Waurn Ponds Limestone, Victoria, Australia] (Oligocene).

Subfamily Brisasterinae MARKOV, 1994

Brisasterinae MARKOV, 1994: 65. Type-genus: *Brisaster* GRAY, 1855. [Australia, New Zealand] (Cenomanian-Recent) {included genera: *Proraster* LAMBERT, 1895; *Pseudolinthia* MARKOV, 1994; *Neoproraster* MARKOV, 1994; *Amphipneustes* KOEHLER, 1900; *Schizocosmus* MARKOV, 1990}.

Family Aeropsidae LAMBERT, 1896

Genus *Sphenaster* JEFFERY in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999

Sphenaster JEFFERY in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 131. Type-species: *S. larumbensis* JEFFERY in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999.

Paleocene

S. larumbensis JEFFERY in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 131-132; figs 42a-d; pl. 11, figs 8-12. [Coraster Beds, Larumbe, Navarra, Spain] (Lower Thanetian) <HT: BMNH EE6073> {based on a single specimen}.

Suborder Micrasterina FISCHER, 1966

Family Micrasteridae LAMBERT, 1920

Genus *Brissopneustes* COTTEAU, 1886

In JEFFERY (1998: 150) the genus *Brissopneustes* is considered as a junior synonym of *Cyclaster* COTTEAU in COTTEAU & LEYMERIE, 1856.

Paleocene

B. vindobonensis KÜHN, 1930: 553-554; pl. 1: figs 8-10. [Bruderndorf Formation, Bruderndorf, Lower Austria, Europe] (Upper Danian) <HT+PT: NHMW 1930V8> {placed into the synonymy of *Cyclaster aturicus* (SEUNES, 1888) by KROH (2001: 404)}.

Maastrichtian

- B. ruegensis* [nom. corr. pro *rügensis*] KUTSCHER, 1978b: 1027-1028, pl. 2, figs 1-5. [Saßnitz, Quarry Wittenfelde, Rügen, Germany] (Late Early Maastricht) <HT: SGWG 60/1>.

Genus *Diplodetus* SCHLÜTER, 1900Subgenus *Protobrissus* LAMBERT, 1907

Paleocene

- D. (Protobrissus) kurkurensis* AZAB & ELATTAAR, 1999: 863-864; tab. 23; pl. 3: figs 16-19. [Gebel El Barga (W of Daraw), Gebel Gharra (W of Aswan) and Kurkur section (at the entrance of Wadi Kurkur), Kurkur area, Egypt] (*Gitolampas abundans* Zone, Late Paleocene) <repository and specimen nos. not given>.
- P. rionensis* MOSKVIN, 1964: 199-201; figs 10-11; pl. 4: figs 3a, б; pl. 5: figs 1a-b, 2a-d. [Western Georgia] (Late Paleocene) <not given> {referred to *Pseudogibbaster* by MOSKVIN (1983)}.

Genus *Isomicraster* LAMBERT, 1901

Late Cretaceous

- I. babatakensis* EGOROV, 1972: 60. [Tadzhikistan] (Maastrichtian, Late Cretaceous) <repository unknown> {nomen novum pro *Epiaster nobilis* SHMIDT, non STOLICKA}.

Genus *Jordaniaster* NEUMANN, 1999

- Jordaniaster* NEUMANN, 1999: 176. Type-species: *J. husseini* NEUMANN, 1999. [Jordan] (Cenomanian).

Cenomanian

- J. husseini* NEUMANN, 1999: 176-178; fig. 2A-D; pl. 1, figs A-D. [Rumeimin Fm., Wadi Salihi, near Rumeimin, north of Amman, Jordan] (Lower Cenomanian) <HT: MBE 2882>.

Genus *Micraster* L. AGASSIZ, 1836

Campanian-Maastrichtian

M. amnicus GREYLING, 1996: 25-32; figs 2-7. [St. Lucia Fm., ESE Mtubatuba, Zululand, South Africa] (Campanian-Maastrichtian) <HT: DNSM-PCZ5892; PT: DNSM-PCZ1685-87, 5887-89>.

Campanian

M. coranguinum simpsoni STOKES, 1975: 65-66; fig. 29f; pl. 3, figs 1-5. [Saltdean, Sussex and Kent, England] (*Marsupites* and *O. pilula* Zone, Lower Campanian) <HT: R. Simpson coll. S54>.

M. coravium POSLAVSKAYA, 1959: 289; pl. 22, figs 1a-д (97). (Early Campanian) <no repository information provided> {cited and illustrated in MELIKOV (in ALI-ZADE, 1988: 208, fig. 60, pl. 8, figs 4a-д), cited by KIER & LAWSON (1978: 110) as *M. coravium* MOSKVIN, 1959, unclear whether this is the same species or a homonym}.

M. solignaci STOKES, 1975: 79; fig. 30e; pl. 11, figs 3-6. [Koudiat Melhab, Tunisia] (Campanian) <HT: coll. Lambert, no specimen no. given>.

M. westlakei STOKES, 1975: 81; fig. 30h; pl. 12, figs 4-6. [Tichbourne Farm, Hampshire, England] (*B. mucronata* Zone, Upper Campanian) <HT: CWUS 3320; PT: CWUS 3321-22>.

Santonian

M. (M.) maleckii HYNDA & MACZYŃSKA, 1979: 22-23; figs 1a-g, 2a-d; pl. 1, figs 1-2; pl. 2, figs 1-3; pl. 3, figs 1-2. [Korzkiew near Cracow, Southern Poland] <HT: PASME MZ VIII Ee960>.

Coniacian

M. (M.) praerogalae OLSZEWSKA-NEJBERT, 2007: 23-55; textf-fig. 54; pl. 27: figs 1a-d. [Shakh-Bogota, about 15 km south of Sarytash, Mangyshlak, Kazakhstan] (*Magadiceramus subquadratus* Zone, Late Coniacian, Late Cretaceous) <HT: IGPUW/E/01/677; PT: IGPUW/E/01/678>.

Late Cretaceous, undifferentiated

M. turkestanensis SHMIDT & SIMAKOV, 1953: 71-72; fig. 29; pl. 4: figs 12-14. [Uzbekistan-Tajikistan-Kyrgyzstan border region, Central Asia] (Senonian) <HT: VNI-GRI 134/336> {transferred to *Sogdiaster* by EGOROV (1972: 59)}.

Subgenus *Gibbaster* GAUTHIER, 1887

Santonian

M. (G.) sonorensis BUITRÓN, 1971: 40-41; pl. 12, figs 3, 5, 7-9. [Cerro de Las Conchas, Arivechi, Sonora, Mexico] (Santonian ?, Late Cretaceous) <HT: IGMUC 2280>.

Genus *Ovulaster* COTTEAU, 1884

SMITH, GALLEMÍ, JEFFERY, ERNST & WARD (1999: 130) placed this genus into the family Corasteridae LAMBERT in LAMBERT & THIÉRY, 1924.

Paleocene

O. protodecimae GIUSBERTI, FANTIN & BUCKERIDGE, 2005: 457-459; figs 2a-b; pl. 1: figs 1-12; pl. 2: figs 6-9. [Scaglia Rossa Formation, Forada creek, near the village Villa di Villa, Lentiai, Belluno, Venetian Prealps, NE Italy] (calcareous nannoplankton zones NP1 and NP4, Danian, Early Paleocene) <HT: MGPD 28877; PR: MGPD 28878, 29013, 29014>.

Maastrichtian

O. reticulatus SMITH & GALLEMÍ in SMITH, GALLEMÍ, JEFFERY, ERNST & WARD, 1999: 130; figs 39a-d, 40a-b; pl. 10, figs 6-10. [Sarasate, Navarra, Spain] (Lower Maastrichtian) <HT: BMNH EE6236; PT: BMNH EE6074>.

Senonian

O. elevatus TZANKOV, 1984: 114-115; pl. 49, figs 4, 4a-c. [Kondel, Dragoman, de Sofia, Bulgaria] <HT: USC CR₂ 1319>.

Genus *Plesiaster* POMEL, 1883

Late Cretaceous

[*P. gibberiensis* EGOROV, 1972]: 60. [Tadzhikistan] (Maastrichtian, Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

[*P. tschaltauensis* EGOROV, 1972]: 60. [Tadzhikistan] (Maastrichtian, Late Cretaceous) <no specimens mentioned> {nomen nudum (no description, illustration, or reference to such included)}.

P. vancouverensis SMITH & MCGUGAN, 1996: 104-107; text-figs 3-5. [Haslam Fm., French Creek, Vancouver Island, British Columbia, Canada] (uppermost Santonian-lowermost Campanian) <HT: BMNH EE5078; PT: BMNH EE5076-77, EE5079-83>.

Genus *Pseudogibbaster* MOSKVIN, 1983

Pseudogibbaster MOSKVIN, 1983: 115; figs 2-3. Type-species: *Protobrissus akkajensis* MOSKVIN & POSLAVSKAYA, 1959. [Crimea, Caucasus and Trans-Caspina Region] (Paleocene) {other species included: *Micraster depressus* KONGIEL, 1937; *Micraster ter-censis* COTTEAU; *Protobrissus indolensis* POSLAVSKAYA & MOSKVIN, 1960; and *Protobrissus rionensis* MOSKVIN, 1964}.

Genus *Sogdiaster* EGOROV, 1972

Sogdiaster EGOROV, 1972: 59. Type-species: *Micraster turkestanensis* SHMIDT, 1953. [former USSR] (Late Cretaceous).

Superfamily Brissidea STOCKLEY, SMITH, LITTLEWOOD,
LESSIOS & MACKENZIE-DODDS, 2005

Brissidea STOCKLEY, SMITH, LITTLEWOOD, LESSIOS & MACKENZIE-DODDS, 2005: 457, 458; fig. 2. {included families: Brissidae, Spatangidae, Loveniidae}.

Family Brissidae GRAY, 1855

Genus *Amoraster* MCNAMARA & AH YEE, 1989

Amoraster MCNAMARA & AH YEE, 1989: 178-179. Type-species: *A. paucituberculata* MCNAMARA & AH YEE, 1989. [South Australia] (Miocene).

Miocene

A. paucituberculata MCNAMARA & AH YEE, 1989: 179-181; figs 2a-b, 3a-b, 4a-e. [Port Campbell Limestone, Portland, Victoria, South Australia] (Bairnsdalian-Mitchellian, Middle-Late Micoene) <HT: WAM 87.303; PT: WAM 85.1271, 87.522, 87.523>.

A. tuberculata MCNAMARA & AH YEE, 1989: 181-183; figs 6a-d. [Mannum Fm., Murray River, South Australia] (Longfordian, Early Micoene) <HT: WAM 87.116; PT: WAM 86.322b, d-g>.

Genus *Antiquibrissus* SZÖRÉNYI, 1955

Antiquibrissus SZÖRÉNYI, 1955: 126-127, 261-262. Type-species: *Antiquibrissus suemeg-ensis* [*süimegensis*] SZÖRÉNYI, 1955. [Hungary] (Mid-Cretaceous).

Cretaceous

- A. suemegensis* [nom. corr. pro *süemegensis*] SZÖRÉNYI, 1955: 127, 262; pl. 22, figs 15-16, 20-21, 26. [groupe de calcaire à *Hippurites*, Sümeg-Kövesdomb, Bakony Mts., Hungary] (Mid-Cretaceous) <HT: GIH Eb/557>.

Genus *Apoxyptalum* MCNAMARA, 1993b

- Apoxyptalum* MCNAMARA, 1993b: 40-42. Type-species: *Apoxyptalum chenjafra* MCNAMARA, 1993b. [Australia] (Oligocene).

Oligocene

- A. chenjafra* MCNAMARA, 1993b: 42-44; figs 2-4. [Waurm Ponds Limestone, Blue Circle Southern Cement Quarry, Reservoir Road, Waurm Ponds, Victoria, Australia] (Janjukian, Late Oligocene) <HT: NMV P135991; PT: NMV P135989, P135990; P135992; P135993; WAM 92.374-382>.

Genus *Brissalius* COPPARD, 2008

- Brissalius* COPPARD, 2008: 3. Type-species: *Brissalius vannaordenburgi* COPPARD, 2008. [Philippines] (Recent) {referred to the family Brissopsidae by COPPARD (2008)}.

Recent

- B. vannaordenburgi* COPPARD, 2008: 3-6; figs 1a-g, 2a-g, 9a-p. [off Siquijor Island, Philippines (depth 200 m)] (Recent) <HT: BMNH 2008.618; PT: BMNH 2008.619>.

Genus *Brissopatagus* COTTEAU, 1863

Recent

- B. relictus* SHIGEL, 1975c: 333-338; figs 1-10, 11a-b, 12, 13a-b, 14-16. [Sagami Bay, Japan Sea, Soyo-Maru Stat. T8 (35°01'0" N, 139°08'0" E), depth 117 m] (Recent) <HT: MMBS Echi 1101>.

Eocene

- B. cotteaudi* LÓPEZ & SILLERO, 2006: 203-204; figs 214a-c. [La Negreta, Tangel, Province Alicante, Spain] (*Schizaster vilanovae* zone, Lutetian, Middle Eocene) <HT: C. Sillero coll'n E-550>.
- B. depressus* LÓPEZ & SILLERO, 2006: 204-205; figs 215, 215A. [Villafranca, Province Alicante, Spain] (Lutetian, Middle Eocene) <HT: C. Sillero coll'n E-554>.

- B. lucentinus* LÓPEZ & SILLERO, 2006: 205-206, figs 216a-d. [Tangel, Province Alicante, Spain] (*Schizaster vilanovae* zone, Lutetian, Middle Eocene) <HT: C. Sillero coll'n E-545>.
- B. palamoensis* LÓPEZ & SILLERO, 2006: 206; figs 217a-d. [Villafranca, Province Alicante, Spain] (*Schizaster vilanovae* zone, Lutetian, Middle Eocene) <HT: C. Sillero coll'n E-543; PT: C. Sillero coll'n E-547>.
- B. parvus* LÓPEZ & SILLERO, 2006: 206-207; figs 218a-b. [Cerro de Garachico, Villafranca, Province Alicante, Spain] (*Schizaster vilanovae* zone, Lutetian, Middle Eocene) <HT: C. Sillero coll'n E-551>.
- B. pulchellus* LÓPEZ & SILLERO, 2006: 207-208; figs 219a-d, 220A-F, un-numbered fig. on p. 207. [Tangel, Province Alicante, and Cantera de los Morteros, Province Aspe, Spain] (*Schizaster vilanovae* zone, Lutetian, Middle Eocene) <HT: C. Sillero coll'n E-549; PT: C. Sillero coll'n E-552, C.López coll'n AM42>.
- B. silleri* SANTOLAYA in SANTOLAYA & SILLERO, 1994: 13-14; two unnumbered figs on p. 14. [Villafranca, Tangel, Province Alicante, Spain] (Lutetian, Middle Eocene) <ST: Collections Sillero, Alicante & Santolaya, Lejona> {re-described in LÓPEZ & SILLERO (2006: 208-209)}.

Genus *Brissopsis* AGASSIZ in AGASSIZ & DESOR, 1847

Miocene

- B. australis* McNAMARA, PHILIP & KRUSE, 1986: 63; figs 7A-E. [Puebla Fm., "Ledge", Torquay, Victoria, Australia] (Longfordian, Early Miocene) <HT: NMV P18557; PT: NMV P18555>.

Genus *Brissus* GRAY, 1825

Pliocene

- B. greifatensis* ELATTAAR, 2001a: 656; fig. 7A; pl. 5: figs 1-4. [Sharm El Arab Member, Shagra Formation, Mersa Um Greifat, S of Wadi Wizr, and Wadi Wizr, 41 km SW Quseir, Red Sea coast, Egypt] (Early-Late Pliocene) <3 syntypes, no numbers mentioned, AUSGM E collection>.
- B. shaimaae* ALI, 1985: 292; figs 4A-C. [Wadi Gabir, Western Egypt] (Early Pliocene) <HT: USNM 339749; PT: USNM 339750>.

Miocene

- B. daviesi* JAIN, 2002: 127-128; pl. 6, figs 4-9. [Gaj Fm., 4.75 km southeast of Rajitpur, Kathiawar, Gujarat, India] (Early Miocene) <HT: GSI 20787; PT: GSI 20784-86>.

B. fosteri MCNAMARA, PHILIP & KRUSE, 1986: 56-59; figs 1-3, 4A. [Mannum Fm., Yarra Glen, South Australia] (Longfordian, Early Miocene) <HT: NMV P55535; PT: NMV P55534, P55536>.

Oligocene

B. bridgeboronensis CARTER, 1987: 1043-1045; figs 2.1-2.3. [Bridgeboro Quarry, Mitchell County, Georgia, USA] (Lower Oligocene) <HT: USNM 402062>.

Genus *Cyclaster* COTTEAU, 1856

Oligocene

C. azemati DEVRIÈS, 1972: 52-54; pl. 1: figs 6-10; pl. 2: figs 9-15. [Horna and Monteagudo, Province Alicante, Spain] (Oligocene) <ST: coll. J. Azema, Paris [no specimen no. provided], coll. J. Leclerc, Paris [no specimen no. provided]>.

Maastrichtian

C. galei JEFFERY, 1997a: 704-705; text-figs 16, 18A-C; pl. 10, figs 5-8. [Koshak and Kyzylsai, Mangyshlak, Kazakhstan] (Uppermost Maastrichtian) <HT: BMNH EE5575; PT: BMNH EE5576, EE5626>.

C. platornatus KUTSCHER, 1978b: 1028-1030; pl. 3, figs 1-5. [Rügen, Germany] (Late Early Maastrichtian) <HT: SGWG 60/2>.

Genus *Eupatagus* L. AGASSIZ, 1847

Subgenus *Eupatagus* L. AGASSIZ, 1847

Recent

E. flindersi BAKER & ROWE, 1990: 291-292; figs 15, 25-27, 91-95. [S. of Goat Id, Victoria, Australia, N. of Eddystone Pt. and Ringarooma Bay, Tasmania] (Recent) <HT: NMNZ 4235; PT: QVM 1985/15/1, 1975/15/2>.

Miocene

E. cetus KRUSE & PHILIP, 1985: 173; fig. 8a-b; pl. 4, figs 4-8. [Mannum Fm., Murray cliffs, Swan Reach to Mannum and Wongulla, South Australia] (Early Miocene) <HT: NMV P18311; PT: NMV P18144, P18142>.

[*E. coronalis* PHILIPPE, 1989]: 32. [Rhône Basin, France] (Miocene) <not given> {no description, figures or repository are given → *nomen nudum*}.

E. ludbrookae KRUSE & PHILIP, 1985: 174; fig. 10a-b; pl. 3, fig. 5; pl. 4, figs 1-3. [Mannum Fm., Murray cliffs, Swan Reach, Mannum and Wongulla; Morgan Limestone, Morgan; all in South Australia] (Early Miocene) <HT: NMV P18241; PT: NMV P55498>.

Oligocene – Miocene

E. collabus KRUSE & PHILIP, 1985: 171; fig. 6a-b; pl. 1, figs 11-13; pl. 2, figs 1-4. [numerous localities in South and West Australia; type from Mannum Fm., Murray cliffs, Swan Reach to Mannum, South Australia] (Late Oligocene – Early Miocene) <HT: NMV P18135; PT: NMV P55484>.

Oligocene

E. planulatus KRUSE & PHILIP, 1985: 175; fig. 11a-b; pl. 4, figs 9-11. [Port Willunga Fm., Aldinga and Port Vincent Limestone, Surveyor Point, Yorke Peninsula, both South Australia] (? Oligocene) <HT: NMV P20492; PT: AUGD S23(3)>.

E. rostratus zitteli HENDERSON, 1975: 45-46; text-figs 10a-e; pl. 11, figs 5-6; pl. 12, figs 1-3. [Cape Farwell, west Nelson; and many other localities, New Zealand] (Whaingaroan – Altonian, Oligocene to Early Miocene) <LT: NHMW 1970/1336; PLT: NHMW 1959/335/50, 1970/1337> {*nomen novum* for *Hemipatagus formosus* ZITTEL, 1864 non DE LORIO, 1863}.

E. (E.) singhi SRIVASTAVA, 1981: 39-40; pl. 1, figs 5-8. [near Ber Mota village, Kutch, India] (*Lepidocyclina (Eulepidina)* zone, Middle Oligocene) <HT: DGUL KTE 1021; PT: DGUL KTE 1022, KTE 1023>.

E. (E.) tandoni SRIVASTAVA, 1981: 40-41; pl. 1, figs 9-11. [near Ber Mota village, Kutch, India] (*Eupatagus (E.) rostratus* zone, Late Oligocene) <HT: DGUL KTE 1030; PT: DGUL KTE 1031-34> {transferred to subgenus *Gymnopatagus* by SRIVASTAVA (1988: 152)}.

Eocene

E. batequensis SQUIRES & DEMETRION, 1992: 46-48; figs 142-145. [Bateque Fm., CSUN-locality 1470, Arroyo San Juan de Abajo, Baja California Sur, Mexico] (Eocene) <HT: IGM 5923 (= LACMIP 8863); PT: IGM 5924 (= LACMIP 8864)>.

E. lawsonae KIER, 1980: 53; pl. 21, figs 1-4. [Castle Hayne Limestone, Ideal Cement Company quarry localities 12, North Carolina, USA] (Middle Eocene) <HT: USNM 264086>.

E. (E.) rajasthanensis SRIVASTAVA & SINGH, 2008: 85-87; figs 2a-g. [Khuiala Formation, about 625 m west of Habur, Jaisalmer district, Rajasthan, India] (Early Eocene) <HT: BSIP 7097/BSIP/39528; PT: BSIP 7097/BSIP/39529 to 7097/BSIP/39547>.

E. texanus ZACHOS in ZACHOS & MOULINEUX, 2003: 504; figs 7.1-7.3. [Weches Formation, Claiborne Group, Localities 24, 30, 42, 57, 62, 73, 75, 77, 80; Houston, Leon, Nacog-

doches, Sabine, and San Augustine Counties, Texas, USA] (Middle Eocene) <HT: TMM 1987TX1; PT: TMM 1987TX2, 1987TX3, 2013TX1 (all from Locality 42)>.

E. wilsoni KIER, 1980: 51-53; figs 25-26; pl. 20, figs 7-11. [Castle Hayne Limestone, Ideal Cement Company quarry localities 12, 16, North Carolina, USA] (Middle Eocene) <HT: USNM 264084; PT: USNM 264085>.

Subgenus *Gymnopatagus* DÖDERLEIN, 1901

Recent

G. parvipetalus BAKER & ROWE, 1990: 288-291; figs 10-13. [N.E. of Cape Brett, New Zealand, 596-604 m] (Recent) <HT: NZOI H452>.

Genus *Fernandezaster* SÁNCHEZ ROIG, 1952

Miocene

F. durhami FISCHER, 1985: 205-206; figs 5/9, 6a-c; pl. 3, fig. 4. [Turrúcares Fm., Pendiente, SW del Cerro Candelaria, Costa Rica] (Middle Miocene) <HT: ECG CO-97>.

Genus *Gualtieria* DESOR, in AGASSIZ & DESOR, 1847

Eocene-Oligocene

G. alicantina SANTOLAYA, 1993: 44-46; figs 1-8. [Villajoysa, Agost and La Romana, Alicante, Spain] (Eocene-Oligocene) <no types defined, repository unknown>.

Subgenus *Blaviaster* LAMBERT, 1920

Eocene

G. (B.) grossouveri productus CHAVANON, 1974: Vol.1: 269; pl. 18: figs 2a-d. [Blayais and Couquèques, Aquitaine Basin, western France] (Middle to Late Eocene) <ST: Castex colln, Fabre colln; no repository nos given for types; figured specimens: 33.C.31, 33.C.32>.

Genus *Hysteraster* McNAMARA & BARRIE, 1993

Hysteraster McNAMARA & BARRIE, 1993: 140. Type-species: *Hysteraster paragrapsimus* McNAMARA & BARRIE, 1993. [South Australia] (Miocene).

Miocene

H. paragrapsimus MCNAMARA & BARRIE, 1993: 140-145; figs 1-4. [Morgan Limestone, Murray River cliffs, downstream from Waikerie at Broken Cliffs, South Australia] (Early to early Middle Miocene) <HT: SAM P32322; PT: SAM P32323, P24260; MV 18039>.

Genus *Macropneustes* L. AGASSIZ, 1847

Subgenus *Deakia* PAVAY, 1875

Palaeogene

M. (D.) xinjiangensis YANG SHENGQIU, 1991: 125-126; pl. 19, figs 1-5. [Upper Qimugen Fm., Kuzigongsu, 5 km E Bashibulake, Wuqia County, Tarim Basin, China] (Early Tertiary) <HT: NIGP 88439>.

Genus *Meoma* GRAY, 1851

Subgenus *Schizobrissus* POMEL, 1869

Eocene

[*M. (S.) chiplonkari* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <repository unknown> {nomen nudum}.

[*M. (S.) cooperi* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <repository unknown > {nomen nudum}.

[*M. (S.) misrai* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <repository unknown> {nomen nudum}.

[*M. (S.) singhi* SRIVASTAVA, 1988]: 151. [India] (*Asterocyclina alticostata* Zone, Middle Eocene) <repository unknown > {nomen nudum}.

Genus *Plagiobrissus* POMEL, 1883

(pro *Plagionotus* AGASSIZ & DESOR, 1847 non Mulsant, 1842)

Pliocene

Plagionotus holmesii McCrady in Tuomey & Holmes, 1855: 9-10; pl. 3: figs 2, 2a. [Smith's, Goose Creek, South Carolina, USA] (Pliocene) <ST: Museum, College of Charleston, South Carolina, [specimen no. not given]> {based on test fragments}.

Plagionotus ravenelianus McCRADY in TUOMEY & HOLMES, 1855: 10-11; pl. 3: figs 3, 3a. [The Groove, Cooper River, South Carolina, USA] (Pliocene) <ST: Museum, College of Charleston, South Carolina, [specimen no. not given]> {based on test fragments}.

Genus *Rhynobrissus* AGASSIZ, 1872

Recent

R. tumulus MCNAMARA, 1982a: 351-358, figs 1-5. [20°40' S, 115°26' E, beach E of Cape Dupuy, NE coast of Barrow Island, Western Australia] (Recent) <HT: WAM 1047-81; PT: WAM 1048-81, 1049-81, 1050-81, 1051-81, BMNH 1981.10.27.1, AM J14770-J14772>.

Genus *Taimanawa* HENDERSON & FELL, 1969

Oligocene

T. rostrata HENDERSON, 1975: 50-51; pl. 14, figs 1-3. [Hazelburn, South Canterbury, New Zealand] (Duntroonian – Waitakian, Oligocene) <HT: NZGS EC.389>.

Eocene

T. prisca HENDERSON, 1975: 49-50; pl. 13, figs 4-6; pl. 14, fig. 6. [Gentle Annie Rock and Perpendicular Point, Punakaikai District, Westland, New Zealand] (Kaiatan – ? Runangan, Late Eocene) <HT: NZGS EC.587; PT: NZGS EC.392, 576, 583-4, 586, 588, 592-9>.

Family Spatangidae GRAY, 1825

Genus *Maretia* GRAY, 1855

Pliocene

M. planulata abbassi ALI, 1985: 294-295; figs 12A-B. [Wadi Abu Abraiki, Western Egypt] (Early Pliocene) <HT: USNM 339748>.

Miocene

M. ranjitpurensis JAIN, 2002: 130-132; pl. 6, figs 11-14. [Gaj Fm.; 4.75 km southeast of Rajitpur; Kuranga Railway Station; and 2.5 km west-southwest of Bhatvadia, all in Kathiawar, Gujarat, India] (Early Miocene) <HT: GSI 20791; PT: GSI 20790, 20792-93>.

Oligocene

M. carolinensis KIER, 1997: 11-13; fig. 6; pl. 9, figs 8-9; pl. 10, figs 1-7; pl. 11, figs 1-2. [Trent Fm., Polocksville state quarry, North Carolina, USA] (Late Oligocene) <HT: USNM 398338; PT: USNM 398337, 398339-398341, 492101-492124>.

Genus *Metalia* GRAY, 1855

Recent

M. angustus DE RIDDER, 1984: 621-623; pl. 2, figs A-D. [Nouvelle-Calédonie] (Recent) <MNHN ECES 8286; PT: MNHM ECES 8287>.

M. kermadecensis BAKER & ROWE, 1990: 293-295; figs 34-35. [Raoul Id., Kermadecs, 90-120 m] (Recent) <HT: NMNZ 3425>.

M. sternalis var. *lata* DOLLFUS & ROMAN, 1981: 124. [Red Sea] (Recent) <HT: not given>.

M. sternalis var. *jousseaumei* DOLLFUS & ROMAN, 1981: 124; pl. 41, figs 2-5. [Red Sea] (Recent) <HT: EM, specimen no. not given>.

Miocene

M. harshadae JAIN, 2002: 128-130; pl. 6, fig. 10; pl. 7, fig. 1. [Gaj Fm., 4.75 km southeast of Rajitpur, Kathiawar, Gujarat, India] (Early Miocene) <ST: GSI 20788-89>.

Genus *Megapneustes* GAUTHIER in FOURTAU, 1899

Eocene

M. jaisalmerensis SRIVASTAVA, RANA & SINGH, 2008: 32-35; pl. 1: figs 1-3, 5-8. [Khuiala Formation, Gharollia Hill section, situated about 2 km northwest of village Pariwar (= Parivar), Jaisalmer district, Rajasthan, India (27°15' N, 70°45' E)] (Early Eocene) <HT: GU/R/KE/2023; PT: GU/R/KE /2024 to GU/R/KE/2026 and GU/R/KE /F – 2027 to GU/R/KE /F to 2031>.

Genus *Peraspatangus* PHILIP & FOSTER, 1971

Peraspatangus PHILIP & FOSTER, 1971: 689-690. Type-species: *Peraspatangus brevis* PHILIP & FOSTER, 1971. [Australia] (Miocene) {other species included: *P. depressus* PHILIP & FOSTER, 1971}.

Miocene

- P. brevis* PHILIP & FOSTER, 1971: 690-692; text-figs 9b-d; pl. 126: fig. 3; pl. 133: fig. 1, 2, 4-6, 8. [Bochara Limestone, junction Grange Burn – Muddy Creek, Hamilton district, Victoria, Australia] (Batesfordian, Lower Miocene) <HT: NMV P27956; PT: NMV P27955, UNE 11758>.
- P. depressus* PHILIP & FOSTER, 1971: 692-693; text-figs 9a; pl. 128: fig. 2; pl. 132: fig. 1, 4, 5, 7; pl. 133: fig. 3. [Rutledge Marl Mb., Port Campbell Limestone, Ingle's Creek, Port Campbell, Victoria, Australia] (Bairnsdalian, Middle Miocene) <HT: NMV P27953; PT: NMV P27954>.

Genus *Spatangobrissus* CLARK, 1923

Attributed to the family Maretiidae by HOLMES, AH YEE & KRAUSE (2005).

Recent

- S. incus* BAKER & ROWE, 1990: 295-297; figs 17, 36-41. [E. of Flinders Island, Bass Strait, between Wilson's Promontory and Glennie Island, Victoria and S. of Bemm River, Victoria, Australia] (Recent) <HT: TM: H515; PT: MV F52356, F52357>.

Miocene

- S. dermodyorum* HOLMES, AH YEE & KRAUSE, 2005: 94-95; figs 6A-E, 7A-C, 8A-E. [Glenforslan Fm., Morgan Grp., 7 km NNE Murray River Lock 1, Blanchetown, South Australia] (Batesfordian, Middle Miocene) <HT: NMV P312570; PT: NMV P312571-P312573>.

Genus *Spatangus* GRAY, 1825

Recent

- S. mathesoni* MCKNIGHT, 1968: 102-107; figs 10-11. [Stat. D 221 (40°06' S, 171°16' E, depth 686 m), D 224 (40°47' S, 169°41' E, depth 903 m), D 226 (39°54' S, 168°40' E, depth 823 m), D 231 (37°53' S, 169°45' E, depth 772 m), Tasman Sea] (Recent) <HT: NZOI 27 (Stat. D 226); PT: NZOI P 42, P 43 (Stat. D 231)>.

Subgenus *Phymapatagus* LAMBERT, 1910

Eocene

S. (P.) grandituberculatus LEWIS, 1989: 34-39; figs 11, 12 a-c; pl. 7: figs 1 a-e, 2 a-c. [Barton-on-Sea, Hampshire, England] (Bartonian) <HT: BMNH E 76821; PT: E 37320-3, E 76817-20, E 76459>.

Genus *Trachypatagus* POMEL, 1869

Miocene

T. nehaleae ALI, 1989b: 60; fig. 8a-d. [Marmarica Limestone, Wadi El Habis, W of Mersa Matruh, Egypt] (Langhian-Serravallian, Middle Miocene) <HT: MUGM 8619; PT: MUGM 8620>.

Subfamily Eurypataginae KROH, 2007

Eurypataginae KROH, 2007: 174. Type-genus: *Eurypatagus* MORTENSEN, 1948. [Indo-Pacific Ocean] (Miocene to Recent) {other genera included: *Paramaretia* MORTENSEN, 1950, ? *Platybrissus* GRUBE, 1866}.

Family Loveniidae LAMBERT, 1905

Genus *Araeolampas* SERAFY, 1974

Araeolampas SERAFY, 1974: 43. Type-species: *Homolampas fulva* A. AGASSIZ, 1879 [Northern Atlantic] (Recent) {other species included: *Araeolampas atlantica* SERAFY, 1974}.

Recent

A. atlantica SERAFY, 1974: 44-46; figs 1a-c, 2a-b. [Albatross Stat. 2105 off Virginia (37°50' N, 43°03.8' W, depth 2575 m); Atlantis Stat. 20 off the Azores (37°50.5' N, 26°00' W, depth 2585 m); west of Dry Tortugas, Gulf of Mexico (depth 3545 m); Pillsbury Stat. P-120 west of Georgia (31°48' N, 76°38' W, depth 1920 m), Stat. P-292, west of Gabon, Gulf of Guinea (0°12' N, 5°11' E, depth 3595 m), Stat. P-1429, north of Haiti (21°19.2' N, 73°45.5' W, depth 2532 m); all in the Atlantic Ocean] (Recent) <HT: USNM 24598 (from Albatross Stat. 2105)>.

Genus *Bahariya* ELBASSYONY, 2005

Bahariya ELBASSYONY, 2005: 321-324. Type-species: *Bahariya teetotumensi* ELBASSYONY, 2005. [Egypt] (Eocene) {considered a junior subjective synonym of *Gualtieria* L. AGASSIZ, 1847 by Smith, The Echinoid Directory (accessed 20.I.2009)}.

Eocene

B. teetotumensi ELBASSYONY, 2005: 324-326; pl. 1: figs 1a-d, 2-6; pl. 2: figs 1a-c, 2. [Mokattam Formation, Teetotum hill, E of Qazzun sand dunes, Teetotum section, NE plateau of Bahariya Oasis, Egypt] (Bartonian, Middle Eocene) <HT: specimen figured in pl. 1: figs 1a-; PT: 20 specimens (type material deposited at: ASU GM under 019-001 to 019-007, Ahmed EL-BASSYONY in litt., 23.04.2007)>.

Genus *Breynia* DESOR, 1847

Recent

B. neanika McNAMARA, 1982b: 188-192; figs 12-13. [Arafura Sea; Albany Passage, Torres Strait; Bowen, Queensland, Northern Australia] <HT: AM J14324; PT: AM J14325, AM J14326a-d>.

Genus *Echinocardium* GRAY, 1825

(Genus *Amphidetus* L. AGASSIZ, 1836, objective junior synonym of *Echinocardium* GRAY, 1825)

Recent

E. meteorensis MIRONOV, 2006: 120-125; figs 11A-I, 12A-I, 13A, 14A-C. [NE Atlantic Seamounts (Atlantis, Great Meteor, Seine); R/V “Moskovsky Universitet” Stat. 19-2, Seine Seamount, NE Atlantic Ocean (33°45' N, 14°20' W, depth 180-200 m); R/V “Akademik Petrovsky”, Cruise 12, Stat. 36; R/V “Vityaz 2”, Cruise 2, Stat. 154B; R/V “Meteor”, Stat. 161a-KD62; R/V “Le Suroit”, expedition “Seamount 2”, Stat. DW 147, Stat. DW 189; R/V “Ikhtiandr”, Cruise 8, Stat. 21A, unnamed seamount (11°37'5 S, 05°11'9 W, depth 300-450 m)] (Recent) <HT: IORAS XV-69-33 (from “Moskovsky Universitet” Stat. 19-2); PT: 2 specimens, no numbers given (from “Meteor” Stat. 161a-KD62 and “Akademik Petrovsky” Stat. 36 respectively)>.

Pliocene

A. ampliflorus McCrady in Tuomey & Holmes, 1855: 6-7; pl. 2: figs 2, 2a. [The Groove, Cooper River, South Carolina, USA] (Pliocene) <not given>.

E. kelloggi KIER, 1983: 500-501; fig. 1; pl. 1, figs 1-5. [Yorktown Fm., Lee Creek Mine, North Carolina, USA] (Lower Pliocene) <HT: USNM 186521>.

Miocene

E. biaense MIHÁLY, 1985: 245-246, 262; pl. 5, figs 3-5. [Leitha Limestone, Bia, Alsó-mjor hegyoldal; Rákos District, Budapest; and Örs vezér tere site, all in Hungary] (Late Badenian (= Early Serravallian), Middle Miocene) <HT: MAFI Ech 115; PT: MAFI Ech 129, 427, 428> {according to RADWAŃSKI & WYSOCKA (2001: 307) this species is a synonym of *Echinocardium peroni* COTTEAU, 1877}.

E. leopolitanum RADWAŃSKI & WYSOCKA, 2001: 304-306; pl. 3: figs 1a-c, 2a-c, 3a-c, 4a-c; pl. 4: figs 1a-c, 2-5, 6a, b; ül. 5: figs 1a-c, 2a, b, 3a, b; pl. 6: figs 1a-d, 2a-c, 3. [Gleboviti (= Chlebowice), near Bibrka, c. 30 km SE Lviv, Ukraine] (Early Badenian [= Langhian], Middle Miocene) <HT: Radwański, no repository no. given [pl. 6: fig. 1a-d]; PT: 2 specimens, Radwański, no repository no. given [pl. 6: fig. 2a-c, 3]>.

E. marylandiense KIER, 1972b: 16-18; fig. 6; pl. 7, figs 2-3; pl. 8, figs 1-2. [Choptank Fm., Scientists' Cliffs, Kenwood Beach, Calvert County, Maryland, USA] (Middle Miocene) <HT: USNM 174460>.

E. olisiponensis KOTCHETOFF, KOTCHETOFF & VEIGA FERREIRA, 1975: 67-75; pl. 3, figs 1-2; pl. 4, figs 3-4; pl. 5, figs 5-9. [Penedo cliff, north of Cap d'Espichel, Portugal] ("Helvetian", Miocene) <not given>.

Genus *Lovenia* DESOR, 1847

Recent

L. lata SHIGEL, 1981: 81-87; figs 1-9. [Aoyama-dashi, Sagami Bay, Japan] (Recent) <HT & PT: MMBS> {no specimen nos. given}.

Miocene

L. bagheerae IRWIN in IRWIN & ARCHBOLD, 1994: 10-12; figs 10A-L. [Port Campbell Limestone, Portland, Victoria, Australia] (Mitchellian, Tortonian, early Late Miocene) <HT: NMV P79247; PT: NMV P79233-P79270, P78918, P78927, P78935, P78941, P78943, P78953>.

Incertis sedis

Genus *Murraypneustes* HOLMES, AH YEE & KRAUSE, 2005

Murraypneustes HOLMES, AH YEE & KRAUSE, 2005: 92-93. Type-species: *Murraypneustes biannulatus* HOLMES, AH YEE & KRAUSE, 2005. [South Australia] (Miocene).

Miocene

M. biannulatus HOLMES, AH YEE & KRAUSE, 2005: 93-94; figs 2A-F, 3A-C, 4A-B, 5. [Glenforslan Fm., Morgan Grp., 7 km NNE Murray River Lock 1, Blanchetown, South Australia] (Batesfordian, Middle Miocene) <HT: NMV P312370; PT: NMV P312371, P312372>.

Suborder Asterostomatina FISCHER, 1966

Family Asterostomatidae PICTET, 1857

Genus *Acanthotrema* BAKER & ROWE, 1990

Acanthotrema BAKER & ROWE, 1990: 312. Type-species: *Acanthotrema siculum* BAKER & ROWE, 1990. [Subantarctic Islands] (Recent) {BAKER (1998: 285) proposed the name *Kermabrissoides* for this genus, since the name *Acanthotrema* is preoccupied}.

Recent

A. siculum BAKER & ROWE, 1990: 312-314; figs 22-24, 84-87. [off Raoul Island, Kermadecs and off Norfolk Island] (Recent) <HT: NMNZ 4254; PT: NZOI P689>.

Genus *Antillaster* LAMBERT, 1909

Eocene

A. albeari KIER, 1984b: 135; pl. 80, figs 1-2; pl. 81, figs 1-2. [Palmer loc. 1003, N of Carretera Central, 3.5 km, on road to San Diego de los Baños, Pinar del Rio Province, Cuba] (Middle to Late Eocene) <HT: ANSP 16631>.

A. bagmanovi McNAMARA & MELIKOV, 2002: 160-164; figs 2, 3a-b, 4, 5a-b, 6, 7. [Paradah Group, Daralik, Nakhichevan region, Azerbaijan] (Late Lutetian to Bartonian, Eocene) <HT: WAM 99.428; PT: 99.429, 99.432>.

Genus *Asterostoma* L. AGASSIZ, 1847

Eocene

A. pawsoni KIER, 1984b: 128-129; fig. 40; pl. 78, figs 1-6. [hillsides above the Yallahs River, St. Thomas, in the Lucky Hill region in St. Mary, and in the Spring Mount region, St. James, Jamaica] (Eocene) <HT: USNM 301378a; PT: USNM 301378b, 301378c>

Genus *Heterobrissus* MANZONI & MAZZETTI, 1877

Recent

H. erinaceus BAKER & ROWE, 1990: 307-309; figs 19, 73-77, 109-111. [off Cape Hawke, E. of Brush Island, E. of Sydney, E. of Sugarloaf Point, S.E. of Jervis Bay, New South Wales and E. of Peregian Beach, southern Queensland, Australia] (Recent) <HT: AM J11032; PT: AM J14450, J14452-8, J11082, J10823, J14460, J5819, J14461; NMNZ 6153, 5104, 1995>.

H. gigas BAKER & ROWE, 1990: 309-310; figs 19, 78-80, 112-114. [E. of Brush Island, New South Wales, off Cape Everard, Victoria, E. of Kiama, off Broken Bay, off Sydney, off Collaroy, off Cape Hawke and between Sydney and Port Stephens, New South Wales, Australia] (Recent) <HT: AM J10824; PT: AM J6383, J10819-J10820, J10822, J10825, J10871, J11031, J12432, J14466, J14470, J14472-J14474; NMNZ 3921-3922>.

Genus *Kermabrissoides* BAKER, 1998

Kermabrissoides BAKER, 1998: 285. Type-species: *A. siculum* BAKER & ROWE, 1990. {nomen nov. pro *Acanthotrema* BAKER & ROWE, 1990 non TAVASSOS, 1928}.

Genus *Palaeotropus* LOVEN, 1874

Recent

P. uniporum MIRONOV, 2006: 117-119; figs 8A-D, 9A-G. [R/V "Akademik Mstislav Keldysh" Stat. 521 Atlantis Seamount, NE Atlantic Ocean (34°23'5-34°23'5 N, 29°59'6-29°59'1 W, depth 1560-1900 m)] (Recent) <HT: IORAS XV-69-32>.

Genus *Plesiozonus* DE MEIJERE, 1902

Recent

P. tenuis DAVID & DE RIDDER, 1989: 210-213; figs 4-6; pl. 3, figs 1-6. [Musorstom 2 st. 52; 14° 00,7' N, 120° 18,7' E; Philippines] (Recent) <HT: MNHN> {no specimen no. given}.

Order Neolampadoida PHILIP, 1963

Family Neolampadidae LAMBERT, 1918

Genus *Actapericulum* HOLMES, 1995

Actapericulum HOLMES, 1995: 120. Type-species: *Actapericulum bicarinatum* HOLMES, 1995. [South Australia] (Miocene).

Miocene

A. bicarinatum HOLMES, 1995: 120-124; figs 6A-M, 8I-L. [Zeally Limestone Mb., Puebla Fm., Point Danger, Torquay, Victoria; Gambier Limestone, at Mt. Gambier and near Nelson; Mannum Fm, near Bow Hill, South Australia] (Longfordian – ? Batesfordian, Burdigalian – Langhian, late Early Miocene) <HT: NMV P140924; PT: NMV P73689-P73703, P73710-P73724, P140925, P140926, P140929, P73683, P73684-P73688, P73704-P73709>.

Genus *Aphanophora* DE MEIJERE, 1903

Eocene

A. ? bassoris HOLMES, 1995: 116-119; figs 2A-B, 3A-M, 5I-L. [Kingscote Limestone, Kingscote, Kangaroo Island, South Australia] (Aldingan, Bartonian?-Priabonian, Late Eocene) <HT: NMV P140922; PT: NMV P73675-P73682, P133073, P140923>.

Incertis sedis**Genus *Desorella* COTTEAU, 1855**

Late Jurassic – Early Cretaceous

D. moravica BLASCHKE, 1911: 180-181; pl. 6: figs 11a, b. [whitish variant of Stramberk Limestone, Štramberk (= Stramberg), Czech Republic] (Tithonian to Berriasian,) <HT: NHMW 1910/0010/0036>.

Genus *Mizunamia* ISHIJIMA & HATAI, 1973

Mizunamia ISHIJIMA & HATAI, 1973: 67. Type-species: *Mizunamia tokiensis* ISHIJIMA & HATAI, 1973. [Japan] (Miocene).

Miocene

M. tokiensis ISHIJIMA & HATAI, 1973: 67 (description: on p. 65); fig. 1. [Akeyo Formation, Mizunami Group, Obora, west of Akeyo and Northeast of Toki, Toki-Mizunami Dis-

trict, Gifu Prefecture, Central Japan] (Miocene) <no types defined, material housed in the collection of Kotora Hatai> {problematic microfossil known from single cross section only, resembles echinoid spine cross section}.

Echinoid Traces (formally named ichnotaxa)

Genus *Bichordites* PLAZIAT & MAHMOUDI, 1988

Bichordites PLAZIAT & MAHMOUDI, 1988: 223-224. Types-species: *Bichordites monastiriensis* PLAZIAT & MAHMOUDI, 1988. [Tunisia] (Pleistocene). {echinoid burrows}.

Pleistocene

B. monastiriensis PLAZIAT & MAHMOUDI, 1988: 224-225; text-figs 5a-c, 8a-e; pl. 1: figs A1-A3, B1-B3, C-F; pl. 2: fig. B. [Khnis Quartzose, abandoned quarry at Khnis, 7 km SW of Monastir, S of Hergla, East-Central Tunisia] (Thyrrhenian, Late Pleistocene) <HT: UPS P.2201; PT: UPS P.2202-2216>.

Genus *Cardioichnus* SMITH & CRIMES, 1983

Cardioichnus SMITH & CRIMES, 1983: 90. Type-species: *Cardioichnus planus* SMITH & CRIMES, 1983. [Europe] (Cretaceous-Miocene). {echinoid burrows}.

Miocene

C. reniformis MAYORAL & MUÑIZ, 2001: 71-75; figs 2A-D, 3A-C, 4A1-A2. [Lepe, SW Huelva Province, Guadalquivir Basin, Spain] (Late Miocene) <HT: GMUS LE11/Cr1 (figs 3A-C); PT: GMUS LE11/Cr2 (figs 4A1-A2)>.

Eocene

C. foradadaensis PLAZIAT & MAHMOUDI, 1988: 225; text-figs 12d, 12f, 12 h; pl. 2: figs C, D. [Foradada-de-Toscar, Huesca, Northern Spain] (Late Ilerdian-Early Cuisian, Early Eocene) <HT: UPS 2217>.

C. ovalis SMITH & CRIMES, 1983: 91; fig. 7B. [exact locality unknown, near Czarna, Wiselka, Silesia, Poland] (Cretaceous-Eocene) <HT: BMNH T762>.

C. planus SMITH & CRIMES, 1983: 90-91; figs 7A, C. [Zollhaus Quarry, ca. 25 km W of Lake Thun, Prealps, Switzerland] (Ilerdian, Early Eocene) <HT: BMNH T763 (middle of three traces); PT: BMNH T763 (outer two traces)>.

Genus *Ereipichnus* MONACO, GIANNETTI, CARACUEL & YEBENES, 2005

Ereipichnus MONACO, GIANNETTI, CARACUEL & YEBENES, 2005: 344. Type-species: *Ereipichnus geladensis* MONACO, GIANNETTI, CARACUEL & YEBENES, 2005. [Spain] (Albian) {echinoid burrows}.

Cretaceous

E. geladensis MONACO, GIANNETTI, CARACUEL & YEBENES, 2005: 344; figs 3A-D, 4A-G. [Sa'caras Formation, Serra Gelada section (Prebetic of Alicante), Spain] (Early to Middle Albian) <HT: EGSG T109; PT: EGSG T110 to T112> {Ichnofossil, horizontal meandering burrows with backfill structure}.

Genus *Gnathichnus* BROMLEY, 1975

Gnathichnus BROMLEY, 1975: 738. Type-species: *Gnathichnus pentax* BROMLEY, 1975. [worldwide] (Jurassic to Recent). {echinoid grazing traces}.

Pleistocene

G. pentax BROMLEY, 1975: 738; pl. 85, figs 1-3; pl. 86, figs 3-5; pl. 87, figs 1-7; pl. 88, figs 1-7. [type-loc.: Sgourou Fm., Kritika, Rhodes, Greece – range: worldwide] (type-strat.: lowermost Pleistocene – range: Jurassic – Recent) <HT: MMH 13386; PT: IGS 115027, 115029> {echinoid grazing traces on bivalves, belemnites, echinoids and pebbles}.

Cretaceous

G. stellarum BRETON, NÉRAUDEAU & CUENCA-BOULAT, 1992: 220-223; pl. 88, figs 1-5. [Falaise du Caillard, Tulmont, Charentes-Maritimes, France] (Upper Campanian) <HT: MHNN 8562; PT: MHNN 8563-8566> {echinoid grazing traces on asteroid marginalia of *Metopaster*; junior synonym of *G. pentax* according to RADWANSKA, 1999: 356}.

Genus *Roderosignus* MICHALÍK, 1977

Roderosignus MICHALÍK, 1977: 340. Type-species: *Roderosignus quinqueradialis* MICHALÍK, 1977 [Slovak Republic] (Late Triassic). {echinoid grazing traces}.

Triassic

R. quinqueradialis MICHALÍK, 1977: 340-342; fig. 14.1-3. [Kössen Fm., Hybe, Low Tatra Hills, Slovak Republic] (Rhaetian, Late Triassic) <HT: SNM XXXX> {echinoid graz-

ing traces on the brachiopod *Rhaetina pyriformis*; junior synonym of *G. pentax* according to RADWANSKA, 1999: 356}.

Formally named Echinoid Growth Anomalies

forma aegra *disparradiata* KRÜGER, 2002: 176; figs 1-3; pl. 1. {deviation from pentamery (e.g. specimens with 3, 4, 6, or 7 ambulacral columns); documented for *Galerites* (*G.*) *vulgaris* (LESKE, 1778) from Northern Europe; KRÜGER (2006: tab. 1) provided a list with numerous additional examples}.

forma aegra *inconstans* KRÜGER, 1993: 317; figs 2a-c, 5a-f, 11, 12, 13. {abnorm insertion of additional plates in ambulacral or interambulacral columns; documented for *Echinocorys gibbus oviformis*, *E. humilis*, *E. ovata*, *E. subglobosa*, *E. sulcata*, *E. cf. limburgica*, *Holaster subglobosus* from the Late Cretaceous of Northern Europe}.

forma aegra *insecta* KRÜGER, 2006: 59-60; figs 1-7. {abnorm pinching/constriction of the two poriferous zones in an ambulacral column; documented for *Arbacia aequituberculata*, *Echinolampas beaumonti*, *E. kleini*, *E. ovalis*, *Echinometra lucunter*, *Echinoneus cyclostomus*, *Encope grandis*, *Hemiaster batnensis*, *Lytechinus veriegatus*, *Micraster* (*Gibbaster*) *gibbus*, *M. stolleyi*, *Paracentrotus lividus*, *Phymechinus mirabilis*, *Spatangus purpureus*, and *Sphaerechinus granularis*}.

References

- ABDELHAMID, M.A.M. & AZAB, M.M. (2003): Aptian-Cenomanian Echinoids from Egypt. – *Revue de Paléobiologie*, **22**/2: 851-876.
- ADEGOKE, O.S. (1977): Stratigraphy and paleontology of the Ewekoro Formation (Paleocene) of southwestern Nigeria. – *Bulletin of American Paleontology*, **71**/295: 5-379.
- AISENBERG, D.E. (1974): [A new representative of the genus *Archaeocidaris* M'COY from the Middle Carboniferous of the Donetz Basin]. – *Vikopni Fauna Flora Ukraine*, **2**: 76-79.
- ALI, M.S.M. (1983): A new *Echinolampas* species (Cassiduloida: Echinoidea) from the Middle Eocene of Egypt. – *Journal of the Palaeontological Society of India*, **28**: 56-58.
- (1985): On some Pliocene echinoids from the area north of Mersa Alam, Red Sea coast, Egypt. – *Paläontologische Zeitschrift*, **59**/3-4: 277-300.
- (1989a): Late Cretaceous echinoids from Gebel El Rowdah, Hatta area, U.A.E. – *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, **177**/3: 393-412.
- (1989b): Middle Miocene *Trachypatagus* (Echinodermata, Echinoidea), from Egypt. – *Journal of African Earth Sciences*, **8**/1: 57-63.

- (1990): Cenomanian echinoids from Ras al Khaimah, United Arab Emirates. – *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, **179/1**: 97-116.
 - (1992a): Additional echinoids from the Late Maestrichtian (Cretaceous) of Gebel el Rowdah, Oman — United Arab Emirates. – *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, **1992/2**: 65-74.
 - (1992b): A new genus and species of regular echinoid from the late cretaceous of Gebel el Rowdah, Hatta area, Oman – U.A.E. – *Journal of Paleontology*, **66/4**: 693-692.
- AMARD, B., COLLIGNON, M., & ROMAN, J. (1983): Etude stratigraphique et paleontologique du Cretace superieur et Paleocene du Tinrhert-W et Tademaït-E (Sahara Algerien). – *Documents des Laboratoires de Géologie de la Faculte des Sciences de Lyon Hors Série*, **6 (1981)**: 15-173.
- AZAB, M.M. (1989): A new *Ditremaster* species (Spatangoida: Echinoidea) from the Middle Eocene of Egypt. – *Middle East Research Center Ain Shams University, Earth Science Series*, **3**: 179-183.
- & ELATTAAR, A.A.E.H.A. (1990): *Pseudopygaulus romani* (Cassiduloidea, Echinoidea): a new species from the Upper Libyan (Lower Eocene) rocks of Egypt. – *Middle East Research Center Ain Shams University, Earth Science Series*, **4**: 215-221.
 - & ELATTAAR, A.A.E.H.A. (1999): Paleocene/Lower Eocene Echinoids of Egypt. – In: *International Conference on Geology of the Arab World (GAW4)*, Cairo University, Egypt, 21.-25 Feb. 1998. Volume 2. – pp. 837-872, Cairo (Cairo University).
- AZIZ, S.A. (1991): Some echinoid remains from Dalmiapuram Formation and Kallakudi reefoidal limestone (Lower Cretaceous), Trichinopoly subbasin, south India. – *ONGC (Oil & Natural Gas Commission) Bulletin*, **25/2**: 11-30.
- & BADVE, R.M. (1990): Occurrence of *Hemipneustes* AGASSIZ (echinoid) from Tiruchirappalli Cretaceous, Tamil Nadu, India. – *Journal of the Geological Society of India*, **36/3**: 326-330.
 - & BADVE, R.M. (2001): New echinoid fauna fom the South Indian Cretaceous deposits. – *Gondwana Geological Magazine*, **16/2**: 43-59.
- BADVE, R.M. & AZIZ, A.S. (1983): Revision of cassiduloid genus *Gongrochanus* KIER (Echinodermata: Echinoidea) from south Indian Cretaceous deposits. – *Proceedings of the Indian Academy of Sciences Animal Sciences*, **92/3**: 231-246.
- BAKER, A.N. (1983): A new apatopygid echinoid genus from New Zealand (Echinodermata: Cassiduloidea). – *National Museum of New Zealand Records*, **2/15**: 163-173.
- (1998): The rediscovery of *Echinus elevatus* HUTTON 1872 in New Zealand, and a new name for *Acanthotrema* BAKER & ROWE 1990 (Echinodermata: Echinoidea). – *Journal of the Royal Society of New Zealand*, **28/2**: 281-286.
 - & Rowe, F.W.E. (1990): Atelostomatid sea urchins from Australian and New Zealand waters (Echinoidea: Cassiduloidea, Holasteroidea, Spatangoida, Neoplampadoidea). – *Invertebrate Taxonomy*, **4/2**: 281-316.

- BANDEL, K. & GEYS, J.F. (1985): Regular echinoids in the Upper Cretaceous of the Hashemite Kingdom of Jordan. – *Annales de la Société Géologique du Nord*, **104**: 97-115.
- BARRAS, C.G. (2006): British Jurassic irregular echinoids. – *Monographs of the Palaeontographical Society*, **159/625**: 1-168.
- (2007): Phylogeny of the Jurassic to Early Cretaceous “disasteroid” echinoids (Echinoidea; Echinodermata) and the origins of spatangoids and holasteroids. – *Journal of Systematic Palaeontology*, **5/2**: 134-161.
- BAUMEISTER, J.G. (1999): Vergleich der Funktionsmorphologie und Paläoökologie zweier Rhabdicariden (Echinodermata: Cidaridae). – *Paläontologische Zeitschrift*, **73/3-4**: 319-326.
- BEADLE, S.C. (1991): The Biogeography of Origin and Radiation: Dendrasterid Sand Dollars in the Northeastern Pacific. – *Paleobiology*, **17/4**: 325-339.
- BLAKE, D.B. & ZINSMEISTER, W.J. (1991): A new marsupiate cidaroid echinoid from the Maastrichtian of Antarctica. – *Palaeontology*, **34/3**: 629-635.
- BLASCHKE, F. (1911): Zur Tithonfauna von Stramberg in Mähren. – *Annalen des k. k. Naturhistorischen Hofmuseums*, **25/1-2**: 143-222.
- BOCZAROWSKI, A. (2001): Isolated Sclerites of Devonian non-pelmatozoan Echinoderms. – *Palaeontologia Polonica*, **59**: 1-218.
- BODA, J. (1964): *Catalogus Originalium Fossilium Hungariae. Pars Zoologica*. – 229 pp., Budapest (M. Áll. Földtani Intézet).
- BORGHI, E. (1995): Segnalazione di una nuova forma affine al *Genocidaris maculata* A. AGASSIZ, 1869. Echinodermi fossili emiliani. – In: FRANCOU, C. (ed.): *Bibliotheca V. Studi di scienze naturali*. – pp. 1-14, Piacenza & Vigevano (Tip.Le.Co. Editore).
- (1999): Echinodermi fossili dell’Emilia: i Cidaridi del Plio-Pleistocene. – *Parva Naturalia*, **1999**: 105-120.
- BOYKO, C.B. (2008): Case 3419. *Corystidae* FOSTER & PHILIP, 1978 (Echinodermata, Echinoidea): proposed emendation of spelling to *Corystusidae* to remove homonymy with *Corystidae* SAMOUELLE, 1819 (Crustacea, Brachyura). – *Bulletin of Zoological Nomenclature*, **65/2**: 114-118.
- BRETON, G., NÉRAUDEAU, D., & CUENCA-BOULAT, C. (1992): *Gnatichnus stellarum* ichnosp. nov., trace de broutagne d’un échinide du Campanien des Charentes (France). – *Revue de Paléobiologie*, **11/1**: 219-229.
- BRITO, I.M. (1979): Clipeasteroides cenozoicos do Brasil (Echinodermata Echinoidea). – *Anais da Academia Brasileira de Ciências*, **41/4**: 729-740.
- (1981a): Contribuição á paleontologia do Estado do Pará. A ocorrência de *Abertella* (Echinoidea, Clypearsteroida) na Formação Pirabas. – *Boletim do Museu Paraense Emílio Goeldi, Nova Série, Geologia*, **23**: 1-8.

- (1981b): O gênero *Hemiaster* no Cretáceo do Brasil (Echinoidea Spatagoidea). – In: SANGUINETTI, Y.T. (ed.): Anais do II. Congresso Latino-Americano Paleontologia, Porto Alegre, Rio Grande do Sul, 26-30 Abril 1981, Volume 1. – pp. 399-414, Porto Alegre (Universidade Federal do Rio Grande do Sul).
- (1981c): Os Equinóides Fósseis do Brasil. III – Espatangóides, Exceto Hemiasteridae. – Anais da Academia Brasileira de Ciências, **53/3**: 569-578.
- & RAMIRES, L.V.O. (1974): Equinóides do Inferior do Norte do Brasil. – Anais da Academia Brasileira de Ciências, **46/2**: 263-274.
- BROMLEY, R.G. (1975): Comparative analysis of fossil and recent echinoid bioerosion. – *Palaeontology*, **18/4**: 725-739.
- BUDIN, I.N. (1980): [A new genus of sand dollars of the family Fibulariidae (Echinoidea)]. – *Zoologicheskii Zhurnal*, **59/2**: 305-308.
- (1983): [A new species of sand dollars of the genus *Scaphechinus* from Peter the Great Bay in the Sea of Japan]. – *Biologiya Morya*, **5**: 55-56.
- BUITRÓN, B.E. (1971): Equinoides del Cretácico del norte de México. – *Paleontología Mexicana*, **31**: 5-71.
- (1973): *Tetragramma gloriae* n. sp., Equinoide de la Formación San Lucas (Hauteriviano-Aptiano) de la región de Huetamo, Michoacán. – *Paleontología Mexicana*, **36**: 33-41.
- (1974): Algunas especies de la Familia Conulidae LAMBERT, 1911 (Echinoidea) del Cretácico superior de Chiapas. – *Paleontología Mexicana*, **39**: 5-27.
- CARRASCO, J.F. (2003): Una nueva especie del género *Pericosmus* (Echinoidea, Eoceno). – *Batalleria – Revista de Paleontología*, **11** (Dec. 2002-2003): 25-30.
- (2005): *Calzadaster friasi*, un nuevo genero y una nueva especie de Spatangoida (Echinoidea) del Eoceno de Cataluna. – *Batalleria – Revista de Paleontología*, **12**: 49-56.
- (2006): Una nueva especie del género *Baueria* (Echinoidea) del Eoceno inferior de la cuenca de Tremp-Graus (Zona central surpirenaica). – *Scripta Musei Geologici Seminarii Barcinonensis, Series Palaeontologica*, **1**: 19-34.
- (2007): Una nueva especie del género *Ambipleurus* (Echinoidea) del Eoceno medio del Noreste de España. Primera cita del género en la Península Ibérica. – *Scripta Musei Geologici Seminarii Barcinonensis, Series Palaeontologica*, **2**: 3-18.
- CARTER, B.D. (1987): *Brissus bridgeboroensis*, a new spatangoid echinoid from the Lower Oligocene of southwestern Georgia. – *Journal of Paleontology*, **61/5**: 1043-1046.
- CASO, M.E. (1981): Contribución al estudio de los Echinozoa de México. La familia Mellitidae Stefanini. Descripción de una nueva especie del género *Mellita*. *Mellita eduardobarrosoi* sp. nov. – *Anales del Centro de Ciencias del Mar y Limnología Universidad Nacional Autónoma de México*, **7** (1980)/2: 141-180.

- CHAVANON, S. (1974): L'approche statistique dans l'interprétation de la systématique de l'évolution et de la paléocologie des Échinides des formations du Cénozoïque Bordelais. – Mémoires de l'Institut de Géologie du Bassin d'Aquitaine, **6/1-2**: 1-457.
- CHESHER, R.H. (1972): A new *Paraster* (Echinoidea: Spatangoida) from the Caribbean (Biological results of the University of Miami deep-sea expeditions. 86). – Bulletin of Marine Science, **22/1**: 10-25.
- CHESNUT, D.R.J. & ETTENSOHN, F.R. (1988): Hombergian (Chesterian) echinoderm paleontology and paleoecology, south-central Kentucky. – Bulletins of American Paleontology, **95/330**: 1-102.
- CHEVET, P. & RIGOLLET, L. (1989): Les Oursins fossiles du Callovien de la Sarthe et des confins de l'Orne. – 64 pp., Le Mans (C.E. Renault & Ville du Mans).
- CHIPLONKAR, G.W. & BADVE, R.M. (1972): Palaeontology of the Bagh Beds. II. Echinoidea. – Proceedings of the Indian Academy of Science, Section B, **76/4**: 133-152.
- & BADVE, R.M. (1974): *Malwaster*, a new Cretaceous subgenus of *Hemiaster*. – Bulletin of the Indian Geological Association, **7/1**: 51-53.
- CLAVEL, B. (1989): Échinides du Crétacé Inférieur Jurassien: Stratigraphie, paléocologie, Paléontologie. – Mémoires de la Société neuchâteloise des Sciences naturelles, **11**: 149-182.
- COLLIGNON, M. (1983): Faune Albo-Cénomaniennne de la Formation des marnes de Kazhdumi, région du Fars-Khuzestan (Iran). – Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon Hors Série, **6 (1981)**: 251-291.
- COLLINS, L.S., COATES, A.G., BERGGREN, W.A., AUBRY, M.-P., & ZHANG, J. (1996): The late Miocene Panama isthmian strait. *Geology*, **24/8**: 687-690.
- COPPARD, S.E. (2008): A comparative analysis of the spatangoid echinoid genera *Brissopsis* and *Metalia*: a new genus and species of spatangoid (Echinodermata: Echinoidea: Brissopsidae) from the Philippines and reassignment of *Brissopsis persica* to *Metalia*. – *Zootaxa*, **1760**: 1-23.
- & SCHULTZ, H.A.G. (2006): A new species of *Coelopleurus* (Echinodermata: Echinoidea: Arbaciidae) from New Caledonia. – *Zootaxa*, **1281**: 1-19.
- & VAN NOORDENBURG, H. (2007): A new species of *Lissocidaris* (Echinodermata: Echinoidea: Cidaridae) from the Philippines: convergent evolution among smooth-spined cidaroids. – *Zootaxa*, **1493**: 53-65.
- CUTRESS, B.M. (1976): A new *Prionocidaris* (Echinodermata: Echinoidea) from the Middle Miocene of Florida. – Proceedings of the Biological Society of Washington, **89/12**: 191-198.
- (1980): Cretaceous and Tertiary Cidaroida (Echinodermata: Echinoidea) of the Caribbean area. – Bulletin of American Paleontology, **77/309**: 1-221.
- DAFNI, J. (1983): A new subspecies of *Tripneustes gratilla* from the northern Red Sea (Echinodermata: Echinoidea: Toxopneustidae). – Israel Journal of Zoology, **32/1**: 1-12.

- DARTNELL, A.J. (1972): A brooding echinoid from Tasmania. – Proceedings of the Linnean Society of New South Wales, **97/1** (no. 429): 30-34.
- DAVID, B. & DE RIDDER, C. (1989): Echinodermes: Échinides irréguliers. – In: FOREST, J. (ed.): Résultats des Campagnes Musorstom, Volume 4. – Mémoires du Muséum National d'Histoire Naturelle, Série A (Zoologie), **143/5**: 203-2227.
- & MOOI, R. (1990): An echinoid that “gives birth”: morphology and systematics of a new Antarctic species, *Urechinus mortenseni* (Echinodermata, Holasteroidea). – Zoomorphology, **110/2**: 75-89.
- & MOOI, R. (2000): A new species of subantarctic *Plexechinus* and its phylogenetic position within the Holasteroidea (Echinodermata: Echinoidea). – Polar Biology, **23/3**: 166-172.
- DE RIDDER, C. (1984): Clypeastéroïdes et spatangoïdes littoraux de Nouvelle-Calédonie (Echinodermata). – Bulletin du Muséum national d'Histoire naturelle, 4^e série, Section A (Zoologie, Biologie et Écologie animales), **6/3**: 617-624.
- DE VILLOUTREYS, O. & BIDAR, A. (1973): *Pseudananchys alpina*, nouvelle espèce d'échinide du Cénomanien Des Alpes de Haute-Provence. – Comptes rendus hebdomadaires des séances, Académie des Sciences Paris, Série D, **276/13**: 1977-1979.
- DEL RÍO, C.J., MARTÍNEZ, S.A., STILWELL, J.D., & CONCHEYRO, A. (2007): Palaeontology of the Cerros Bayos section, Roca Formation (Danian), La Pampa Province, Argentina. – Alcheringa, **31/3**: 241-269.
- DEVRIÈS, A. (1972): Contribution à l'étude des échinides fossiles d'Espagne. – Bulletin des Sciences de la Terre de l'Université de Poitiers, **12**: 1-60.
- (1973): Suite aux Echinides d'Algérie. – Publication du Service de la Carte géologique d'Algérie (Paléontologie), Nouvelle Série, Bulletin, **44**: 45-100.
- DOLLFUS, R.-P. & ROMAN, J. (1981): Les Échinides de la mer Rouge. – Ministère des Universités, Comité des Travaux Historiques et Scientifiques, Mémoires de la Section des Sciences, **9**: 1-145.
- DONOVAN, S.K. & ROWE, D.-A.C. (2000): Spatangoid Echinoids from the Eocene of Jamaica. – Journal of Paleontology, **74/4**: 654-661.
- DRAGOMIR, B.-P. (1997): Nouvelles espèces de Salenoides de l'Aptien de Banat. – Acta Palaeontologica Romaniae, **1/29**: 219-220.
- DUDICOURT, J.-C., NÉRAUDEAU, D., NICOLLEAU, P., CEULEMANS, L., & BOUTIN, F. (2005): Une faune remarquable d'échinides marsupiaux dans le Pliocène de Vendée (Ouest de la France). – Bulletin de la Société Géologique de France, **176/6**: 545-557.
- DURHAM, J.W. (1994): Fossil *Encope* (Echinoidea) from the Pacific coast of southern Mexico. – Revista Mexicana de Ciencias Geológicas, **11/1**: 113-116.
- & MORGAN, S.R. (1978): New sand dollars (Echinoidea) of the genera *Merriamaster* and *Dendraster* from Purisima Formation, California. – Proceedings of the California Academy of Sciences, **41/11**: 297-305.

- EGOROV, E.V. (1972): [Stratigraphical distribution of Upper Cretaceous echinoids of Tadzhikistan]. – Doklady Akademii Nauk Tadzhikskoj SSR, **15/5**: 59-61.
- EGOYAN, V.L. (1955): [Upper Cretaceous deposits of south-western Armyanskog SSR]. – 271 pp., Erevan (Akademiya Nauk Armyanskog SSR).
- ELATTAAR, A.A.E.H.A. (2001a): Pliocene-Quaternary echinoids from the Red Sea coast, between Safaga and Quseir, Egypt. – The Second International Conference of the Geology of Africa, Assiut, Egypt, **1**: 635-668.
- (2001b): *Temnotrema* and *Fibularia* (Temnopleuroidea and Clypeasteroidea, Echinoidea) from the Pliocene Rocks of the Red Sea Coast. – Egyptian Journal of Paleontology, **1**: 81-96.
- & STROUGO, A. (2001): Echinoids from the lower and upper Eocene of Sinai. – Middle East Research Center Ain Shams University, Earth Science Series, **15**: 61-78.
- ELBASSYONY, A.A. (2005): *Bahariya teetotumensis* n. gen. n. sp. from the Middle Eocene of Egypt. – Revue de Paléobiologie, **24/1**: 319-329.
- ENDELMAN, L.G. (1980a): [To the revision of the order Hololectypoida (Echinoidea)]. – Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Geologicheskii, **55/1**: 93-103.
- (1980b): [New species of *Neoglobator* (Echinoidea, Hololectypoida) from the Danian-Eocene of the south of the USSR]. – Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Geologicheskii, **55/3**: 93-103.
- ENGEL, H. (1972): *Phymosoma maastrichtensis* spec. nov., a fossil echinoid from the Cretaceous of Maastricht (Echinacea, Phymosomatoida, Phymosomatidae). – Zoologische Mededelingen, **47**: 540-544.
- (1976): On a new echinid from the Cretaceous of Maastricht: *Scutellina supramarginalis* (Echinoidea, Gnathostomata, Clypeasteroidea, Laganina, Fibulariidae). – Bulletin Zoologisch Museum Universiteit van Amsterdam, **5/7**: 55-57.
- ERNST, G. (1971): Biometrische Untersuchungen über die Ontogenie und Phylogenie der *Offaster*/*Galeola*-Stammesreihe (Echin.) aus der nordwesteuropäischen Oberkreide. – Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, **139**: 169-225.
- & SCHULZ, M.-G. (1971): Die Entwicklungsgeschichte der hochspezialisierten Echiniden-Reihe *Infulaster* – *Hagenowia* in der borealen Oberkreide. – Paläontologische Zeitschrift, **45/3-4**: 120-143.
- FELL, F.J. (1975): The echinoid genus *Centrostephanus* in the South Pacific Ocean with a description of a new species. – Journal of the Royal Society of New Zealand, **5/2**: 179-193.
- FISCHER, R. (1985): La fauna de la Formación Turrúcares (Mioceno, Valle Central, Costa Rica). – Geologica et Palaeontologica, **19**: 191-225.
- FORESI, L.M., MAZZEI, R., SALVATORINI, G., & DONIA, F. (2008): Biostratigraphy and chronostratigraphy of the Maltese Lower Globigerina Limestone Member (*Globigerina* Limestone

- Formation): new preliminary data based on calcareous plankton. – *Bollettino della Società Paleontologica Italiana*, **46/2-3**: 175-181.
- FOSTER, R.J. & PHILIP, G.M. (1978): Tertiary holasteroid echinoids from Australia and New Zealand. – *Palaeontology*, **21/4**: 791-822.
- FREST, T.J. & STRIMPLE, H.L. (1977): *Praepholidocidaris*, a new echinoid from the Pella Formation (Mississippian) of Iowa. – *Proceedings of the Iowa Academy of Sciences*, **84/3**: 98-105.
- GABB, W.M. (1881): Descriptions of Caribbean Miocene Fossils. – *Journal of the Academy of Nature Sciences of Philadelphia*, 2nd Series, **8**: 337-348.
- GARRAFIELO, F.J.M. & CARREIRA, M.M.H. (1994): Uma nova espécie de cassidulóide (Echinodermata-Echinoidea) da Formação Pirabas (Mioceno inferior), Pará, Brasil: *Rhyncholampas candidoi* n. sp. – *Boletim do Museu Paraense Emilio Goeldi, Serie Ciencias da Terra*, **6**: 53-58.
- GÄSSER, Z. (1994): Un nou Clypeasteroid de l'Eocè català. – *Batalleria*, **4 (1990-1993)**: 13-16.
- GATT, M. (2005): A new late Oligocene to early Miocene species of *Apatopygus* (Echinoidea: Cassiduloida) from the Maltese Islands. – *Alcheringa*, **29/1**: 101-112.
- GERASIMOV, P.A. (1955): [Manual for Mesozoic fossils from the central region of the European part of the SSSR. Volume 2. Echinoidea, Arthropoda, Annelida, Bryozoa and Corals of the Jurassic Beds]. – 90 pp., Moskva (Ministry for Geology & Mining, SSSR).
- GEYS, J.F. (1979): Salenioid Echinoids from the Maastrichtian (Upper Cretaceous) of Belgium and The Netherlands. – *Paläontologische Zeitschrift*, **53/3-4**: 296-322.
- (1980): Phymosomatoid echinoids from the Campanian and the Maastrichtian of Belgium and the Netherlands. – *Paläontologische Zeitschrift*, **54/3-4**: 199-224.
- (1982): Regular echinoids from the Turonian and the Coniacian (Upper Cretaceous) of the Mons Basin (Belgium). – *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **53/3**: 1-20.
- (1983): *Rachiosoma gigasei* nov. spec. an addition to the echinoderm fauna of the Maastrichtian (Upper Cretaceous) of Belgium. – *Bulletin de la Société Belge de Géologie*, **92/4**: 255-259.
- (1984): *Porosoma kaspicum* n. sp. (Echinoidea, Phymosomatidae) from the Campanian of Mangyshlak (Kazakhstan, U.S.S.R.). – *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, **1984/1**: 27-32.
- (1985): Regular echinoids from the Cenomanian of Hainaut (Belgium and France). – *Bulletin de la Société belge de Géologie*, **94/2**: 129-157.
- (1987): The genus *Typocidaris* (Cidaroida; Echinoidea) in the Upper Cretaceous of the Maastricht area (Belgium and the Netherlands). – *Bulletin de l'Institut Royal des Sciences naturelles de Belgique – Sciences de la Terre*, **57**: 201-215.

- (1989): Hemicidaroid echinoids from Upper Cretaceous deposits in the Wadi Qena-area (Eastern Desert, Egypt). – Bulletin de l'Institut Royal des Sciences naturelles de Belgique – Sciences de la Terre, **59**: 127-135.
- (1992): Regular echinoids, other than Hemicidaroida, from Upper Cretaceous deposits in the Wadi Qena-area (Eastern Desert, Egypt). – Bulletin de l'Institut Royal des Sciences naturelles de Belgique – Sciences de la Terre, **62**: 139-154.
- & MACHALSKI, M. (1992): A new salenioid echinoid, *Salenia sigillata pozaryskae* subsp. n., from the uppermost Maastrichtian of the middle Vistula Valley, central Poland. – Acta Geologica Polonica, **42**/1-2: 135-139.
- GIUSBERTI, L., FANTIN, M., & BUCKERIDGE, J. (2005): *Ovulaster protodecimae* n. sp. (Echinoidea, Spatangoida) and associated epifauna (Cirripedia, Verrucidae) from the Danian of northeastern Italy. – Rivista Italiana di Paleontologia e Stratigrafia, **111**/3: 455-465.
- GLAUERT, L. (1923): *Cidaris comptoni* sp. nov. A Cretaceous echinoid from Gingin. – Journal of the Royal Society of Western Australia, **9**/1: 48-52.
- GONÇALVES, F. (1971): *Echinolampas antunesi* nov. sp. (Cassidulidae s.l.), échinide nouveau du Miocène de la région de Luanda, Angola. – Revista da Faculdade de Ciências, Universidade de Lisboa, Serie 2^a (C, Ciências naturais), **16**/2: 307-310.
- GONGADZE, G.S. (1972): [The representatives of the genus *Conulus* in the Upper Cretaceous of Georgia]. – Trudy Tbilisskogo Gosudavstvennogo Universiteta, Fizico-Matematicheskie i Estestvennye Nauki [Proceedings of Tbilisi University, mathematical and natural sciences], **A3 (144)**: 223-234.
- (1976): [A new species of the genus *Echinocorys* from the Upper Maastrichtian of the southern border region of the Dzirulski massiv]. – Trudy Tbilisskogo Gosudavstvennogo Universiteta, Geografiya-Geologiya [Proceedings of Tbilisi University, Geography & Geology], **172**: 39-42.
- (1979): [The upper Cretaceous Echinoids of Georgia and their stratigraphic value]. – 151 pp., Tbilisi (Tbilisi University Press).
- GRAWE-BAUMEISTER, J., SCHWEIGERT, G., & DIETL, G. (2000): Echiniden aus dem Nusplinger Plattenkalk (Ober-Kimmeridgium, Südwestdeutschland). – Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie), **286**: 1-39.
- GREYLING, E.H. (1996): A new species of *Micraster* (Echinoidea: Spatangoida) and the first record of *Hemimaster* (*Bolbaster*) Pomel from the Upper Cretaceous (Campanian/Maastrichtian) of Zululand. – Durban Museum Novitates, **21**: 25-36.
- & COOPER, M.R. (1993): A new cassiduloid echinoid from the Lower Cretaceous (Upper Albian) of Zululand. – Durban Museum Novitates, **18**: 13-20.
- & COOPER, M.R. (1995): Two new irregular echinoids from the Upper Cretaceous (mid-Campanian) of Angola. – Durban Museum Novitates, **20**: 63-71.
- GUENSBURG, T.E. (1984): Echinodermata of the Middle Ordovician Lebanon Limestone, Central Tennessee. – Bulletins of American Paleontology, **86**/319: 1-100.

- HAGDORN, H. (1995): Die Seeigel des germanischen oberen Muschelkalks. – Geologisch-Paläontologische Mitteilungen Innsbruck, **20**: 245-281.
- HAM, R.W.J.M. van der (1995): *Hemiaster (Leymeriaster) eluvialis*, a new echinoid from the late Maastrichtian of NE Belgium and SE Netherlands. – Bulletin de l'Institut Royal des Sciences naturelles de Belgique – Sciences de la Terre, **65**: 153-164.
- HAROLD, A.S. & TELFORD, M. (1990): Systematics, phylogeny and biogeography of the genus *Melilita* (Echinoidea: Clypeasteroidea). – Journal of Natural History, **24/4**: 987-1026.
- HARPER, J.A. & SHAAK, G.D. (1974): A new *Echinolampas* (Echinodermata, Echinoidea) from the Ocala Limestone of Florida. – Journal of Paleontology, **48/6**: 1166-1169.
- HAUDE, R. (1999): Ein seltener paläozoischer Seeigel aus der Endzeit des Iberg-Atolls (Harz, Adorf-Stufe). – Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, **214/1-2**: 149-167.
- & THOMAS, E. (1994): Eleutherozoen (Echinodermata) aus dem Unter-Karbon von Aprath im Bergischen Land. – In: HACKLER, C., HEINRICH, A., & KRAUSE, E.-B. (eds): Geologie, Paläontologie und Vor- und Frühgeschichte zwischen Lippe und Wupper. – Archäologie im Ruhrgebiet, **2**: 115-132.
- HENDERSON, R.A. (1975): Cenozoic spatangoid echinoids from New Zealand. – Paleontological Bulletin, New Zealand geological Survey, **46**: 1-90.
- HESS, H. (1971): Über einige Echiniden aus Dogger und Malm des Schweizer Juras. – Eclogae Geologicae Helvetiae, **64/3**: 611-633.
- (1972): Eine Echinodermen-Fauna aus dem mittleren Dogger des Aargauer Juras. – Schweizer paläontologische Abhandlungen, **92**: 1-87.
- HOGGETT, A.K. & ROWE, F.W.E. (1986): South-west Pacific cidarid echinoids (Echinodermata) including two new species. – Indo-Malayan Zoology, **3/1**: 1-13.
- HOLMES, F.C. (1995): Australian Tertiary Neolampadidae (Echinoidea): a review and description of two new species. – Proceedings of the Royal Society of Victoria, **107/2**: 113-128.
- (1999): Australian Tertiary Apatopygidae (Echinoidea). – Proceedings of the Royal Society Victoria, **111/1**: 51-70.
- (2004): A new Late Eocene cassiduloid (Echinoidea) from Yorke Peninsula, South Australia. – Memoirs of the Museum Victoria, **61/2**: 209-216.
- , AH YEE, C., & KRAUSE, J. (2005): Two new Middle Miocene spatangoids (Echinoidea) from the Murray Basin, South Australia. – Memoirs of the Museum Victoria, **62/1**: 91-99.
- HYNA, V.A. & MACZYNSKA, S. (1979): *Micraster (Micraster) maleckii* sp. n. from the Santonian of the environs of Cracow, a new echinoid ex gr. *Micraster (Micraster) vaginalae* Nowak. – Prace Muzeum Ziemi, **32**: 21-27.

- ICZN (2009): Opinion 2238 (Case 3419). Corystidae Foster & Philip, 1978 (Echinodermata, Echinoidea) spelling emended to Corystusidae to remove homonymy with Corystidae Samouelle, 1819 (Crustacea, Brachyura). – Bulletin of Zoological Nomenclature, **66**/4: 371-372.
- ILIEVA, T.K. (1998): [Change of the echinoid fauna across the Cretaceous-Tertiary boundary section in the area of the town Byala, Varna District]. – Review of the Bulgarian Geological Society, **59**/3: 133-138.
- IRWIN, R.P. (1995): New family and genus of Clypeasteroidea (Echinoidea) from the Australian Tertiary. – Association of Australasian Palaeontologists, Memoir, **18**: 189-198.
- & ARCHBOLD, N.W. (1994): The spatangoid echinoid *Lovenia* from the Tertiary of southeastern Australia. – Proceedings of the Royal Society of Victoria, **106**: 1-15.
- ISHIJIMA, W. & HATAI, K. (1973): A Problematic Microfossil from the Miocene Mizunami Group, Toki-Mizunami District, Gifu Prefecture, Japan. – St. Paul's Review of Science (= Rikkyo Daigaku Kenkyu Hokoku), **3**/3: 65-68.
- JAGT, J.W.M. (2000): Late Cretaceous-Early Palaeogene echinoderms and the K/T boundary in the southeast Netherlands and northeast Belgium – Part 4: Echinoids. – Scripta Geologica, **121**: 181-375.
- , ANDERSEN, S.B., & JAKOBSEN, S.L. (1998): The echinoid *Gauthieria alterna* from the Uppermost Maastrichtian of Denmark. – Bulletin of the Geological Society of Denmark, **45**/1: 23-26.
- & HAM, R.W.J.M. van der (1995): First record of the echinoid genus *Tylocidaris* Pomel 1883 from the type Maastrichtian of the Netherlands. – Paläontologische Zeitschrift, **69**/1-2: 233-239.
- JAIN, R.L. (2002): Echinoids from the Gaj Formation (Early and Middle Miocene) of Kathiawar, Gujarat, India. – Journal of the Palaeontological Society of India, **47**: 107-135.
- JEFFERY, C.H. (1997a): All change at the K-T boundary? Echinoids from the Maastrichtian and Danian of the Mangyshlak Peninsula, Kazakhstan. – Palaeontology, **40**/3: 659-712.
- (1997b): Revision of the echinoid *Enallopneustes* from the Upper Cretaceous of North Africa. – Cretaceous Research, **18**/2: 237-248.
- (1999): A reappraisal of the phylogenetic relationships of somaliasterid echinoids. – Palaeontology, **42**/6: 1027-1041.
- JENSEN, M. (1982): Morphology and classification of Euechinoidea BRONN, 1860 – a cladistic analysis. – Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening, **143** (1981): 7-99.
- JESIONEK-SZYMAŃSKA, W. (1978): On a new galeropygid genus (Echinoidea) from the Jurassic (Upper Lias) of Morocco. – Acta Palaeontologica Polonica, **23**/2: 187-193.

- (1979): Morphology and microstructure of oligolamellar teeth in Paleozoic echinoids. Part 1. Teeth of some early lepidocentrid echinoids. – *Acta Palaeontologica Polonica*, **24/2**: 275-293.
- KADIL'NIKOVA, Y.I. & MOSKVIN, M.M. (1984): [Danian and Paleocene species of the sea urchins *Studeria* and *Hypsopygaster*]. – *Paleontologicheskii Zhurnal*, **1984/1**: 37-46.
- KIER, P.M. (1972a): Tertiary and Mesozoic Echinoids of Saudi Arabia. – *Smithsonian Contributions to Paleobiology*, **10**: 1-105.
- (1972b): Upper Miocene Echinoids from the Yorktown Formation of Virginia and their environmental significance. – *Smithsonian Contributions to Paleobiology*, **13**: 1-41.
- (1973): A new Silurian echinoid genus from Scotland. – *Palaeontology*, **16/4**: 651-663.
- (1977): Triassic echinoids. – *Smithsonian Contributions to Paleobiology*, **30**: iv+1-88.
- (1980): The Echinoids of the Middle Eocene Warley Hill Formation, Santee Limestone, and Castle Hayne Limestone of North and South Carolina. – *Smithsonian Contributions to Paleobiology*, **39**: 1-102.
- (1982): Echinoids. – In: SPRINKLE, J. (ed.): *Echinoderm Faunas from the Bromide Formation (Middle Ordovician) of Oklahoma*. – *University of Kansas Paleontological Contributions, Monograph*, **1**: 310-314.
- (1983): Upper Cenozoic Echinoids from the Lee Creek Mine. – In: RAY, C.E. (ed.): *Geology and Paleontology of the Lee Creek Mine, North Carolina, I*. – *Smithsonian Contributions to Paleobiology*, **53**: 499-507.
- (1984a): Echinoids from the Triassic (St. Cassian) of Italy, their lantern supports, and a revised phylogeny of Triassic echinoids. – *Smithsonian Contributions to Paleobiology*, **56**: iii+1-41.
- (1984b): Fossil spatangoid echinoids of Cuba. – *Smithsonian Contributions to Paleobiology*, **55**: vi+1-336.
- (1992): Neogene paleontology in the northern Dominican Republic. 13. The Class Echinoidea (Echinodermata). – *Bulletin of American Paleontology*, **102/339**: 13-27.
- (1997): Oligocene Echinoids of North Carolina. – *Smithsonian Contributions to Paleobiology*, **83**: 1-37.
- KINMAN, K.E. (1994): The Kinman system: towards a stable cladistic-eclectic classification of organisms (living and extinct; 48 phyla, 269 classes, 1,719 orders). – ii+88 pp., Hays, KS (Kinman).
- KOLATA, D.R. (1975): Middle Ordovician echinoderms from Northern Illinois and Southern Wisconsin. – *Journal of Paleontology – The Palaeontological Society Memoir*, **7**: 1-74.
- , STRIMPLE, H.L., & LEVORSON, C.O. (1977): A new species of *Bothriocidaris* (Echinoidea) from the Cincinnati Maquoketa Group of Iowa. – *Proceedings of the Iowa Academy of Sciences*, **84/4**: 146-149.

- KÖNIG, H. (1982): Unterpermische Seeigel aus Kreta (Griechenland). – Neues Jahrbuch für Geologie und Paläontologie, Monatshefte, **1982/3**: 165-175.
- KOTCHETOFF, B., KOTCHETOFF, Y., & VEIGA FERREIRA, O. de (1975): Contribution à la connaissance des gisements fossilifères miocènes au Nord du Cap d'Espichel. – Comunicações Serviços Geológicos de Portugal, **59**: 59-106.
- KROH, A. (2004): *Cottreaucorys kollmanni* n. sp. (Echinodermata: Echinoidea) and other new records from the Early Paleocene Bruderndorf Formation in Austria. – Annalen des Naturhistorischen Museums in Wien, Serie A, **106**: 309-325.
- (2005): Catalogus Fossilium Austriae. Band 2. Echinoidea neogenica. – lvi+210 pp., Wien (Österreichische Akademie der Wissenschaften).
- (2007): *Hemipatagus*, a misinterpreted Loveniid (Echinodermata: Echinoidea). – Journal of Systematic Palaeontology, **5/2**: 163-192.
- & JAGT, J.W.M. (2004): Additional records of Late Cretaceous and Early Palaeogene echinoids from the Gschlifgraben (Ultrahelvetic, Upper Austria). – Acta Geologica Polonica, **54/ 4**: 551-571.
- KRÜGER, F.J. (1993): Plattenanomalien bei *Echinocorys sulcata* (Goldfuss 1826) aus dem Dan (Alttertiär) von Dalbyover, Dänemark. – Paläontologische Zeitschrift, **67/3-4**: 305-321.
- (2002): x-mal Feuerstein: Zweiter Fund eines Galerites mit sieben Ambulakren. – Der Geschiebe-Sammler, **34/4**: 173-181.
- (2006): Ambulakrale Anomalien bei fossilen und rezenten Echinoidea (Echinodermata). – Der Aufschluss, **57**: 59-64.
- KRUSE, P.D. & PHILIP, G.M. (1985): Tertiary species of echinoid genus *Eupatangus* from southern Australia. – Special Publications, South Australian Department of Mines and Energy, **5**: 167-185.
- KÜHN, O. (1930): Das Danien der äusseren Klippenzone bei Wien. – Geologische und Palaeontologische Abhandlungen, Neue Folge, **17/5**: 495-576.
- KUTSCHER, M. (1978a): Neue Echiniden aus dem Unter-Maastricht der Insel Rügen. I. Holasteridae DURHAM & MELVILLE. – Zeitschrift für geologische Wissenschaften, **6/5**: 627-639.
- (1978b): Neue Echiniden aus dem Unter-Maastricht der Insel Rügen. II Spatangoida CLAUS, 1876. – Zeitschrift für geologische Wissenschaften, **6/8**: 1025-1037.
- (1985a): Die Echinodermen des Magdeburger Grünsandes (Mittel-Oligozän). – Abhandlungen und Berichte für Naturkunde und Vorgeschichte, **12/6**: 3-14.
- (1985b): Neue Echiniden aus dem Unter-Maastricht der Insel Rügen. Vertreter der Ordnungen Echinothurioida CLAUS, 1880, Diadematoidea DUNCAN, 1889 und Phymosomatoida MORTENSEN, 1904. – Zeitschrift für geologische Wissenschaften, **13/2**: 235-247.
- (1985c): Neue Echiniden aus dem Unter-Maastricht der Insel Rügen. Vertreter der Phymosomatidae POMEL, 1883. – Zeitschrift für geologische Wissenschaften, **13/4**: 521-532.

- LAMBERT, J. (1911): Description des échinides crétacés de la Belgique, principalement de ceux conservés au Musée royal de Bruxelles. II. Note sur les Échinides de l'étage Sémonien. – Mémoire du Musée Royal d'Histoire Naturelle de Belgique, **4**: 1-81.
- (1926): Sur des *Rhabdocidaris orbigny* Agassiz, communiqués par M. G. Schneider de Bâle. – Eclogae Geologicae Helvetiae, **19**: 755-763.
- (1931): Echinides Crétacés de la région d'Héraclée. – Annales de la Société Géologique de Belgique, **54 (Mémoires)**: M1-M12.
- LARRAIN, A.P. (1975): Los equinoideos regulares fosiles y recientes de Chile. – Gayana Zoolgia, **35**: 5-189.
- (1985a): *Brachysternaster*, new genus and *Brachysternaster chesheri*, new species of Antarctic echinoid (Spatangoida, Schizasteridae). – Polar Biology, **4/3**: 121-124.
- (1985b): A new, early *Hemiaster* (Echinodermata: Echinoidea) from the Lower Cretaceous of Antofagasta, northern Chile. – Journal of Paleontology, **59/6**: 1401-1408.
- (1986): A new species of subantarctic echinoid (Echinoidea: Schizasteridae). – Boletín de la sociedad de Biología de Concepción, **56 (1985)**: 115-119.
- & BIRÓ-BAGÓCZKY, L. (1985): New *Pygurus* (Echinodermata: Echinoidea) from the Tithonian of central Chile: first record from the Jurassic of the Southern Hemisphere. – Journal of Paleontology, **59/6**: 1409-1413.
- LEWIS, D.N. (1986): The Cretaceous echinoid *Boletechinus*, with notes on the phylogeny of the Glyphocyphidae and Temnopleuridae. – Bulletin of the British Museum (Natural History), Geology Series, **40/2**: 59-90.
- (1989): Fossil Echinoidea from the Barton Beds (Eocene, Bartonian) of the type locality at Barton-on-Sea in the Hampshire Basin, England. – Tertiary Research, **11/1**: 1-47.
- & ENSOM, P.C. (1982): *Archaeocidaris whatleyensis* sp. nov. (Echinoidea) from the Carboniferous Limestone of Somerset, and notes on echinoid phylogeny. – Bulletin of the British Museum (Natural History), Geology Series, **36**: 77-104.
- & JEFFERIES, R.P.S. (1980): *Salenia trisuranalis* sp. nov. (Echinoidea) from the Eocene (London Clay) of Essex, and notes on its phylogeny. – Bulletin of the British Museum (Natural History), Geology Series, **33**: 115-121.
- LI, F., ZHANG, X., HE, L., & ZHANG, K. (1985): Paleozoic echinoids of south China. – Earth Science, Journal of the Wuhan College of Geology, **10/Special issue**: 97-103.
- LIAO, Y. (1978): The echinoderms of the Xisha islands, Guangdong province, China. III. Echinoidea. – Studia Marina Sinica, **12**: 107-132.
- (1979): A new genus of clypeasteroid sea-urchin from Huang Hai. – Oceanologia et Limnologia Sinica, **10/1**: 67-72.
- & LIN, C. (1981): [A new echinoid with sexual dimorphism from the Late Tertiary deposits of Beibuwan, Guangxi]. – Acta Palaeontologica Sinica, **20/5**: 482-484.

- LINDER, R.A., DURHAM, J.W., & ORR, W.N. (1988): New Late Oligocene echinoids from the central Western Cascades of Oregon. – *Journal of Paleontology*, **62/6**: 945-958.
- LINDLEY, I.D. (2003): Echinoids of the Kairuku Formation (Lower Pliocene), Yule Island, Papua New Guinea: Spatangoida. – *Proceedings of the Linnean Society of New South Wales*, **124**: 153-162.
- (2004): Some living and fossil echinoderms from the Bismark Archipelago, Papua New Guinea and two new echinoid species. – *Proceedings of the Linnean Society of New South Wales*, **125**: 115-139.
- LOBATSHEVA, S.V. (1961): [Representatives of the genus *Toxaster* from the Neocomian of Comet-Dag and Bolshogo Balkhan]. – In: [Problems of the oil and gas-bearing deposits of Central Asia]. – *Trudy Vsesoiuznogo nauchno-issledovatel'skogo geologicheskogo instituta*, **46**: 151-173. [not seen]
- LÓPEZ, J. & SILLERO, C. (2006): Equinoides fósiles de la Provincia de Alicante. – *Memorias del Mupe*, **2**: 1-254.
- LUCAS, S.G. (2000): A new species of the echinoid *Hemiaster* from the Lower Cretaceous (Albian) of southwestern New Mexico. – *New Mexico Museum of Natural History and Science, Bulletin*, **16**: 101-106.
- MADON-SENEZ, C. (2002): A new species of marsupiate Antarctic echinoid: *Amphipneustes davidi* (Echinodermata: Spatangoida: Schizasteridae). – *Proceedings of the Biological Society of Washington*, **115/1**: 51-56.
- MALINOWSKA, L. (ed.) (1989): *Gology of Poland. Volume III. Atlas of Guide and Characteristic Fossils Part 2c Mesozoic, Cretaceous*. – Warsaw (Wydawnictwa Geologiczne).
- MAMEDOV, T.A. & MELIKOV, O.G. (1976): Novye vidy morskikh ezheg iz Ehotsenovykh otlozhenig Respubliki Mali (Zapadiaya Afrika). – *Voprosy Paleontologii i Stratigrafii Azerbajdzana*, **1**: 156-166.
- MANSO, C.L.d.C. & SOUZA-LIMA, W. (2007): *Holosalenia bahiensis*, um novo equinoide (Echinodermata) do Albiano da bacia de Camamu, Brasil. – *Revista Brasileira de Paleontologia*, **10/1**: 27-34.
- MARCOPOULOU-DIACANTONI, A. (1972): Échinides (*Clypeaster*, *Schizaster*, *Spatangus*, *Brissopsis*) de l'Helvétien de l'île de Crète centrale et Orientale. – *Annales Géologiques des Pays Helléniques*, **24**: 133-160.
- MÄRKEL, K. (1969): Morphologie der Seeigelzähne. II. Die gekielten Zähne der Echinacea (Echinodermata, Echinoidea). – *Zeitschrift für Morphologie der Tiere*, **66**: 1-50.
- MARKOV, A.V. (1988a): [Deep-sea urchins of the genus *Salenia* (Echinoidea)]. – *Zoologicheskii Zhurnal*, **67/3**: 375-383.
- (1988b): [Sea urchins of the family Saleniidae: composition and distribution]. – *Zoologicheskii Zhurnal*, **67/4**: 567-575.

- (1989): [Species composition of the genus *Trigonocidaris* (Echinoidea)]. – Zoologicheskii Zhurnal, **68/8**: 75-84.
- (1990): A new genus of heart urchins from south Atlantic. – Zoologicheskii Zhurnal, **69/1**: 54-60.
- (1991): [The most deep-sea representative of the family Schizasteridae (Echinoidea)]. – Zoologicheskii Zhurnal, **70/8**: 153-155.
- (1994): [Morphology, systematics and phylogeny of schizasterid sea urchins]. – Trudy Paleontologicheskogo Instituta, **258**: 1-94.
- & Endelman, L.G. (1991): [New findings of Upper Cretaceous and Eocene temnopleurids (Echinoidea)]. – Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody. Otdel Geologicheskii, **66/1**: 74-86.
- & SOLOVJEV, A.N. (2001): [Echinoids of the family Paleopneustidae (Echinoidea, Spatangoida): morphology, taxonomy, phylogeny]. – Geos – Trudy Paleontologicheskogo Instituta, Rossikaia Akademia Nauk, **280**: 1-108.
- MARTÍNEZ, S. (1984): *Amplaster coloniensis* n. g. n. sp. (Echinoidea: Monophorasteridae) del Mioceno de Uruguay. – In: DEL CARMEN, P.M. (ed.): Memoria III Congreso Latinoamericano de Paleontología. – pp. 505-508, Mexico, D.F. (Instituto de Geología de la UNAM).
- , REICHLER, V., & MOOI, R. (2005): A new species of *Abertella* (Echinoidea: Scutellina) from the Gran Bajo del Gualicho Formation (Late Early Miocene-Early Middle Miocene), Río Negro Province, Argentina. – Journal of Paleontology, **79/6**: 1229-1233.
- MAYORAL, E. & MUNIZ, F. (2001): New Ichnospecies of *Cardioichnus* from the Miocene of the Guadalquivir Basin, Huelva, Spain. – Ichnos, **8/1**: 69-76.
- McKINNEY, M.L., McNAMARA, K.J., & WIEDMAN, L.A. (1988): Echinoids from the La Meseta Formation (Eocene), Seymour Island, Antarctica. – In: FELDMANN, R.M. & WOODBURNE, M.O. (eds): Geology and paleontology of Seymour Island, Antarctic Peninsula. – Geological Society of America, Memoir, **169**: 499-503.
- McKNIGHT, D.G. (1968): Additions to the echinoid fauna of New Zealand. – New Zealand Journal of marine and freshwater Research, **2/1**: 90-110.
- (1974): Some echinoids new to New Zealand waters. – New Zealand Oceanographic Institution Records, **2/3**: 25-44.
- McNAMARA, K.J. (1982a): A new species of the echinoid *Rhynobrissus* (Spatangoida: Brissidae) from North-West Australia. – Records of the Western Australian Museum, **9/4**: 349-360.
- (1982b): Taxonomy and evolution of living species of *Breynia* (Echinoidea: Spatangoida) from Australia. – Records of the Western Australian Museum, **10/2**: 167-197.
- (1984): Living Australian species of the echinoid *Pericosmus* (Spatangoida: Pericosmidae). – Records of the Western Australian Museum, **11/2**: 87-100.

- (1985a): The spatangoid echinoid *Linthia* from the Late Eocene of Southern Australia. – Transactions of the Royal Society of South Australia, **109**/4: 161-165.
- (1985b): Taxonomy and evolution of the Cainozoic spatangoid echinoid *Protenaster*. – Palaeontology, **28**/2: 311-330.
- (1986): First Mesozoic record of the cidaroid echinoid *Goniocidaris*. – Alcheringa, **10**/4: 353-354.
- (1987a): An enigmatic *Echinolampas* (Echinoidea) from the Miocene of Victoria. – Proceedings of the Royal Society of Victoria, **99**/ 3-4: 109-110.
- (1987b): The holasteroid echinoid *Echinocorys* from the Maastrichtian of Western Australia. – Records of the Western Australian Museum, **13**/3: 419-426.
- (1987c): Taxonomy, evolution, and functional morphology of southern Australian Tertiary hemiasterid echinoids. – Palaeontology, **30**: 319-352.
- (1989): *Echinolampas laubei* (Echinoidea), a replacement name for *Echinolampas duncani* McNamara, 1987. – Journal of Paleontology, **63**/2: 257.
- (1993a): First record of a spatangoid echinoid from the Paleocene of the Perth Basin, Western Australia. – Records of the Western Australian Museum, **16**/3: 331-336.
- (1993b): A new genus of brissid echinoid from the Upper Oligocene of Victoria. – Proceedings of the Royal Society of Victoria, **105**/1: 39-48.
- (1996): Plio-Pleistocene *Peronella* (Echinoidea: Clypeasteroidea) from Western Australia. – Records of the Western Australian Museum, **18**/2: 193-202.
- & AH YEE, C. (1989): A new genus of brissid echinoid from the Miocene of Australia. – Geological Magazine, **126** /2: 177-186.
- & BARRIE, D.J. (1993): A new genus of marsupiate spatangoid echinoid from the Miocene of South Australia. – Records of the South Australian Museum, **26** (1992)/2: 139-147.
- & KENDRICK, G.W. (1994): Cenozoic molluscs and echinoids of Barrow Island, Western Australia. – Records of the Western Australian Museum Supplement, **51**: 1-50.
- & MELIKOV, O.H. (2002): The asterostomatid echinoid *Antillaster* from the Paradash Group (Middle Eocene) of the Nakhichevan region of Azerbaijan. – Records of the Western Australian Museum, **21**/2: 157-165.
- & PHILIP, G.M. (1980a): Australian Tertiary schizasteroid echinoids. – Alcheringa, **4**/1-2: 47-65.
- & PHILIP, G.M. (1980b): Living Australian schizasterid echinoids. – Proceedings of the Linnean Society of New South Wales, **104**(1979)/2: 127-146.
- & PHILIP, G.M. (1980c): Tertiary species of *Echinolampas* (Echinoidea) from southern Australia. – Memoirs of the National Museum of Victoria, **41**: 1-14.

- & PHILIP, G.M. (1984): A Revision of the Spatangoid Echinoid *Pericosmus* from the Tertiary of Australia. – Records of the Western Australian Museum, **11**/4: 319-356.
- , PHILIP, G.M., & KRUSE, P.D. (1986): Tertiary Brissid echinoids of southern Australia. – Alcheringa, **10**/1: 55-86.
- MELIKOV, O.H. (1989): [Morphological peculiarity of the new species of the genus *Heteraster* Orbigny, 1853 (Echinoidea) from the Urganian facies of Azerbaijan]. – In: KALO, D.D., ROZHNOV, S.V., SOLOVJEV, A.N., STUKALINA, G.A., & KHINTS, I.M. (eds): Fossil and recent echinoderm researches. – pp. 156-160, Tallin (Academy of Sciences of the Estonian SSR, Institute of Geology).
- & KHLILOV, A.G. (1988): [Echinoidea]. – In: ALI-ZADE, A. (ed.): [Cretaceous fauna of Azerbaijan]. – pp. 184-213, Baku (Akademiya Nauk Azerbajdzhanskoj SSR, Institut Geologii I.M. Gubkina / Izdatel'vstvo Elm).
- MICHALÍK, J. (1977): Systematics and ecology of *Zeilleria* BAYLE and other brachiopods in the Uppermost Triassic of the West Carpathians. – Geologický Zborník – Geologica Carpathica, **28**/2: 323-346.
- MIHÁLY, S. (1980): Echinoidea maradványok a Bükk hegység felsőpermjéből. – A Magyar Állami Földtani Intézet Évi Jelentése, **1978**: 399-412.
- (1985): Felső-Bádeni Echinoideák Budapesti új Feltárásokból. – A Magyar Állami Földtani Intézet Évi Jelentése, **1983**: 235-262.
- (1990a): Két új Echinoidea faj a Magyarországi Miocénből. – A Magyar Állami Földtani Intézet Évi Jelentése, **1987**: 237-249.
- (1990b): Új Echinoidea faj a Bükk hegység felső-karbonjából. – A Magyar Állami Földtani Intézet Évi Jelentése, **1987**: 251-257.
- (1981): Az aggteleki-karszt középső-Triász echinoideái. – A Magyar Állami Földtani Intézet Évi Jelentése, **1979**: 297-331.
- MIRONOV, A.N. (1971): [Soft sea urchins of the family Echinothuriidae collected by the R/V “Vityaz” and the “Academician Kurchatov” in the Pacific and Indian Oceans]. – Trudy Instituta Okeanologii Akademii Nauk SSSR, **92**: 317-325.
- (1973): [New deep-sea species of sea urchins of the genus *Echiocrepis* and the distribution of the family Pourtalesiidae (Echinoidea, Meridosterniana)]. – Trudy Instituta Okeanologii Akademii Nauk SSSR, **91**: 240-247.
- (1974a): [Pourtalesiid sea urchins of the Antarctic and Subantarctic (Echinoidea: Pourtalesiidae)]. – Trudy Instituta Okeanologii Akademii Nauk SSSR, **98**: 240-252.
- (1974b): [New species of the abyssal sea urchins of the genus *Echinosigra* (Echinoidea, Pourtalesiidae)]. – Zoologicheskii Zhurnal, **53**/12: 1803-1806.
- (1975): [Deep-sea urchins (Echinodermata, Echinoidea) collected during the 14th cruise of the R/V “Akademik Kurchatov”]. – Trudy Instituta Okeanologii, Akademii Nauk SSSR, **100**: 205-214.

- (1976): [Deep-sea urchins of the northern Pacific]. – Trudy Instituta Okeanologii, Akademii Nauk SSSR, **99**: 140-164.
- (1978a): [Meridosternous echinoids (Echinoidea: Meridosternina) collected during the 16th cruise of the R/V “Dm. Mendeleev”]. – Trudy Instituta Okeanologii, Akademii Nauk SSSR, **113**: 208-226.
- (1978b): [The most deep-sea species of sea urchins (Echinoidea, Pourtalesiidae)]. – Zoologicheskii Zhurnal, **57/5**: 721-726.
- (1981): Sea-urchins (Echinoidea). – In: KUZNETSOV, A.P. & MIRONOV, A.N. (eds): Benthos of the submarine mountains Marcus-Necker and adjacent pacific regions. – pp. 131-140, Moscow (Akademiya Nauk SSSR).
- (1993): [Deep-sea echinoids (Echinodermata: Echinoidea) of the South Atlantic]. – Trudy Instituta Okeanologii, Akademii Nauk SSSR, **127**: 218-227.
- (1995a): [Holasteroid sea urchins. 1. Morphological diversity of *Pourtalesia jeffreysi*]. – Zoologicheskii Zhurnal, **74/11**: 68-77.
- (1995b): [Holasteroid echinoids. 2. The genus *Pourtalesia*]. – Zoologicheskii Zhurnal, **74/12**: 59-75.
- (1996): [Holasteroid sea urchins. 3. *Helgocystis* and a new genus *Rictocystis*]. – Zoologicheskii Zhurnal, **75/8**: 1109-1119.
- (1997): [Holasteroid sea urchins. 4. *Echinosigra*]. – Zoologicheskii Zhurnal, **76/2**: 173-186.
- & SAGAIDACHNY, A.Y. (1984): [Morphology and distribution of the recent echinoids of the genus *Echinocyamus* (Echinoidea: Fibulariidae)]. – Trudy Instituta Okeanologii, Akademii Nauk SSSR, **119**: 179-204.
- (2008): Pourtalesiid sea urchins (Echinodermata: Echinoidea) of the northern Mid-Atlantic Ridge. – Marine Biology Research, **4/1-2**: 3-24.
- MONACO, P., GIANNETTI, A., CARACUEL, J.E., & YEBENES, A. (2005): Lower Cretaceous (Albian) shell-armoured and associated echinoid trace fossils from the Sacaras Formation, Serra Gelada area, southeast Spain. – Lethaia, **28/4**: 333-344.
- MOOI, R. (1990a): A New “Living Fossil” Echinoid (Echinodermata) and the Ecology and Palaeobiology of Caribbean Cassiduloids. – Bulletin of Marine Science, **46/3**: 688-700.
- (1990b): Progenetic miniaturization in the sand dollar *Sinaechinocyamus*: implications for clypeasteroid phylogeny. – In: DE RIDDER, C., DUBOIS, P., LAHAYE, M.-C., & JANGOUX, M. (eds): Echinoderm Research. – Proc. 2nd European Echinoderm Conf., Brussels, 18-21 Sept. 1989. – pp. 137-143, Rotterdam (A. A. Balkema).
- & DAVID, B. (1996): Phylogenetic analysis of extreme morphologies: deep-sea holasteroid echinoids. – Journal of Natural History, **30/6**: 913-953.

- , DAVID, B., FELL, J.F., & CHONE, T. (2000): Three new species of bathyal cidaroids (Echinodermata: Echinoidea) from the Antarctic region. – Proceedings of the Biological Society Washington, **113**/1: 224-237.
- , MARTÍNEZ, S., & PARMA, S.G. (2000): Phylogenetic systematics of Tertiary monophorasterid sand dollars (Clypeasteroidea: Echinoidea) from South America. – Journal of Paleontology, **74**/2: 263-281.
- & PETERSON, D. (2000): A new species of *Leodia* (Clypeasteroidea: Echinoidea) from the Neogene of Venezuela and its importance in the phylogeny of mellitid sand dollars. – Journal of Paleontology, **74**/6: 1083-1092.
- MOSKVIN, M.M. (1964): [Echinoids of the genus *Protobrissus* from the Danian and Paleocene sediments of the USSR]. – In: Sbornik v chest' akademika Y. S. Yovcheva. – pp. 185-202, Sofia.
- (1982): New Upper Cretaceous and Paleocene sea urchins of the family Palaeostomatidae. – Paleontological Journal, **16**: 100-107.
- (1983): [*Pseudogibbaster*, a new genus of Danian and Paleocene echinoids]. – Paleontologicheskii Zhurnal, **1983**/3: 112-115.
- (1984): [The echinoids *Domechinus* and *Hardouinia* from the Upper Cretaceous of the Middle Asia]. – Paleontologicheskii Sbornik, **21**: 63-69.
- & SHIMANSKAYA, N.V. (1981): [Palaeocene *Echinocorys* (sea urchins) from the territory of USSR]. – Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody. Otdel Geologicheskii, **56**/3: 127-137.
- & SHIMANSKAYA, N.V. (1993): [Genus *Echinocorys* (Echinoidea) in the Danian of eastern part of the European paleobiogeographic province. Part 2]. – Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody. Otdel Geologicheskii, **68**/2: 44-59.
- & SHIMANSKAYA, N.V. (1997): [New species of echinoid from Upper Maastrichtian Mangyshlak]. – Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Geologicheskii, **72**/1: 70-71.
- MU, E. & WU, Y. (1976): [Cretaceous and Tertiary echinoids from the Mount Jolmo Lungma Region]. – In: Academica sinica. Tibetan scientific expeditional team (eds): [A report of the scientific expedition in the Mount Jolmo Lungma Region (1966-1968). Palaeontology fasc. 2]. – pp. 357-386, Beijing (Science Press).
- MACZYŃSKA, S. (1984): Typ Echinodermata. – In: MALINOWSKA, L. (ed.): Budowa geologiczna Polski, tom III; Atlas skamieniałości przewodnich i charakterystycznych części 2c; Mesozoik, Kreda. – pp. 425-461, Warszawa (Wydawnictwa Geologiczne).
- (1989): Type Echinodermata. – In: MALINOWSKA, L. (ed.): Geology of Poland. Volume III. Atlas of Guide and Characteristic Fossils Part 2c Mesozoic, Cretaceous. – pp. 300-315, Warszawa (Wydawnictwa Geologiczne).

- NASTAJ, W. (1985): *Pygaulus althi* (ZARĘCZNY) and *Pygaulus richardi* sp. n. (Echinoidea, Cassiduloidea) from Cenomanian conglomerates near Cracow. – *Zeszyty Naukowe Akademii Gorniczo-Hutnicznej im Stanislawia Staszica Geologia*, **11/1**: 17-23.
- NÉRAUDEAU, D., BARBE, S., MERCIER, D., & ROMAN, J. (2003): Signatures paléoclimatiques des échinides du Néogène final atlantique à faciès redonien. – *Annales de Paléontologie*, **89**: 153-170.
- , CRAME, J.A., & KOOSER, M. (2000): Upper Cretaceous Echinoids from Ross Basin, Antarctica. – *Géobios*, **33/4**: 455-466.
- NESTLER, H. (1978): Echiniden aus dem Unter-Maastricht der Insel Rügen. VI. Ein neuer Vertreter der Temnopleuridae. – *Zeitschrift für geologische Wissenschaften*, **6/5**: 619-625.
- (1981): Ein neuer *Palaechinus* (Echinoidea) aus dem Devon. – *Zeitschrift für geologische Wissenschaften*, **9/5**: 553-559.
- (1998): Kretazische und eozane Echiniden aus dem Südjemem. – *Zeitschrift fuer Geologische Wissenschaften*, **26/5-6**: 563-607.
- NEUMANN, C. (1999): New spatangoid echninoids (Echinodermata) from the Upper Cretaceous of Jordan: their taxonomy and phylogenetic importance. – *Berliner geowissenschaftliche Abhandlungen*, **E30**: 175-189.
- NICOLLEAU, P. & VADET, A. (1995): Les oursins des Marnes à Spongiaires de l'Oxfordien du Poitou. – In: BRANGER, P., NICOLLEAU, P., & VADET, A. (eds): *Les ammonites et les oursins de l'Oxfordien du Poitou (faciès à spongiaires de l'Oxfordien moyen et supérieur)*. – pp. 55-81, Aiffres (Musées de la Ville de Niort & Assoc. Promotion Géol. Paléont. Poitou-Charentes-Vendée).
- OLSZEWSKA-NEJBERT, D. (2007): Late Cretaceous (Turonian – Coniacian) irregular echinoids of western Kazakhstan (Mangyshlak) and southern Poland (Opole). – *Acta Geologica Polonica*, **57/ 1**: 1-87.
- OYEN, C.W. & PORTELL, R.W. (1996): A new species of *Rhyncholampas* (Echinoidea: Cassidulidae) from the Chipola Formation: The first confirmed Member of the Genus from the Miocene of the Southeastern USA and the Caribbean. – *Tulane Studies in Geology and Paleontology*, **29/2**: 59-66.
- PARMA, S.G. (1985): *Eoscutella* GRANT y HERTLEIN (Echinodermata: Clypeasteroidea) en el Patagiano (Terciario Inferior) de la Provincia de Santa Cruz, Republica Argentina. – *Ameghiana* – *Revista de la Asociación Paleontología Argentina*, **22/1-2**: 35-41.
- (1989): Los primeros Cassidulidae (Echinodermata: Cassiduloidea) de la Formacion Roca (Paleoceno), Casa de Piedra (limite entre las provincias de Rio Negro y La Pampa), Rep. Argentina. – In: *IV Congreso Argentino de Paleontología y Bioestratigrafía: Mendoza, Noviembre 23-27, 1986. Actas 4.* – pp. 93-98, Mendoza.
- & CASADÍO, S. (2005): Upper Cretaceous-Paleocene echinoids from Northern Patagonia, Argentina. – *Journal of Paleontology*, **79/6**: 1072-1087.

- & MARTÍNEZ, S. (1995): Nuevo género de Monophorasteridae (Echinodermata) del Terciario de Patagonia. – In: IV Congreso Argentino de Paleontología y Bioestratigrafía: Trelew-Chubut, Abril 3-8, 1994, Resúmenes. – pp. 65-66, Museo Paleontológico Egidio Feruglio).
- PAWSON, D.L. (1964): The echinoid genus *Caenopedina* in New Zealand. – Transactions of the Royal Society of New Zealand, Zoology, **5**/5: 63–66.
- (1978): The echinoderm fauna of Ascension Island, South Atlantic Ocean. – Smithsonian Contributions to the Marine Sciences, **2**: i-iv, 1-31.
- & PHELAN, T.H. (1979): *Clypeaster kieri*, a new species of clypeasteroid (Echinodermata: Echinoidea) from off Bombay, India. – Proceedings of the Biological Society of Washington, **92**/4: 796-800.
- PHELAN, T.H. (1972): Comments on the echinoid genus *Encope* and a new subgenus. – Proceedings of the Biological Society of Washington, **85**/8: 109-130.
- PHILIP, G.M. & FOSTER, R.J. (1971): Marsupiate Tertiary echinoids from south-eastern Australia and their zoogeographic significance. – Palaeontology, **14**/4: 666-695.
- PHILIPPE, M. (1989): Révision des Échinides miocènes du Bassin du Rhône: résultats d'ordre systématique. – Actes 6ème Séminaire international Echinodermes, Les Embiez, 1988; Vie Marine, Hors Serie, **10**: 24-35.
- (1998): Les Échinides miocènes du Bassin du Rhône: révision systématique. – Nouvelles Archives du Muséum d'Histoire Naturelle de Lyon, **36**/1-2: 3-241, 249-441.
- PLAZIAT, J.-C. & MAHMOUDI, M. (1988): Trace fossils attributed to burrowing echinoids: a revision including new ichnogenus and ichnospecies. – Géobios, **21**/2: 209-233.
- PLEDGE, N.S. & SADLER, T. (1990): A new subspecies of the sea urchin *Peronella lesueuri* from the Quaternary of South Australia. – Transactions of the Royal Society of South Australia, **114**/2: 103-104.
- PORETSKAYA, E.S. (1972): [New Late Callovian *Pygurus* from Turkmenia]. – In: ZANINA, I.E. (ed.): [New species of fossil plants and invertebrates of USSR]. – pp. 291-292, Moskva (Akademia Nauka SSSR).
- (1974a): [Echinoidea]. – In: AKOPJANA, V.T. (ed.): [Atlas of the fossil fauna of the Armyanskog SSR]. – pp. 360-368, Erevan (Akademiya Nauk Armyanskog SSR).
- (1974b): [New echinoids from the Upper Oxfordian of the Kopetdag range and on the problem of classification and phylogeny of the disasterids]. – Voprosy Stratigrafii, **1**: 85-91.
- (1983): [*Polydiadema* (Echinoidea) from the Lower Valanginian of Mangyshlak]. – Paleontologicheskii Zhurnal, **1983**/3: 116-119.
- (1989): [New echinoid species *Codechinus prosorovskyi* from the Barremian of Turkmenistan]. – In: KALO, D.D., ROZHNOV, S.V., SOLOVJEV, A.N., STUKALINA, G.A., & KHINTS, I.M. (eds): Fossil and recent echinoderm researches. – pp. 161-165, Tallin (Academy of Sciences of the Estonian SSR, Institute of Geology).

- & LOBATSHEVA, S.V. (1988): [Section Echinodermata. Class Echinoidea]. – In: KRIMHOLZ, G.J. (ed.): [Berriasian of Mangyshlak]. – Akademiya Nauk SSSR, Mezhvedomstvennyi Stratigraficheskii Komitet SSSR, Trudy, **17**: 165-173.
- RADWAŃSKI, A. & WYSOCKA, A. (2001): Mass aggregation of Middle Miocene spine-coated echinoids Echinocardium and their integrated eco-taphonomy. – Acta Geologica Polonica, **51/4**: 295-316.
- RADWAŃSKA, U. (1996): A new Echinoid from the Eocene La Meseta Formation of Seymour Island, Antarctic Peninsula. – In: GAZDZICKI, A. (ed.): Palaeontological results of the Polish Antarctic Expedition, Part II. – Palaeontologia Polonica, **55**: 117-125.
- (1999): Lower Kimmeridgian echinoids of Poland. – Acta Geologica Polonica, **49/4**: 287-364.
- (2003): A monograph of the Polish Oxfordian echinoids; Part 1, Subclass Cidaroidea Claus, 1880. – Acta Palaeontologica Polonica, **53/2**: 143-165.
- REGUANT, S., ROMAN, J., & VILLATTE, J. (1970): Echinides de l'Eocene moyen de la region de Vic (Barcelone). – Bulletin de la Société Géologique de France, Série 7, **12/5**: 894-912.
- REICH, M., VILLIER, L., & KUTSCHER, M. (2004): The echinoderms of the Rügen White Chalk (Maastrichtian, Germany). – In: HEINZELLER, T. & NEBELSICK, J.H. (eds): Echinoderms: München. – Proc. 11th Internat. Echinoderm Conf., Munich/Germany/6-10 Oct. 2003. – pp. 495-501, Leiden/London/New York/Philadelphia/Singapore (A.A. Balkema Publishers (Taylor & Francis Grp.)).
- REY, J., TAJ-EDDINE, K., & WITAM, O. (1989): Échinides néocomiens de Safi (Maroc). – Géobios, **22/5**: 599-611.
- ROBERT, P. (1991): Notes sur les Echinides Oxfordiens de Druyes, 1ère partie. – In: Echinochroniques n° 1. – Bulletin de l'Association Echinologique, **1990**
- (1994): Révision des Echinides fossiles de l'Oxfordien moyen des environs de Druyes (Yonne) et notamment des moules internes rencontrés dans les Calcaires à Chailles. – iii+142 pp., Auxerre (Association Européenne et francophone pour le développement et la sauvegarde des recherches échinologiques).
- ROMAN, J. (1976): Echinides Eocenes et Miocenes du Qatar (Golfe Persique). – Annales de Paléontologie, **62/1**: 49-85.
- (1983): Échinides «Marsupiaux» (Fam. Temnopleuridae) dans le Néogène de l'Ouest Européen. – Annales de Paléontologie, **69/1**: 13-42.
- (1989): Échinoïdes Paléogènes du Bassin de Paris: aperçus nouveaux. – Actes du 114^e Congrès National des Sociétés Savantes, Section des Sciences, Paris, **114**: 293-304.
- (1991): *Trochodiadema? dhofarensis* nov. sp. (Echinoidea, Echinacea) de l'Albien du Sultanat d'Oman. – Annales de Paléontologie, **77/1**: 51-59.

- & STROUGO, A. (1988): Échinoids de l'étage Mokattamien (Éocène moyen et supérieur) de la région du Grand Caire (Égypte). – Bulletin du Museum National d'Histoire naturelle, Section C: Sciences de la Terre: Paleontologie, Geologie, Mineralogie, **10/2**: 137-161.
- & STROUGO, A. (1994): Echinoïdes du Libyen (Eocene Inferieur) d'Égypte. – Revue de Paléobiologie, **13/1**: 29-57.
- ROSSI, de G.E. & LEVY, R. (1989): Presencia de *Amplaster* n. sp. (Echinodermata, Clypeasteroidea) en el Terciario de Patagonia. – In: IV Congreso Argentino de Paleontología y Bioestratigrafía: Mendoza, Noviembre 23-27, 1986. Actas 4. – pp. 89-92, Mendoza.
- ROWE, F.W.E. (1989): Nine New Deep-water Species of Echinodermata from Norfolk Island and Wanganella Bank, northeastern Tasman Sea, with a Checklist of the Echinoderm Fauna. – Proceedings of the Linnean Society of New South Wales, **111/4**: 257-291.
- & HOGGETT, A.K. (1986): The cidarid echinoids (Echinodermata) of New South Wales. – Proceedings of the Linnean Society of New South Wales, **108 (1985)/4**: 225-261.
- SALAH, A.A. (1982): Die Temnocidariden (reg. Echiniden) der Maastricht-Stufe von NW-Deutschland. – Geologisches Jahrbuch, **A61**: 207-223.
- & SCHMID, F. (1982): Die Tylocidariden (reg. Echiniden) der Ober-Maastricht-Stufe von Dänemark und NW-Deutschland. – Geologisches Jahrbuch, **A61**: 177-205.
- SANTOLAYA, J.M. (1993): *Gualtieria*: un nuevo equinido del Terciario alicantino. – Cidaris – Revista Ilicitana de Paleontología y Mineralogía, **2/4**: 44-47.
- & SILLERO, C. (1994): Guía Ilustrada de los equinoideos fósiles de la Provincia de Alicante (I). – Cidaris – Revista Ilicitana de Paleontología y Mineralogía, **3/5**: 4-43.
- SARWAR, M. (1988): A New Fossil Echinoid from Eocene Rocks of Sindh Pakistan. – Pakistan Journal of Science, **39-40**: 75-83.
- (1989): Description of a new echinoid species of the genus *Hemipneustes*. – Pakistan Journal of Zoology, **21/4**: 375-380.
- SAUCÈDE, T., MOOI, R., & DAVID, B. (2007): Phylogeny and origin of Jurassic irregular echinoids (Echinodermata: Echinoidea). – Geological Magazine, **144/2**: 333-359.
- SAVCHINSKAYA, O.V. (1974): Tip Echinodermata. – In: KRYMGOLTZ, G.Y. (ed.): [Atlas of Upper Cretaceous Fauna of the Don Basin]. – pp. 303-333, Moscow (Nedra).
- (1987): Echinodermata. – In: SOBETSKIG, V.A., NEKHRIKOVA, N.I., BALAN, T.M., PLAMADY-ALA, G.S., MASLENNIKOVA, L.N., SAVCHINSKAYA, O.V., KOZBIMICHEVA, E.I., BEIBYAMOVSKIG, V.N., & VOLCHETOORSKIG, L.F. (eds): [Atlas of Invertebrate Fauna of the Late Cretaceous Sea of the Prekasian Basin]. – Trudy Paleontologiceskogo Instituta, **187**: 229-240.
- SCHLOSSER, M. (1923): Revision der Unteroligocänfauna von Häring und Reut im Winkel. – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beilagen Band, **47**: 254-294.

- SCHNEIDER, C.L., SPRINKLE, J., & RYDER, D. (2005): Pennsylvanian (Late Carboniferous) echinoids from the Winchell Formation, north-central Texas, USA. – *Journal of Paleontology*, **79**/4: 745-762.
- SCHULZ, M.-G. (1983): Die Echinodermen der Inoceramen-Mergel (Buntmergelserie, Ultrahelvetikum, Unter-Maastricht) des Moos-Grabens SE Siegsdorf (Oberbayern). – *Zitteliana*, **10**: 715-723.
- (1985): Die Evolution der Echiniden-Gattung *Galerites* im Campan und Maastricht Norddeutschlands. – *Geologisches Jahrbuch, Reihe A*, **80**: 3-93.
- SERAFY, D.K. (1971): A new species of *Clypeaster* (Echinodermata, Echinoidea) from San Felix Island, with a key to the recent species of the eastern Pacific Ocean. – *Pacific Science*, **25**/1: 165-170.
- (1974): A new heart-urchin, *Araeolampas atlantica*, new genus, new species, from the North Atlantic and notes on the closely related *Homolampas*. – *Proceedings of the Biological Society of Washington*, **87**/6: 41-48.
- SHIGEI, M. (1975a): A new species of the cidarid sea-urchins (Echinoidea: Cidaroida) from the East China Sea. – *Journal of the Faculty of Science, Tokyo University, Section 4, Zoology*, **13**/3: 321-328.
- (1975b): A new species of a new genus of the arbaciid sea-urchins (Echinoidea: Arbacioida) from the North Pacific, off Hachijo Island. – *Journal of the Faculty of Science, Tokyo University, Section 4, Zoology*, **13**/3: 329-332.
- (1975c): A new species of the heart urchins, an extant species of *Brissopatagus* (Echinoidea: Spatangoida), from Sagami Bay. – *Journal of the Faculty of Science, Tokyo University, Section 4, Zoology*, **13**/3: 333-339.
- (1977a): A new species of the echinothurioid sea-urchins, *Sperosoma nudum*, from the North Pacific, off central Japan. – *Journal of the Faculty of Science, Tokyo University, Section 4, Zoology*, **14**/1: 71-77.
- (1977b): A new species of the aspidodiadematoïd sea-urchins, *Aspidodiadema intermedium*, from the East China Sea, off southern Japan. – *Journal of the Faculty of Science, Tokyo University, Section 4, Zoology*, **14**/1: 79-84.
- (1981): A new species of the spatangoid sea-urchin (Echinoidea; Spatangoida), *Lovenia lata*, from Sagami Bay. – *Journal of the Faculty of Science University of Tokyo, Section IV Zoology*, **15**/1: 81-87.
- (1982): A new species of the fibulariid sea-urchin, *Fibularia japonica*, from Japanese waters. – *Publications of Seto Marine Biological Laboratory*, **27**/1-3: 11-16.
- SHIMANSKAYA, N.V. (1974): [A new Cenomanian species of the genus *Echinocorys* from the Crimea]. – *Paleontologicheskij Zhurnal*, **1974**/3: 149-152.
- SHMIDT, O.I. (1975): [On the new subgenus of the family Eoscutellidae DURHAM in the Paleogene of Kamchatka (Yuzhno Sakhalinsk)]. – *Trudy Sachalinskij Kompleksnyj Naučno-Issledovatel'skij Institut*, **36**: 29-31.

- (1976): [First notes on Echinodermata from Upper Cretaceous strata in the Soviet Arctic]. – Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (N.S.), **388**: 151-157.
- (1984): Chapter II. Sea urchins. – In: MENNER, V.V. (ed.): Atlas of fauna and flora of the Neogene sediments of the Far East. Tochilinski key section of western Kamchatka. – Trudy Geologicheskogo Instituta Akademiya Nauk SSSR, **385**: 147-150.
- , GORBATOV, A.M., & ZHELEZKO, V.I. (1979): [Sea urchins from Cenomanian deposits in Mangyshlak Mountains]. – In: MESEZHNIKOV, M.S. & KIRICHKOVA, A.I. (eds): [Stratigraphy of the Lower Cretaceous deposits of the petroleum regions of the USSR]. – Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (N.S.), 101-122.
- & SIMAKOV, S.N. (1953): [Late Cretaceous sea urchins from south-eastern Middle Asia]. – Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (N.S.), **66/1**: 5-75.
- & SINYELNIKOVA, V.N. (1971): [On sea urchins of Kavran series (western Kamchatka)]. – Doklady Akademii Nauk SSSR, **199/4**: 909-912.
- SILLERO, C. (2002): *Leiopedina molinai* sp. nov., un nueva equínido del Eoceno de Alicante. – Cidaris – Revista Ilicitana de Paleontología, **21-22**: 13-16.
- SILVA, G.H.d. (1959): Fósseis do Miocénico marinho da Ilha de Porto-Santo. – Memórias e Notícias, Coimra, **48**: 1-22.
- SMITH, A.B. (1994): Triassic echinoids from Peru. – Palaeontographica, Abteilung A (Palaeozoologie-Stratigraphie), **233/1-6**: 177-202.
- (1995): Late Campanian-Maastrichtian echinoids from the United Arab Emirates-Oman border region. – Bulletin of the Natural History Museum, Geology Series, **51/2**: 121-240.
- & BENGTON, P. (1991): Cretaceous echinoids from north-eastern Brazil. – Fossils and Strata, **31**: 1-88.
- & CRIMES, T.P. (1983): Trace fossils formed by heart urchins – a study of *Scolicia* and related traces. – Lethaia, **16**: 79-92.
- , GALLEMÍ, J., JEFFERY, C.H., ERNST, G., & WARD, P.D. (1999): Late Cretaceous-early Tertiary echinoids from northern Spain: implications for the Cretaceous-Tertiary extinction event. – Bulletin of the Natural History Museum, Geology Series, **55/2**: 81-137.
- & JEFFERY, C.H. (2000): Maastrichtian and Palaeocene echinoids: a key to world faunas. – Special Papers in Palaeontology, **63**: 1-406.
- & MCGUGAN, A. (1996): A new deep-water spatangoid echinoid from the Cretaceous of British Columbia, Canada. – Bulletin of the Natural History Museum, Geology Series, **52/2**: 103-107.

- & SAVILL, J.J. (2001): *Bromidechinus*, a new Ordovician echinozoan (Echinodermata), and its bearing on the early history of echinoids. – Transactions of the Royal Society of Edinburgh: Earth Sciences, **92/2**: 137-147.
- , SIMMONS, M.D., & RACEY, A. (1990): Cenomanian echinoids, larger foraminifera and calcareous algae from the Natih Formation, central Oman Mountains. – Cretaceous Research, **11/ 1**: 29-69.
- & WRIGHT, C.W. (1989): British Cretaceous echinoids. Part 1, General introduction and Cidaroida. – Monographs of the Palaeontographical Society, **141 (578)**: 1-101.
- & WRIGHT, C.W. (1990): British Cretaceous echinoids. Part 2, Echinothurioida, Diadematoidea and Stirodonta (1, Calycina). – Monographs of the Palaeontographical Society, **143 (583)**: 101-198.
- & WRIGHT, C.W. (1993): British Cretaceous echinoids. Part 3, Stirodonta 2 (Hemicidaroida, Arbacioidea and Phymosomatoida, Part 1). – Monographs of the Palaeontographical Society, **147 (593)**: 199-267.
- & WRIGHT, C.W. (1996): British Cretaceous echinoids. Part 4, Stirodonta 3 (Phymosomatidae, Pseudodiademmatidae) and Camarodonta. – Monographs of the Palaeontographical Society, **150 (602)**: 268-341.
- & WRIGHT, C.W. (1999): British Cretaceous echinoids. Part 5, Holoctypoida, Echinoneoida. – Monographs of the Palaeontographical Society, **153 (612)**: 342-390.
- & WRIGHT, C.W. (2000): British Cretaceous echinoids. Part 6, Neognathostomata (Casiduloids). – Monographs of the Palaeontographical Society, **154 (615)**: 391-429.
- & WRIGHT, C.W. (2003): British Cretaceous echinoids. Part 7, Atelostomata, 1. Holasteroida. – Monographs of the Palaeontographical Society, **156 (619)**: 440-568.
- & WRIGHT, C.W. (2008): British Cretaceous echinoids. Part 8, Atelostomata, 2. Spatangoida (1). – Monographs of the Palaeontographical Society, **162 (630)**: 569-635.
- SMUTS, W.J. (1988): A recently discovered clypeasteroid, *Echinodiscus colchesterensis* species novum, from the Alexandria Formation (Late Tertiary), South Africa. – Palaeontologia Africana, **26/8**: 93-98.
- SOLOVJEV, A.N. (1966): [Morphology and evolution Collyritidae and Disasteridae and the Systematics of sea urchins of the order Spatangoida]. – In: Proceedings of the 12th USSR Palaeontological Society, Leningrad. Pp. 45-46.
- (1971): [Late Jurassic and Early Cretaceous disasterid echinoids of the USSR]. – Trudy Paleontologiceskogo Instituta, **131**: 1-124.
- (1989): [New echinoid genus *Eoholaster* (Spatangoida, Holasteridae) from the Berriasian]. – In: KALO, D.D., ROZHOV, S.V., SOLOVJEV, A.N., STUKALINA, G.A., & KHINTS, I.M. (eds): Fossil and recent echinoderm researches. – pp. 148-155, Tallin (Academy of Sciences of the Estonian SSR, Institute of Geology).

- SQUIRES, R.L. & DEMETRION, R.A. (1992): Paleontology of the Eocene Bateque Formation, Baja California Sur, Mexico. – Contributions in Science, **434**: 1-55.
- & DEMETRION, R.A. (1993): A new species of the clypeasteroid echinoid *Astrodapsis* from the Miocene Isidro Formation, Baja California Sur, Mexico. – Journal of Paleontology, **67/2**: 258-263.
- & DEMETRION, R.A. (1994): A new species of the oligopygoid echinoid *Haimea* from the Lower Eocene of Baja California Sur, Mexico. – Journal of Paleontology, **68/4**: 846-851.
- & DEMETRION, R.A. (1995): A new genus of cassiduloid echinoid from the Lower Eocene of the Pacific Coast of Western North America and a new report of *Cassidulus ellipticus* KEW, 1920, from the Lower Eocene of Baja California Sur, Mexico. – Journal of Paleontology, **69/3**: 509-515.
- SRIVASTAVA, D.K. (1978): Microechinoids from the Oligo-Miocene rocks of Kutch, India. – In: RASHEED, D.A. (ed.): Proceedings of the VII Indian Colloquium on Micropalaeontology and Stratigraphy. – pp. 421-428, Madras (Dept. of Geology, University of Madras).
- (1981): Echinoid genus *Eupatagus* in Tertiary rocks of Kutch, India. – Journal of The Palaeontological Society of India, **25**: 38-41.
- (1982): A record of echinoid genus *Salenia* from the Middle Eocene rocks of southwestern Kutch, India. – Records of the Geological Survey of India, **112/7**: 23-25.
- (1988): Tertiary Echinoids of India – a Review. – Geological Survey of India Special Publication, **11/2**: 147-162.
- (2003): A new species of Cassiduloid echinoid *Gongrochanus* KIER, 1962 from the Late Cretaceous (Maastrichtian) rocks of the Ariyalur Area, Tamil Nadu, India. – Journal of the Palaeontological Society of India, **48**: 59-64.
- , GUPTA, A., & JAUHRI, A.K. (2008): A new regular echinoid from the Middle Eocene of Kachchh, Western India. – Journal of the Palaeontological Society of India, **53/1**: 107-110.
- & MATHUR, U.B. (1996): New fibulariid echinoid from the middle Eocene rocks of Rajasthan, India. – Journal of the Palaeontological Society of India, **41**: 53-56.
- , MISHRA, V.P., & SRIVASTAVA, M.C. (1992): *Echinolampas jigniensis*, a new species of cassiduloid echinoid from the Eocene rocks of Jammu Himalaya. – Geophytology, **20** (1990)/2: 98-102.
- , RANA, R.S., & SINGH, H. (2008): Record of *Megapneustes* GAUTHIER (Brissid echinoid) from the Khuiala Formation, Jaisalmer District, Rajasthan, India. – Journal of the Palaeontological Society of India, **53/1**: 31-36.
- , SINGH, A.P., TIWARI, R.P., & JAUHRI, A.K. (2008): Cassiduloids (Echinoidea) from the Siju Formation (late Lutetian-early Bartonian) of the South Garo Hills, Meghalaya, India. – Revue de Paléobiologie, **27/2**: 511-523.

- & SINGH, H. (2008): Brissid echinoid *Eupatagus* L. AGASSIZ, 1847 from the Khuiala Formation, Jaisalmer district, Rajasthan, India. – Earth Science India, **1/2**: 83-91.
- & SINGH, S.K. (1999): *Echinolampas* – a cassiduloid echinoid from the Tertiary rocks of Kachchh, India. – Journal of the Palaeontological Society of India, **44**: 25-47.
- & SINGH, S.K. (2001): Some species of *Amblypygus* AGASSIZ, 1840 and an indeterminate holectypoid echinoid from the Middle Eocene rocks of Kachchh (= Kutch), India. – Journal of the Palaeontological Society of India, **46**: 25-36.
- STILLER, F. (2001): Echinoid spines from the Anisian (Middle Triassic) of Qingyan, south-western China. – Palaeontology, **44/3**: 529-551.
- STOCKLEY, B., SMITH, A.B., LITTLEWOOD, T., LESSIOS, H.A., & MACKENZIE-DODDS, J.A. (2005): Phylogenetic relationships of spatangoid sea urchins (Echinoidea): taxon sampling density and congruence between morphological and molecular estimates. – Zoologica Scripta, **34/5**: 447-468.
- STOKES, R.B. (1975): Royaumes et provinces fauniques du Crétacé établis sur la base d'une étude systématique du genre *Micraster*. – Mémoires du Muséum National d'Histoire Naturelle, Nouvelle Série, Série C (Sciences de la Terre), **31**: 1-94.
- SZÖRÉNYI, E. (1952): Két új *Echinocyamus* faj a Dununántúli Eocénből. Deux nouvelles espèces du genre *Echinocyamus* de l'Éocène Transdanubien. – Földtani Közlöny, **82**: 289-293.
- (1955): Échinides Crétacés de la Bakony. – Geologica Hungarica, Series Palaeontologica, **26**: 1-332.
- TANAKA, K. (1984a): Hemiasterid echinoids from the Upper Cretaceous of Japan. – Transactions and Proceedings of the Palaeontological Society of Japan, New Series, **135**: 427-444.
- (1984b): Some Cretaceous Echinoids from Japan. – Bulletin of the Geological Survey of Japan, **35/5**: 189-202.
- , KANIE, Y., & OBATA, I. (1979): Maastrichtian and Danian echinoids from northwestern Madagascar. – Bulletin of the National Science Museum Series C (Geology & Paleontology), **5/ 2**: 25-50.
- & KOZAI, T. (1982): Some Cretaceous echinoids from the Monobe area, central Shikoku. – Transactions and Proceedings of the Palaeontological Society of Japan, New Series, **126**: 341-355.
- , NODA, M., & TANAKA, H. (1984): Echinoids from the Cretaceous Haidateyama Group, eastern Kyushu. – Transactions and Proceedings of the Palaeontological Society of Japan, New Series, **136**: 445-454.
- & OBATA, I. (1982): Selected echinoid fossils from the Miyako Group (Lower Cretaceous), northeast Honshu, Japan. – Bulletin of the National Science Museum, Series C (Geology & Paleontology), **8/3**: 117-143.
- TANDON, K.K. (1973): A new species of *Echinolampas* from Oligocene rocks of Kutch, India. – Proceedings of the Indian Science Congress, **60/3**: 195.

- & SRIVASTAVA, D.K. (1980): A new genus and species of the clypeasteroid echinoid from the middle Eocene rocks of Kutch, India. – *Journal of the Palaeontological Society of India*, **23-24**: 1-3.
- TATARENKO, D.E. & POLTARAUS, A.B. (1993): Affiliation of sea urchin, *Pseudocentrotus depressus*, to the family Strongylocentrotidae and description of a new genus *Mesocentrotus* belonging to this group based on DNA-DNA hybridization and comparative morphological data. – *Zoologicheskii Zhurnal*, **72/2**: 61-72.
- TESSIER, F. & ROMAN, J. (1973): Études paléontologiques et géologiques sur les Falaises de Fresco (Côte d'Ivoire). 6. Échinides. – *Annales de la Faculté des Sciences, Dakar*, **26**: 139-172.
- THIÉRY, P., LAMBERT, J., & COLLIGNON, M. (1928): Note sur quelques échinides de la région de Voulte. – *Travaux du Laboratoire de Géologie de la Faculté des Sciences de Lyon*, **13/Mémoire 11**: 81-103.
- TOULA, F. (1911): Die jungtertiäre Fauna von Gatun am Panamakanal. II. Theil. – *Jahrbuch der kaiserlich-königlichen Geologischen Reichsanstalt*, **61**: 487-530.
- TRAUTSCHOLD, H. (1859): Ueber Petrefacten vom Aralsee. – *Bulletin de la Société Impériale des Naturalistes de Moscou*, **32/2**: 303-322.
- TUOMEY, M. & HOLMES, F.S. (1855-1857): Pleiocene fossils of South Carolina containing descriptions and figures of the polyparia, echinodermata and mollusca. – i-xvi (1857); 1-30, pls. 1-8 (1855); 31-144, pls. 25-28 (1856); 145-152, pls. 29-30 (1857) pp., Charleston (James & Williams).
- TZANKOV, V. (1982): Les fossiles de Bulgarie. Va. Crétacé Supérieur: Cephalopoda (Nautiloidea, Ammonoidea) et Echinodermata (Echinoidea). – 136 pp., Sofia (Académie Bulgare des Sciences).
- VADET, A. (1988): Révision des principaux «*Cidaris*» de l'Oxfordien et du Kimméridgien inférieur Européens. – *Mémoires de la Société Académique du Boulonnais*, **4**: 1-148.
- (1991): Révision des «*Cidaris*» du Lias et du Dogger Européens. – *Mémoires de la Société Académique du Boulonnais*, **10**: 1-175.
- (1993): Les Oursins du Bathonien et du Bajocien de Normandie. – *Mémoires de la Société Académique du Boulonnais*, **6/3-4**: 1-104.
- (1995): *Cottaldia paquettei* nov. sp. un *Cottaldia* nouveau dans les calcaires à spongiaires du Poitou. – *Mémoires de la Société Académique du Boulonnais*, **16**: 114-115.
- (1997): Échinides fossiles du Callovien de la Sarthe et des confins de l'Orne. II-A: Les échinides irréguliers: Analyse et évolution. – *Mémoires de la Société Académique du Boulonnais*, **18**: 1-140.
- (1999a): La Classification des oursins réguliers du Lias et du Jura. – *Bulletin de la Société Académique du Boulonnais*, **4/3**: 70-83.
- (1999b): Les Echinides de la Sarthe: supplément. – *Bulletin de la Société Académique du Boulonnais*, **4/3**: 84-92.

- (1999c): Révision des Échinides de Saint Cassian. – Mémoires de la Société Académique du Boulonnais, **20**/1-2: 1-116.
- (2000): Un Salenioida primitif dans le Kimmeridgien de l’Ain. – Bulletin de la Société Académique du Boulonnais, **4**/4: 93-97.
- (2003): Echinides fossiles du Jurassique du Boulonnais. Première partie: les«*Cidaris*» [also cited as: “Echinides fossiles du Boulonnais. Du Bajocien au Tithonien: les«*Cidaris*».” – Title on front cover differs from title inside]. – Annales de la Société d’Histoire Naturelle du Boulonnais, **3**/2: 2-35.
- (2005a): Echinides fossiles du Boulonnais: Du Bajocien au Tithonien. Echinothurioidea, Pedinoida, Orthopsida, Hemicidaroida. – Annales de la Société d’Histoire Naturelle du Boulonnais, **4**/1: 38-108.
- (2005b): Echinides fossiles du Boulonnais: Du Bajocien au Tithonien. Phymosomatoida, Stomechinoida, Arbacioida. – Annales de la Société d’Histoire Naturelle du Boulonnais, **4**/2: 110-156.
- (2007): Echinides fossiles du Boulonnais: Du Bajocien au Tithonien. Les irréguliers. – Annales de la Société d’Histoire Naturelle du Boulonnais, **5**/2: 157-220.
- & CHESNIER, M. (2003): Les echinides fossiles de Normandie du Sinémurien au Bajocien inférieur. – Annales de la Société d’Histoire Naturelle du Boulonnais, **3**/1: 3-48.
- , EDERLÉ, R., & ROBERT, P. (1995): Les oursins du Bathonien de l’Yonne. – Mémoires de la Société Académique du Boulonnais, **16**: 99-107.
- , MARNIGNAC, C., NICOLLEAU, P., & REBOUL, R. (2007): Les echinides Berriasso – Valanginiens de la falaise de Safi (Maroc). [also cited as: «Les Echinides de Safi (Maroc).» – Title on front cover; differs from title inside]. – Annales de la Société d’Histoire Naturelle du Boulonnais, **6**/2: 1-42.
- & NICOLLEAU, P. (1995): *Pygopyrina thibaudi* sp. nov. de l’Oxfordien de Dompcevrin. – Mémoires de la Société Académique du Boulonnais, **16**: 8-88.
- & NICOLLEAU, P. (2000): Un *Acrocidaris* nouveau dans le Callovien du Portugal. – Bulletin de la Société Académique du Boulonnais, **4**/4: 107-109.
- & NICOLLEAU, P. (2002): Un *Gymnocidaris* nouveaux dans le Kimmeridgien supérieur de la Région d’Angoulême (Charente). – In: Echinides nouveaux peu ou mal connus Tome II fascicule 3. – Annales de la Société d’Histoire Naturelle du Boulonnais, **2**/1: 3-7.
- & NICOLLEAU, P. (2005): Echinides nouveaux peu ou mal connus. Tome II, fascicule 5. Les oursins domériens de la région de Rich (Maroc). – Annales de la Société d’Histoire Naturelle du Boulonnais, **5**/1: 1-36.
- , NICOLLEAU, P., & PINEAU, J.-P. (1996): Échinides fossiles du Callovien de la Sarthe et des confins de l’Orne. I: Les échinides réguliers. – Mémoires de la Société Académique du Boulonnais, **17**: 1-184.

- , NICOLLEAU, P., & PINEAU, J.-P. (1998): Échinides fossiles du Callovien de la Sarthe et de l'Orne. II-B: Les échinides irréguliers: Description. – Mémoires de la Société Académique du Boulonnais, **19**: 1-128.
- , NICOLLEAU, P., & REBOUL, R. (2008): Echinides nouveaux peu ou mal connus. Tome II, fascicule 6. *Masrouraster* nov. gen. du Domérien de l'Atlas Marocain. – Annales de la Société d'Histoire Naturelle du Boulonnais, **7/2**: 15-17.
- , NICOLLEAU, P., & RIGOLLET, L. (2002): Les Échinides fossiles du Kimmeridgien des récifs coralliens de la Pointe du Chay (près de La Rochelle, Charente Maritime, France). – Annales de la Société d'Histoire Naturelle du Boulonnais, **2/2**: 1-71.
- & RÉMY, N. (2004): Echinides nouveaux peu ou mal connus. Tome II, fascicule 4. Les oursins du Bajocien des Ardennes. – Annales de la Société d'Histoire Naturelle du Boulonnais, **3/3**: 1-20.
- & SLOWIK, D. (2001): Les oursins Bajocien de Liocourt. – Mémoires de la Société Académique du Boulonnais, Série Histoire Naturelle, **22/1**: 1-48.
- & WILLE, E. (2002): Quelques oursins du Lusitanien du Portugal. – In: Echinides nouveaux peu ou mal connus Tome II fascicule 3. – Annales de la Société d'Histoire Naturelle du Boulonnais, **2/1**: 8-18.
- VALDINUCCI, A. (1974): «*Aliaster*» nuovo genere di echinide istituito per un piccolo gruppo di specie tratte dai generi *Trachyaster* ed *Opissaster* (Famiglia Hemiasteridae Clark). – Bolletino del Servizio Geologico d'Italia, **94** (1973)/2: 454-479.
- VEIGA FERREIRA, O. da (1962): Nota sobre a presença do género «*Agassizia*» no Miocénico do Algarve. – Comunicações dos Serviços Geológicos de Portugal, **16**: 293-295.
- VENZO, S. (1934): Nuove forme del cattiano di monte Brione. – Studi Trentini di Scienze naturali, Annata, **15/2-3**: 103-116.
- VIALOV, O.S. (1945): [New variety of *Collyrites analis* DESM. from the Balkhan Jurassic]. – Doklady Akademii Nauk SSSR, **50**: 467-469.
- WANG, C.-C. (1982a): On the early Miocene sand dollar *Echinodiscus yeliuensis* n.sp. from the Talio Formation of Yeliu, northern Taiwan. – Proceedings of the Geological Society of China, **25**: 150-157.
- (1982b): On the fossil laganid echinoids from Taiwan with a discussion of the “genus” *Peronellites* Hayasaka & Morishita. – Acta Geologica Taiwanica, **21**: 140-156.
- (1983): A new species of *Astriclypeus* from the Wuchihshan Formation near Chilung, Taiwan. – Bulletin of the Central Geological Survey, **2**: 113-120.
- (1984a): Fossil *Echinodiscus* from Taiwan. – Bulletin of the Central Geological Survey, **3**: 107-115.
- (1984b): New classification of clypeasteroid echinoids. – Proceedings of the Geological Society of China, **27**: 119-152.

- (1984c): A new species of *Pericosmus* (Echinoidea) from the Miocene Nankang Sandstone of northern Taiwan. – *Jingjibu Zhongyang Dizhi Diaochasuo Tekan* = Special Publication of the Central Geological Survey, **3**: 249-256.
- (1985): Fossil clypeasterid echinoids from Taiwan. – *Proceedings of the Geological Society of China*, **28**: 143-176.
- (1986): Fossil astriclypeid echinoids from Taiwan. – *Proceedings of the Geological Society of China*, **29**: 149-183.
- (1994): Phylogenetic analysis of the fossulasterid echinoids (Order Clypeasteroida) from Australia. – *Journal of the Geological Society of China*, **37/2**: 225-246.
- WEBSTER, G.D. (1997): Lower Carboniferous echinoderms from northern Utah and western Wyoming. – *Bulletin Utah Geological Survey*, **128**/Paleontology Series Vol. 1: 1-65.
- WEINBERG, S. & DE RIDDER, C. (1998): *Asthenosoma marisrubri* n. sp. (Echinodermata, Echinoidea) from the Red Sea. – *Beaufortia*, **48/3**: 27-46.
- WIESBAUR, J. (1874): Fossile Pflanzen im marinen Tertiär Conglomerate zu Kalksburg bei Wien. – *Verhandlungen der kaiserlich-königlichen Geologischen Reichsanstalt*, **8/7**: 157-165.
- WOODRING, W.P. (1957): Geology and Paleontology of Canal Zone and adjoining Parts of Panama. – *Geology and Description of Tertiary Mollusks (Gastropods: Trochidae to Turritellidae)*. – *Geological Survey Professional Papers*, **306-A**: 1-145.
- YANG SHENGQIU (1991): [Late Cretaceous and Early Tertiary Echinoids from Tarim Basin]. – In: PAN HUAZHANG, YANG SHENGQIU, & SUN DONGLI (eds): [Late Cretaceous and Early Tertiary Gastropoda, Echinoids and Brachiopoda from Tarim Basin]. – pp. 101-132, Beijing (Science Press).
- ZACHOS, L.G. & MOLINEUX, A. (2003): Eocene echinoids of Texas. – *Journal of Paleontology*, **77/3**: 491-508.
- ZARDINI, R. (1973): Fossili di Cortina: Atlante degli echinodermi cassiani (Trias Medio-superiore) della regione dolomitica attorno a Cortina d'Ampezzo. – 29 pp., Cortina d'Ampezzo (Foto Ghedina).
- ŽITĚ, J. (1986): *Magnosia* Michelin and *Magnosiopsis* gen. n., (Echinoidea) from the Lower Cretaceous of Štramberk (Czechoslovakia). – *Časopis pro mineralogii a geologii*, **31/ 4**: 371-385.
- (1994): Some oligopygoids of the genus *Haimea* MICHELIN (Echinoidea) from the Eocene of Cuba. – *Acta Musei Nationalis Pragae*, **49B (1993)**/1-4: 1-40.
- & GEYS, J.F. (2003): *Novasalenia* gen. nov.: a remarkable Late Cenomanian echinoid from the Bohemian Cretaceous Basin (Czech Republic). – *Cretaceous Research*, **24/1**: 23-30.

Index

<i>Aarchaeocidaris</i>	209	<i>Adelopneustes ernsti</i>	282
<i>Aarchaeocidaris strawberryensis</i>	209	<i>Adelopneustes montainvillensis</i>	285
<i>abatoides, Pericosmus</i>	375	<i>adrianae, Miocidaris</i>	210, 276
<i>abatoides, Schizocosmus</i>	375	<i>adunca, Cidaris</i>	221
<i>Abatus</i>	366	<i>aegyptiaca, Scutella</i>	296
<i>Abatus kieri</i>	366	Aeropsidae	376
<i>Abelleicidaris</i>	227, 228	<i>affinidouvillei, Iranianster</i>	350
<i>Abelleicidaris borjnadorensis</i>	226, 228	<i>affinimorgani, Iranianster</i>	350
<i>Abelleicidaris bradfordensis</i>	228	<i>africana, Linthia</i>	370
<i>Abelleicidaris saemanni</i>	228	<i>Agassizia</i>	367
<i>Abertella</i>	305	<i>Agassizia algarbiensis</i>	367
<i>Abertella complanata</i>	305	<i>Agassizia eugeniae</i>	367
<i>Abertella gualichensis</i>	305	<i>Agassizia powersi</i>	367
Abertellidae	305	<i>Agassizia (Anisaster)</i>	367
<i>abiquaensis, Arbacia</i>	263	<i>Agassizia (Anisaster) arabica</i>	367
<i>abyssale, Kamptosoma</i>	235	<i>Agassizia (Anisaster) wilmingtonica</i>	
<i>Acanthocidaris</i>	225	<i>inflata</i>	367
<i>Acanthotrema</i>	393	<i>aggtelekensis, Cidaris</i>	221
<i>Acanthotrema siculum</i>	393, 394	<i>Aguayoaster</i>	367
<i>Acriaster</i>	323	<i>Aguayoaster schickleri</i>	367
<i>Acriaster sergipensis</i>	323	<i>aialensis, Cidaris</i>	218
<i>Acrocidaris</i>	251	<i>aialensis fusiformis, Cidaris</i>	218
<i>Acrocidaris marocense</i>	251	<i>aichinoi, Aliaster</i>	361
<i>Acrocidaris nicolleau</i>	252	<i>aichinoi, Trachyaster</i>	361
Acroechinoidea	233	<i>akdjarensis, Conulus</i>	282
<i>Acropeltis</i>	263	<i>akkajensis, Neoglobator</i>	284
<i>Acropeltis atlantica</i>	263	<i>akkajensis, Protobrissus</i>	380
<i>Acrosalenia</i>	242	<i>akkajensis, Pseudogibbaster</i>	380
<i>Acrosalenia arabica</i>	242	<i>akkaptschigensis, Parahemiaster</i>	362
<i>Acrosalenia bowersi</i>	242	<i>aksuensis, Pygaulus</i>	313
<i>Acrosalenia chartroni</i>	233	<i>aktaschensis, Hemiaster</i>	356
<i>Acrosalenia dhrumaensis</i>	242	<i>alanbakeri, Caenopedina</i>	239
<i>Acrosalenia marcoui</i>	243	<i>alata brevicaulis, Cidaris</i>	218
<i>Acrosalenia marratensis</i>	242	<i>alata canaliculata, Cidaris</i>	218
Acrosaleniiidae	242	<i>alatus, Amplaster</i>	301
Acrosalenioidea	241	<i>alatus, Lunulaster</i>	301
<i>Actapericulum</i>	395	<i>albeari, Antillaster</i>	393
<i>Actapericulum bicarinatum</i>	395	<i>Albertechinus</i>	206
<i>aculeata, Cidaris</i>	218	<i>Albertechinus devonicus</i>	206
<i>aculeata ailaensis, Cidaris</i>	218	<i>alexandri forma arctambulacrum,</i>	
<i>acuminata, Echinolampas</i>	316	<i>Echinolampas</i>	316
<i>acutus, Gongrochanus?</i>	320	<i>algarbiensis, Agassizia</i>	367
<i>acutus, Pygorhynchus</i>	314	<i>Aliaster</i>	361
<i>Adelopneustes</i>	282, 284	<i>Aliaster aichinoi</i>	361

<i>Aliaster almerai</i>	361	<i>Amphiope bioculata</i>	304
<i>Aliaster cotteaudi</i>	361	<i>Amphiope caupianensis</i>	303
<i>Aliaster cotteri</i>	361	<i>Amphiope miocenica</i>	303
<i>Aliaster jourdyi</i>	361	<i>Amphipneustes</i>	367, 376
<i>Aliaster lovisatoi</i>	361	<i>Amphipneustes davidi</i>	367
<i>alicantina, Gualtieria</i>	385	<i>Amphipneustes mironovi</i>	368
<i>alievi, Orthaster</i>	364	<i>amphora, Echinosisgra</i>	347
<i>Allomma</i>	255	<i>amphora amphora, Echinogutta</i>	349
<i>Allomma kalon</i>	255	<i>amphora amphora, Echinosisgra</i>	347
<i>Allomma wrighti</i>	255	<i>amphora antarctica, Echinosisgra</i>	348
<i>almerai, Aliaster</i>	361	<i>amphora fabrefacta, Echinosisgra</i>	348
<i>almerai, Opissaster</i>	361	<i>amphora indica, Echinogutta</i>	349
<i>Almucidaris</i>	216	<i>amphora indica, Echinosisgra</i>	348
<i>Almucidaris durhami</i>	216	<i>amphora valvaedentata, Echinosisgra</i>	348
<i>alphonsei, Schizaster</i>	372	<i>Amplaster</i>	301
<i>alpina, Cidaris</i>	218	<i>Amplaster alatus</i>	301
<i>alpina, Pseudanachys</i>	341	<i>Amplaster coloniensis</i>	301
<i>alta, Brissopsis</i>	366	<i>Amplaster ellipticus</i>	301
<i>alta, Eopericosmus</i>	366	<i>ampliflorus, Amphidetus</i>	391
<i>alternans, Rolliericidaris</i>	227	<i>amplus, Leioechinus</i>	260
<i>alternus, Gauthieria</i>	256	<i>amudariensis, Hemiaster</i>	356
<i>alternus, Hemithylus</i>	256	<i> analis dimota, Collyrites</i>	332
<i>altus, Angusticlypeus</i>	308	<i>Anaulcidaris</i>	232
<i>altus, Catopygus</i>	311	<i>ancileformis, Echinocorys</i>	335
<i>altus, Clypeus</i>	308	<i>andinus, Pygurus</i>	310
<i>altus, Plesiechinus</i>	241	<i>angelae, Baueria</i>	263
<i>ambiguum, Polydiadema</i>	253	<i>angolanus, Leiostomaster</i>	364
<i>Ambipleurus</i>	267	<i>anguillensis, Tretocidaris</i>	224
<i>Ambipleurus viladensis</i>	267	<i>angulatus, Conulus</i>	282
<i>Ambipleurus? quaylei</i>	267	<i>angulosus, Epiaster</i>	352
<i>Amblypygidae</i>	286	<i>Angusticlypeus</i>	308
<i>Amblypygoida</i>	286	<i>Angusticlypeus altus</i>	308
<i>Amblypygus</i>	278, 286	<i>Angusticlypeus angustiporus</i>	308
<i>Amblypygus moriensi</i>	279	<i>Angusticlypeus babeau</i>	308
<i>Amblypygus moriensis</i>	279	<i>Angusticlypeus davoustianus</i>	308
<i>Amblypygus ratiparensis</i>	279	<i>Angusticlypeus mulleri</i>	308
<i>Amblypygus (Paramblypygus)</i>		<i>Angusticlypeus rathieri</i>	308
<i>houphoueti</i>	279	<i>angustiporus, Angusticlypeus</i>	308
<i>ammonitorum, Salenia</i>	245	<i>angustiporus, Clypeus</i>	308
<i>amnicus, Micraster</i>	378	<i>angustisulcus, Pliotoxaster</i>	355
<i>Amoraster</i>	380	<i>angustus, Metalia</i>	388
<i>Amoraster paucituberculata</i>	380	<i>Anisaster</i>	367
<i>Amoraster tuberculata</i>	380	<i>Anisocidaridae</i>	208
<i>ampezzana, Miocidaris</i>	210	<i>Anisocidaris</i>	208, 229
<i>Amphidetus</i>	391	<i>Anisocidaris leberti</i>	218
<i>Amphidetus ampliflorus</i>	391	<i>anisopetalus, Hemiaster</i>	356
<i>Amphiope</i>	303, 304	<i>annae, Parapygus</i>	319

<i>annae</i> , <i>Rostropygus</i>	319	<i>arabicus</i> , <i>Polycyphus</i>	261
<i>annosus</i> , <i>Pericosmus</i>	365	<i>arabicus</i> , <i>Pygurus</i>	310
Anorthopygidae	281	<i>arachnius</i> , <i>Hemiaster</i>	356
<i>Anorthopygus</i>	281	<i>Araeolampas</i>	390
<i>Anorthopygus arabicus</i>	281	<i>Araeolampas atlantica</i>	390
<i>antarctica</i> , <i>Huttonechinus</i>	344	<i>Araeolampas fulva</i>	390
<i>antartica</i> , <i>Echinogutta</i>	349	<i>Arbacia</i>	263
<i>Antillaster</i>	393	<i>Arbacia abiquaensis</i>	263
<i>Antillaster albeari</i>	393	Arbaciidae	263
<i>Antillaster bagmanovi</i>	393	<i>Arbacina</i>	268
<i>antipodeanus</i> , <i>Urechinus</i>	345	<i>Arbacina emmae</i>	268
<i>antiqua</i> , <i>Pedina</i>	240	<i>Arbacina hugueti</i>	268
<i>antiqua</i> , <i>Thieulinipedina</i>	240	<i>Arbacina pareyni</i>	268
<i>Antiquibrissus</i>	380	Arbacioida	262
<i>Antiquibrissus suemegensis</i>	380, 381	<i>Arbia</i>	267
<i>Antiquibrissus sümegensis</i>	381	Archaeocidaridae	209
<i>Antrechinus</i>	344	<i>Archaeocidaris</i>	209
<i>Antrechinus drygalskii</i>	344	<i>Archaeocidaris apheles</i>	209
<i>Antrechinus mortenseni</i>	344, 345	<i>Archaeocidaris brownwoodensi</i>	209
<i>Antrechinus nordenskjoldi</i>	344	<i>Archaeocidaris faassi</i>	210
<i>antunesi</i> , <i>Echinolampas</i>	316	<i>Archaeocidaris hamata</i>	209
<i>aoteanus</i> , <i>Urechinus</i>	345	<i>Archaeocidaris hemispinifera</i>	210
<i>aotearoa</i> , <i>Ctenocidaris</i>	212	<i>Archaeocidaris schrëteri</i>	209
Apatopygidae	330	<i>Archaeocidaris whatleyensis</i>	210
<i>Apatopygus</i>	330	<i>Archaeocidaris? diadematoïdes</i>	209
<i>Apatopygus gaudensis</i>	330	<i>archiaci complanatus</i> , <i>Schizaster</i>	373
<i>Apatopygus mannumensis</i>	330	<i>Archiacia</i>	323
<i>Apatopygus vincentinus</i>	330	<i>Archiacia hungarica</i>	323
<i>Aphanophora</i>	395	<i>Archiacia magna</i>	323
<i>Aphanophora? bassoris</i>	395	Archiaciidae	323
<i>apheles</i> , <i>Archaeocidaris</i>	209	<i>arcticus</i> , <i>Hemiaster</i>	356
<i>apokryphos</i> , <i>Psephoaster</i>	363	<i>arcus</i> , <i>Conulus</i>	282
<i>Aporocidaris</i>	211	<i>argentinensis</i> , <i>Bolbaster</i>	359
<i>Aporocidaris eltania</i>	211	<i>ariyalurensis</i> , <i>Gongrochanus</i>	320
<i>Aporocidaris usarpi</i>	211	<i>ariyalurensis</i> , <i>Limpasiaster</i>	324
<i>Apoxyptalum</i>	381	<i>ariyalurensis</i> , <i>Progongrochanus</i>	322
<i>Apoxyptalum chenjafra</i>	381	<i>Arnaudaster</i>	315
<i>Aptilechinus</i>	206	<i>Arnaudaster cylindriiformis</i>	315
<i>Aptilechinus caledonensis</i>	206	<i>arnonensis</i> , <i>Mecaster</i>	360
<i>arabica</i> , <i>Acrosalenia</i>	242	<i>arumaensis</i> , <i>Rhynchopygus</i>	327
<i>arabica</i> , <i>Agassizia (Anisaster)</i>	367	<i>ashrafi</i> , <i>Schizaster (Paraster)</i>	374
<i>arabica</i> , <i>Bothryopneustes</i>	309	<i>asiatica</i> , <i>Remondella</i>	300
<i>arabica</i> , <i>Leptosalenia</i>	246	<i>aslaniani</i> , <i>Ditremaster</i>	361
<i>arabica</i> , <i>Salenia</i>	246	<i>asperula</i> , <i>Cidaris</i>	226
<i>arabicus</i> , <i>Anorthopygus</i>	281	<i>asperula</i> , <i>Oedematocidaris</i>	226
<i>arabicus</i> , <i>Goniopygus</i>	265	<i>Aspidodiadema</i>	237
<i>arabicus</i> , <i>Hemipneustes</i>	338	<i>Aspidodiadema intermedium</i>	237

<i>Aspidodiadema montanum</i>	237	<i>Bahariya</i>	391
<i>Aspidodiadema sinuosum</i>	237	<i>Bahariya teetotumensi</i>	391
Aspidodiadematidae	237	<i>bahiensis, Holosalenia</i>	248
<i>Asterostoma</i>	393	<i>bajarunasi, Hypsopygaster</i>	324
<i>Asterostoma pawsoni</i>	393	<i>bajarunasi, Neoproraster</i>	371
Asterostomatidae	393	<i>bajarunasi, Toxaster</i>	355
Asterostomatina	393	<i>bajasurensis, Haimea</i>	285
<i>Asthenosoma</i>	234	<i>bajasurensis, odapsis</i>	298
<i>Asthenosoma marisrubis</i>	234	<i>bajsarensis, Linthia</i>	370
Astriclypeidae	291, 303	<i>bajulus, Pachycentrotus</i>	275
<i>Astriclypeus</i>	303	<i>bakeri, Notocidaris</i>	212
<i>Astriclypeus elegans</i>	303	<i>bakonyense, Pseudodiadema</i>	253
<i>Astriclypeus miaoliensis</i>	303	<i>Balanocidaris</i>	216
<i>Astriclypeus pitouensis</i>	303	<i>Balanocidaris clubis</i>	216
<i>Astriclypeus waiwulunensis</i>	304	<i>Balanocidaris fusiformis</i>	217
<i>Astriclypeus yeliuensis</i>	303	<i>balcanicus, Echinocorys</i>	335
<i>Astrodapsis</i>	298	<i>baldwini, Lytechinus</i>	272
<i>Astrodapsis bajasurensis</i>	298	<i>balli, Desoricidaris</i>	214
<i>Astrodapsis ilyinensis</i>	299	<i>balli, Leiocidaris</i>	214
<i>astrodapsoides, Echinarachnius</i>	299	<i>baltica, Cidaris</i>	227
Atelostomata	307	<i>baltica, Oedematocidaris</i>	227
<i>atlantica, Acropeltis</i>	263	<i>banatica, Salenia</i>	245
<i>atlantica, Araeolampas</i>	390	<i>bandeli, Hemicidaris</i>	249
<i>atlasensis, Diplopodia</i>	252	<i>Bandelicyphus</i>	268
Atopechinae	267	<i>Bandelicyphus genaensis</i>	268
Atopechinaeæ	267	<i>bangtoupensis, Cidaris</i>	221
Atopechinidae	262	<i>baroini, Culozoma</i>	238
<i>Atopechinus</i>	267	<i>Baronechinus</i>	259
<i>australis, Brissopsis</i>	382	<i>Baronechinus baroni</i>	259
<i>australis, Echinocorys</i>	335	<i>baroni, Baronechinus</i>	259
<i>australis, Nordenskjöldaster?</i>	351	<i>bashibulakensis, Nucleopygus</i>	325
<i>Austrocidaris</i>	212	<i>basiprocta, Plesiocorys</i>	340
<i>Austrocidaris pawsoni</i>	212	<i>Basseaster</i>	351
<i>Austrocidaris seymourensis</i>	212	<i>bassoris, Aphanophora?</i>	395
<i>avenionensis nerthensis, Cyathocidaris</i>	222	<i>bastai, Echinolampas</i>	317
<i>azemati, Cyclaster</i>	383	<i>batequensis, Eupatagus</i>	384
<i>azemati, Pygomalus</i>	308	<i>batheri, Cidaris</i>	221
<i>azemati, Pyrina</i>	283	<i>Batheritiaris</i>	275, 276
<i>babatakensis, Isomicraster</i>	377	<i>Batheritiaris incipiens</i>	276
<i>babeaui, Angusticlypeus</i>	308	<i>batnensis, Holosalenia</i>	248
<i>babeaui, Clypeus</i>	308	<i>batnensis, Salenia</i>	248
<i>baconicus, Botriopygus</i>	314	<i>Baudhuinipygus</i>	311
<i>baconicus, Epiaster</i>	352	<i>Baudhuinipygus brodiei</i>	311
<i>baconicus, Hemiaster</i>	358	<i>Baudhuinipygus haimei</i>	311
<i>baconicus, Pseudholaster</i>	341	<i>Baueria</i>	263
<i>bagheerae, Lovenia</i>	392	<i>Baueria angelae</i>	263
<i>bagmanovi, Antillaster</i>	393	<i>baumi, Eurhodia</i>	328

<i>baylissi</i> , <i>Salenia</i>	245	<i>bosei</i> , <i>Polydiadema</i>	253
<i>beatriceae</i> , <i>Tripylus</i>	375	Bothriocidaridae	204
<i>bedakensis</i> , <i>Hemiaster</i>	356	<i>Bothriocidaris</i>	204
<i>beineuensis</i> , <i>Pygurus</i>	310	<i>Bothriocidaris kolatai</i>	204
<i>bella</i> , <i>Salenia</i>	245	<i>Bothriocidaris maquoketensis</i>	204
<i>bella parva</i> , <i>Salenia</i>	245	<i>Bothriocidaris solemi</i>	204
<i>bellissae</i> , <i>Giraliaster</i>	337	<i>Bothriocidaris vulcani</i>	204
<i>bellus</i> , <i>Progongrochanus</i>	322	Bothriocidaroida	204
<i>belyaevi</i> , <i>Pilematechinus</i>	344	<i>Bothryopneustes</i>	308
<i>bermotensis</i> , <i>Kutchypygyus</i>	287	<i>Bothryopneustes arabica</i>	309
<i>bermudezi</i> , <i>Cidaris</i>	217	<i>Bothryopneustes dhurmaensis</i>	309
<i>bermudezi</i> , <i>Tylocidaris</i>	226	<i>Bothryopneustes inflata</i>	308
<i>bernasconiae</i> , <i>Sterechinus</i>	274	<i>Bothryopneustes kauffmani</i>	308
<i>biaense</i> , <i>Echinocardium</i>	392	<i>Botriopygus</i>	314
<i>biannulatus</i> , <i>Murraypneustes</i>	393	<i>Botriopygus baconicus</i>	314
<i>bicarinatum</i> , <i>Actapericulum</i>	395	<i>Botriopygus latipetalus</i>	314
<i>Bichordites</i>	396	<i>Botriopygus neoataxensis</i>	314
<i>Bichordites monastiriensis</i>	396	<i>Botriopygus ovalis</i>	314
<i>biconica</i> , <i>Cidaris</i>	218	<i>Botriopygus pappi</i>	314
<i>bigoti</i> , <i>Dicoptella</i>	271	<i>Botriopygus pappi elongatus</i>	315
<i>bigoti</i> , <i>Guittonicidaris</i>	230	<i>Botriopygus pappi kutassyi</i>	315
<i>bigoti</i> , <i>Rhabdocidaris</i>	230	<i>Botriopygus petalodes planus</i>	315
<i>bigoti</i> , <i>Viaudechinus</i>	271	<i>Botriopygus subquadratus</i>	315
<i>bioculata</i> , <i>Amphiope</i>	304	<i>Botriopygus suemegensis</i> [<i>sümegeensis</i>]	315
<i>bipuncata</i> , <i>Diplopodia</i>	252	<i>Botriopygus variabilis</i>	315
<i>bipuncata</i> , <i>Girardema</i>	252	<i>boutini</i> , <i>Gymnocidaris</i>	222
<i>bischoffi</i> , <i>Crassiholaster</i>	334	<i>bowersi</i> , <i>Acrosalenia</i>	242
<i>bischoffi</i> , <i>Holaster</i>	334	<i>bowersi</i> , <i>Iranianster</i>	350
<i>blackdownensis</i> , <i>Pseudholaster</i>	341	<i>Brachysternaster</i>	368
<i>blackmorei anterior</i> , <i>Hagenowia</i>	337	<i>Brachysternaster chesheri</i>	368
<i>blainvillei</i> , <i>Philicidaris</i>	229	<i>bradfordensis</i> , <i>Abelleicidaris</i>	228
<i>blainvillei</i> , <i>Rhabdocidaris</i>	229	<i>bradfordensis</i> , <i>Cidaris</i>	228
<i>Blaviaster</i>	385	<i>Brangema</i>	238
<i>blumenbachii</i> , <i>Cidarites</i>	227	<i>Brangema brangeri</i>	238
<i>blumenbachii</i> , <i>Nenoticidaris</i>	227	<i>brangeri</i> , <i>Brangema</i>	238
<i>bobkovae</i> , <i>Parahemiaster</i>	362	<i>brangeri</i> , <i>Chesniericidaris</i>	217
<i>Bolbaster</i>	358	Braunechinidae	276
<i>Bolbaster argentinensis</i>	359	<i>Braunechinus</i>	277
<i>Bolbaster compressus</i>	359	<i>Braunechinus dorsata</i>	277
<i>Boletechinus</i>	262, 273	<i>Braunechinus similis</i>	277
<i>Boletechinus delawaricus</i>	262	<i>Braunechinus waechteri</i>	277
<i>Boletechinus rowei anglicus</i>	262	<i>brevipetala</i> , <i>Linthia</i>	370
<i>bolschechetensis</i> , <i>Cardiaster</i>	334	<i>brevipetalus</i> , <i>Epiaster</i>	352
<i>bonolai</i> , <i>Desoricidaris</i>	214	<i>brevis</i> , <i>Peraspatangus</i>	389
<i>bononii</i> , <i>Echinolampas (Isolampas)</i>	316	<i>Breynia</i>	391
<i>borjnadorensis</i> , <i>Abelleicidaris</i>	226, 228	<i>Breynia neanika</i>	391
<i>borraeus</i> , <i>Pericosmus</i>	365	<i>bridgeboronensis</i> , <i>Brissus</i>	383

<i>Brisaster</i>	368, 376	<i>bungoensis</i> , <i>Heteraster</i>	353
<i>Brisaster latus</i>	368	<i>burckhardti</i> , <i>Cidarites</i>	217
<i>Brisaster micropetalus</i>	368	<i>butleri</i> , <i>Clypeaster</i>	288
<i>Brisaster moskvini</i>	368	<i>caddoensis</i> , <i>Schizaster</i>	373
<i>Brisaster tasmanicus</i>	368	<i>caementum</i> , <i>Micropsis?</i>	257
Brisasterinae	376	<i>Caenocidaris</i>	231
<i>Brissalius</i>	381	<i>Caenopedina</i>	239
<i>Brissalius vannoordenburgi</i>	381	<i>Caenopedina alanbakeri</i>	239
Brissidae	380	<i>Caenopedina novaezealandiae</i>	239
Brissidea	380	<i>Caenopedina otagoensis</i>	239
<i>Brissopatagus</i>	381	<i>calcalosus</i> , <i>Nucleopygus</i>	325
<i>Brissopatagus cotteau</i>	381	<i>caledonensis</i> , <i>Aptilechinus</i>	206
<i>Brissopatagus depressus</i>	381	<i>californiensis</i> , <i>Calilampas</i>	328
<i>Brissopatagus lucentinus</i>	382	<i>Calilampas</i>	328
<i>Brissopatagus palamoensis</i>	382	<i>Calilampas californiensis</i>	328
<i>Brissopatagus parvus</i>	382	<i>Callaudicidaris</i>	230
<i>Brissopatagus pulchellus</i>	382	<i>Callaudicidaris fowleri</i>	230
<i>Brissopatagus relictus</i>	381	<i>callidus</i> , <i>Hemiaster (Bolbaster)</i>	359
<i>Brissopatagus silleri</i>	382	<i>callista</i> , <i>Prionocidaris</i>	215
<i>Brissopneustes</i>	376	<i>Calocidaris</i>	217
<i>Brissopneustes ruegensis</i>	377	<i>Calocidaris palmeri</i>	217
<i>Brissopneustes rügensis</i>	377	<i>Calycina</i>	241
<i>Brissopneustes vindobonensis</i>	376	<i>Calzadaster</i>	369
<i>Brissopsis</i>	382	<i>Calzadaster friasi</i>	369
<i>Brissopsis alta</i>	366	<i>Camerogalerus</i>	281
<i>Brissopsis australis</i>	382	<i>Camerogalerus cantabricus</i>	281
<i>Brissus</i>	382	<i>candidoi</i> , <i>Rhyncholampas</i>	326
<i>Brissus bridgeboronensis</i>	383	<i>cantabricus</i> , <i>Camerogalerus</i>	281
<i>Brissus daviesi</i>	382	<i>Cardabia</i>	343
<i>Brissus fosteri</i>	383	<i>Cardabia bullarensis</i>	343
<i>Brissus greifatensis</i>	382	<i>Cardiaster</i>	334
<i>Brissus shaimae</i>	382	<i>Cardiaster bolschechetensis</i>	334
<i>Brochopleurus</i>	268	<i>Cardiaster cotteauanus</i>	340
<i>Brochopleurus pliocenicus</i>	268	<i>Cardiaster cotteauanus latohumilis</i>	334
<i>brocki</i> , <i>Heterosalenia</i>	242	<i>Cardiaster protoplanus</i>	334
<i>brodiei</i> , <i>Baudhuinipygus</i>	311	<i>Cardiaster pseudoplanus</i>	334
<i>brodiei</i> , <i>Nucleolites</i>	311	<i>Cardioichnus</i>	396
<i>Bromidechinus</i>	204	<i>Cardioichnus foradadaensis</i>	396
<i>Bromidechinus rimaporus</i>	204	<i>Cardioichnus ovalis</i>	396
<i>bromidensis</i> , <i>Unibothriocidaris</i>	205	<i>Cardioichnus planus</i>	396
<i>brownwoodensis</i> , <i>Archaeocidaris</i>	209	<i>Cardioichnus reniformis</i>	396
<i>brykovi</i> , <i>Scaphechinus</i>	305	<i>Cardiopygus</i>	323
<i>bucharensis</i> , <i>Catopygus</i>	319	<i>Cardiopygus cardus</i>	323
<i>bucharensis</i> , <i>Pseudocatopygus</i>	319	Cardiotaxinae	343
<i>buhaysensis</i> , <i>Heterodiadema</i>	250	<i>Cardiotaxis</i>	343
<i>bulgaricus</i> , <i>Seunaster</i>	342	<i>cardus</i> , <i>Cardiopygus</i>	323
<i>bullarensis</i> , <i>Cardabia</i>	343	<i>Caribbaster</i>	369

<i>Caribbaster dyscritus</i>	369	<i>Cherreauma</i>	238
<i>Caribbaster loveni</i>	369	<i>Cherreauma cherreaui</i>	238
<i>carinatus</i> , <i>Schizaster</i> (<i>Paraster</i>)	375	<i>chesheri</i> , <i>Brachysternaster</i>	368
Carnarechininae	345	<i>Chesniericidaris</i>	217
<i>Carnarechinus</i>	345	<i>Chesniericidaris brangeri</i>	217
<i>Carnarechinus clypeatus</i>	345	<i>Chesniericidaris wrightii</i>	217
<i>carolinensis</i> , <i>Maretia</i>	388	<i>cheveti</i> , <i>Collyrites</i>	332
<i>carolinensis</i> , <i>Psammechinus</i>	274	<i>chiangchunshanensis</i> , <i>Clypeaster</i>	288
<i>carvalhoi</i> , <i>Tholaster</i>	342	<i>chiplonkari</i> , <i>Echinolampas</i>	317
<i>cascadensis</i> , <i>Salenia</i>	244	<i>chiplonkari</i> , <i>Gongrochanus</i>	320
<i>caseyi</i> , <i>Pseudholaster</i>	341	<i>chiplonkari</i> , <i>Meoma</i> (<i>Schizobrissus</i>)	386
Cassidulidae	323	<i>chipolanus</i> , <i>Rhyncholampas</i>	326
Cassiduloidea	307	<i>Chondrocidaris</i>	225
<i>Cassidulus</i>	324	<i>chovaresmicus</i> , <i>Galerites</i>	284
<i>Cassidulus crassus</i>	322	Cidaridae	210
<i>Cassidulus kieri</i>	324	Cidarinae	216
<i>Cassidulus oldhaminus</i>	322	<i>Cidaris</i>	217
<i>Cassidulus santolayae</i>	324	<i>Cidaris aculeata</i>	218
<i>castriauxi</i> , <i>Cenofibula</i>	292	<i>Cidaris aculeata ailaensis</i>	218
<i>catherinae</i> , <i>Phalacrocidaris</i>	213	<i>Cidaris adunca</i>	221
<i>Catopygus</i>	311	<i>Cidaris aggtelekensis</i>	221
<i>Catopygus altus</i>	311	<i>Cidaris aialensis</i>	218
<i>Catopygus bucharensis</i>	319	<i>Cidaris aialensis fusiformis</i>	218
<i>Catopygus dilatatus</i>	311	<i>Cidaris alata brevicaulis</i>	218
<i>Catopygus loevisiformis</i>	311	<i>Cidaris alata canaliculata</i>	218
<i>Catopygus neocylindricus</i>	311	<i>Cidaris alpina</i>	218
<i>Catopygus subcircularis sulcatus</i>	311	<i>Cidaris asperula</i>	226
<i>caupianensis</i> , <i>Amphiope</i>	303	<i>Cidaris baltica</i>	227
<i>cauveriae</i> , <i>Toxaster</i>	354	<i>Cidaris bangtoupensis</i>	221
<i>cearensis</i> , <i>Hemiaster</i>	356	<i>Cidaris batheri</i>	221
<i>celsus</i> , <i>Pericosmus</i>	365	<i>Cidaris bermudezi</i>	217
<i>Cenofibula</i>	292	<i>Cidaris biconica</i>	218
<i>Cenofibula castriauxi</i>	292	<i>Cidaris bradfordensis</i>	228
<i>Centrostephanus</i>	237	<i>Cidaris comptoni</i>	217
<i>Centrostephanus sylviae</i>	237	<i>Cidaris coralliophila</i>	218
<i>Ceratophysa</i>	346	<i>Cidaris costalarensis</i>	218
<i>Ceratophysa ceratopyga valvaecristata</i>	346	<i>Cidaris costata</i>	218
<i>ceratopyga valvaecristata</i> , <i>Ceratophysa</i>	346	<i>Cidaris costata tuberculata</i>	218
<i>cetus</i> , <i>Eupatagus</i>	383	<i>Cidaris costeanensis</i>	219
<i>changchihkengensis</i> , <i>Peronella</i>	295	<i>Cidaris desnoyersi</i>	229
<i>changi</i> , <i>Fossilaster</i>	289	<i>Cidaris desori</i>	228
<i>chargensis</i> , <i>Paraster</i>	375	<i>Cidaris dorsata</i>	277
<i>chartroni</i> , <i>Acrosalenia</i>	233	<i>Cidaris dorsata alata</i>	219
<i>chartroni</i> , <i>Nicolleuma</i>	233	<i>Cidaris dorsata fusiformis</i>	219
<i>chenjafra</i> , <i>Apoxypetalum</i>	381	<i>Cidaris etalloni</i>	227
<i>cherei</i> , <i>Salenia</i>	245	<i>Cidaris fangchui</i>	221
<i>cherreaui</i> , <i>Cherreauma</i>	238	<i>Cidaris flexuosa</i>	276

<i>Cidaris flexuosa brevicostata</i>	219	<i>Cidaris valparolae</i>	221
<i>Cidaris forminensis</i>	219	<i>Cidaris venustespinoza</i>	222
<i>Cidaris forminensis</i>	219	<i>Cidaris verticillata</i>	221
<i>Cidaris foveata</i>	214	<i>Cidaris wachangpoensis</i>	222
<i>Cidaris fowleri</i>	230	<i>Cidaris waechteri</i>	277
<i>Cidaris giauensis</i>	219	<i>Cidaris wrightii</i>	217
<i>Cidaris glabra</i>	219	<i>Cidaris zschokkei</i>	228
<i>Cidaris gladius</i>	221	<i>Cidarites</i>	217
<i>Cidaris gu</i>	221	<i>Cidarites blumenbachii</i>	227
<i>Cidaris haussmanni</i>	277	<i>Cidarites burckhardti</i>	217
<i>Cidaris hirudo</i>	213	<i>Cidarites elegans</i>	228, 229
<i>Cidaris lancedelli</i>	219, 224	<i>Cidarites grandaeus</i>	233
<i>Cidaris leberti</i>	218	<i>Cidarites interpunctata</i>	247
<i>Cidaris leonardi</i>	219	<i>Cidarites maximus</i>	230
<i>Cidaris lineola campanulata</i>	219	<i>Cidaroida</i>	208
<i>Cidaris mafengpoensis</i>	221	<i>Cidaroidea</i>	204
<i>Cidaris major</i>	231	<i>Circopeltinae</i>	258
<i>Cidaris milierensis</i>	219	<i>Circopeltis</i>	258, 259
<i>Cidaris moeschi</i>	227	<i>Circopeltis maastrichtensis</i>	257
<i>Cidaris montanaro</i>	219, 223	<i>Circopeltis? emiratus</i>	259
<i>Cidaris moorei</i>	228	<i>circularis, Gongrochanus</i>	320
<i>Cidaris moralina</i>	230	<i>Claviaster</i>	331
<i>Cidaris morieri</i>	232	<i>Claviasteridae</i>	331
<i>Cidaris ovata</i>	219	<i>Clitopygidae</i>	331
<i>Cidaris palaformis</i>	222	<i>Clitopygus</i>	331
<i>Cidaris parandieri</i>	227	<i>clubis, Balanocidaris</i>	216
<i>Cidaris plana</i>	220	<i>Clypeaster</i>	288
<i>Cidaris pleracantha</i>	227	<i>Clypeaster butleri</i>	288
<i>Cidaris quadrialata</i>	220	<i>Clypeaster chiangchunshanensis</i>	288
<i>Cidaris quadriserrata</i>	220	<i>Clypeaster dondolii</i>	289
<i>Cidaris reticulata</i>	220	<i>Clypeaster gombosae</i>	289
<i>Cidaris rinbianchi</i>	220, 223	<i>Clypeaster hothiensis</i>	289
<i>Cidaris saemanni</i>	228	<i>Clypeaster isolatus</i>	288
<i>Cidaris seelandica</i>	220	<i>Clypeaster kieri</i>	288
<i>Cidaris seelandica misurinensis</i>	220	<i>Clypeaster kurangaensis</i>	289
<i>Cidaris serradentata</i>	222	<i>Clypeaster maanliaoensis</i>	288
<i>Cidaris similis</i>	277	<i>Clypeaster maoadentroensis</i>	288
<i>Cidaris smithii</i>	227	<i>Clypeaster paishatunensis</i>	288
<i>Cidaris spongiosa</i>	220, 221	<i>Clypeaster papilionensis</i>	289
<i>Cidaris sublaevis</i>	228	<i>Clypeaster paraensis</i>	289
<i>Cidaris sulcata</i>	220	<i>Clypeaster pitouensis</i>	288
<i>Cidaris tenuicostata</i>	220	<i>Clypeaster tauricus elatior</i>	289
<i>Cidaris tetraedrica</i>	220	<i>Clypeaster yeni</i>	289
<i>Cidaris tetraedrica appendiculata</i>	220	<i>Clypeaster yingkoensis</i>	289
<i>Cidaris toucasi</i>	230	<i>Clypeasteridae</i>	288
<i>Cidaris trapezoidalis</i>	221	<i>Clypeasterina</i>	288
<i>Cidaris uniformis</i>	213	<i>Clypeasteroida</i>	287, 307

<i>clypeatus</i> , <i>Carnarechinus</i>	345	<i>compressus</i> , <i>Bolbaster</i>	359
<i>clypeatus</i> , <i>Cystechinus</i>	345	<i>comptoni</i> , <i>Cidaris</i>	217
Clypeidae	308	<i>comptoni</i> , <i>Goniocidaris</i>	217
<i>clypeiformis</i> , <i>Pygorhynchus</i>	314	<i>conica</i> , <i>Scutella</i>	297
Clypeolampadidae	327	<i>conicus</i> , <i>Homoeaster</i>	363
<i>Clypeus</i>	309	Conoclypidae	285
<i>Clypeus altus</i>	308	Conoclypina	285
<i>Clypeus angustiporus</i>	308	<i>Conoclypus</i>	285
<i>Clypeus babeaui</i>	308	<i>Conoclypus mittereri</i>	285
<i>Clypeus davoustianus</i>	308	<i>Conoclypus nettoanus</i>	309
<i>Clypeus ederlei</i>	309	Conulidae	281, 286
<i>Clypeus mulleri</i>	308	<i>Conulus</i>	282
<i>Clypeus orbicularis</i>	307	<i>Conulus akdjarensis</i>	282
<i>Clypeus rathieri</i>	308	<i>Conulus angulatus</i>	282
<i>Codechinus</i>	260	<i>Conulus arcus</i>	282
<i>Codechinus prosorovskyi</i>	260	<i>Conulus cookei</i>	282
<i>Codiopsis</i>	264	<i>Conulus djanelidzei</i>	282
<i>Codiopsis lehmannae</i>	264	<i>Conulus kubatliensis</i>	282
<i>Codiopsis rupellensis</i>	264	<i>Conulus lamberti</i>	282
<i>Codiopsis smiseri</i>	264	<i>Conulus sinensis</i>	282
Coelopleuridae	267	<i>cookei</i> , <i>Conulus</i>	282
<i>Coelopleurus</i>	264, 267	<i>cookei</i> , <i>Echinolampas</i>	317
<i>Coelopleurus exquisitus</i>	264	<i>cookei</i> , <i>Prionocidaris</i>	215
<i>Coelopleurus ulugqatensis</i>	264	<i>cooki</i> , <i>Echinolampas</i>	317
Coenholectypinae	280	<i>cooperi</i> , <i>Meoma (Schizobrissus)</i>	386
<i>Coenholectypus</i>	279, 280	<i>Coptodiscus</i>	279, 280
<i>Coenholectypus hideshimensis</i>	279	<i>Coptodiscus magniproctus</i>	279
<i>colchesterensis</i> , <i>Echinodiscus</i>	304	<i>coralliophila</i> , <i>Cidaris</i>	218
<i>collabus</i> , <i>Eupatagus</i>	384	<i>coramandeli</i> , <i>Proisaster</i>	354
<i>collenoti</i> , <i>Nicolleaua</i>	233	<i>coranguinum simpsoni</i> , <i>Micraster</i>	378
<i>collenoti</i> , <i>Pseudodiadema</i>	233	<i>Coraster</i>	350
<i>Colliclypeus</i>	309	<i>Coraster urmaensis</i>	350
<i>Colliclypeus nettoanus</i>	309	<i>Coraster vilanovae</i>	350
<i>Collyrites</i>	332	<i>coravium</i> , <i>Micraster</i>	378
<i>Collyrites analis dimota</i>	332	<i>cordiformis</i> , <i>Orbignyana</i>	333
<i>Collyrites cheveti</i>	332	<i>cordobai</i> , <i>Discoides</i>	281
<i>Collyrites jaccardi</i>	333	<i>coreyi</i> , <i>Vaquerosella</i>	301
Collyritidae	332, 351	<i>coronalis</i> , <i>Eupatagus</i>	383
Collyritoida	351	<i>cortinensis</i> , <i>Miocidaris</i>	210
Collyropsinae	333	Corystidae	343, 351
<i>Collyropsis</i>	332, 333	<i>Corystus</i>	343
<i>Collyropsis ovoides</i>	332	Corystusidae	343
<i>Collyropsis ultimus</i>	332	<i>costalarensis</i> , <i>Cidaris</i>	218
<i>coloniensis</i> , <i>Amplaster</i>	301	<i>costata</i> , <i>Cidaris</i>	218
<i>complanata</i> , <i>Abertella</i>	305	<i>costata tuberculata</i> , <i>Cidaris</i>	218
<i>compressa</i> , <i>Toxaster</i>	354	<i>costatus</i> , <i>Hemiaster</i>	356
<i>compressa</i> , <i>Toxaster</i>	354	<i>costeanensis</i> , <i>Cidaris</i>	219

<i>Cottaldia</i>	264	<i>cyclotus</i> , <i>Pygaulus</i>	313
<i>Cottaldia paquettei</i>	264	<i>cylindriformis</i> , <i>Arnaudaster</i>	315
<i>cotteauanus</i> , <i>Cardiaster</i>	340	<i>Cyphosoma duplicatum</i>	258
<i>cotteauanus</i> , <i>Protocardiaster</i>	340	<i>Cyphosoma legayi</i>	258
<i>cotteauanus latohumilis</i> , <i>Cardiaster</i>	334	<i>Cystechinus clypeatus</i>	345
<i>cotteaui</i> , <i>Aliaster</i>	361	<i>dagestanensis</i> , <i>Orthaster</i>	364
<i>cotteaui</i> , <i>Brissopatagus</i>	381	<i>dalmiapurensis</i> , <i>Phymosoma</i>	257
<i>cotteaui</i> , <i>Hemiaster</i>	361	<i>damensis</i> , <i>Fibularia</i>	293
<i>cotteri</i> , <i>Aliaster</i>	361	<i>danicus</i> , <i>Neoglobator</i>	285
<i>cotteri</i> , <i>Opissaster</i>	361	<i>danubiana</i> , <i>Salenia</i>	245
<i>Cottreaucorys</i>	351	<i>daralagezensis</i> , <i>Echinolampas</i>	317
<i>Cottreaucorys kollmanni</i>	351	<i>davidi</i> , <i>Amphipneustes</i>	367
<i>couvelardi</i> , <i>Couvelardicidaris</i>	228	<i>daviesi</i> , <i>Brissus</i>	382
<i>Couvelardicidaris</i>	227, 228	<i>davoustiana mercieri</i> , <i>Pedina</i>	240
<i>Couvelardicidaris couvelardi</i>	228	<i>davoustianus</i> , <i>Angusticypeus</i>	308
<i>Couvelardicidaris moorei</i>	228	<i>davoustianus</i> , <i>Clypeus</i>	308
<i>Crassiholaster</i>	334	<i>Deakia</i>	386
<i>Crassiholaster bischoffi</i>	334	<i>debensis</i> , <i>Heteraster</i>	353
<i>Crassiholaster sphaericus</i>	334	<i>debensis</i> , <i>Pygorhynchus</i>	314
<i>Crassiholaster subglobosus</i>	334	<i>delawaricus</i> , <i>Boletechinus</i>	262
<i>Crassiholaster subglobosus</i> forma <i>norfolkensis</i>	334	<i>Dendraster</i>	298
<i>Crassiholaster sulciproctus</i>	334, 335	<i>Dendraster sullivanii</i>	298
<i>crassisulcatus</i> , <i>Toxaster</i>	355	<i>Dendrasteridae</i>	291, 298
<i>crassus</i> , <i>Cassidulus</i>	322	<i>densituberculata</i> , <i>Magnosia</i>	266
<i>crassus</i> , <i>Progongrochanus</i>	322	<i>depressa</i> , <i>Fibularia</i>	294
<i>Cravechinidae</i>	208	<i>depressa</i> , <i>Plesiosalenia</i>	247
<i>Cravechinus</i>	208	<i>depressa</i> , <i>Pseudocidaris</i>	251
<i>Cravechinus?</i> <i>frankei</i>	208	<i>depressa</i> , <i>Salenia</i>	247
<i>crenularis</i> , <i>Gagara</i>	273	<i>depressoides</i> , <i>Holectypus</i>	280
<i>crenulata</i> , <i>Farquharsonia</i>	241	<i>depressus</i> , <i>Brissopatagus</i>	381
<i>cretensis</i> , <i>Pronechinus</i>	206	<i>depressus</i> , <i>Diadema</i>	252
<i>crickleyensis</i> , <i>Pronucleolites?</i>	307	<i>depressus</i> , <i>Girardema</i>	252
<i>crispus</i> , <i>Echinocyamus</i> (<i>Lepidocyamus</i>)	293	<i>depressus</i> , <i>Hemicidaris</i>	250
<i>Ctenocidarinae</i>	211	<i>depressus</i> , <i>Micraster</i>	380
<i>Ctenocidaris</i>	212	<i>depressus</i> , <i>Peraspatangus</i>	389
<i>Ctenocidaris aotearoa</i>	212	<i>depressus</i> , <i>Pseudholaster</i>	341
<i>Culozoma</i>	238	<i>depressus</i> , <i>Pseudogibbaster</i>	380
<i>Culozoma baroini</i>	238	<i>dermodyorum</i> , <i>Spatangobrissus</i>	389
<i>curukensis</i> , <i>Hemiaster</i>	356	<i>desnoyersi</i> , <i>Cidaris</i>	229
<i>Cyathocidaris</i>	222	<i>desnoyersi</i> , <i>Philicidaris</i>	229
<i>Cyathocidaris avenionensis nerthensis</i>	222	<i>Desorella</i>	287, 331, 395
<i>Cyclaster</i>	376, 383	<i>Desorella moravica</i>	395
<i>Cyclaster aturicus</i>	376	<i>Desorellidae</i>	287
<i>Cyclaster azemati</i>	383	<i>desori</i> , <i>Cidaris</i>	228
<i>Cyclaster galei</i>	383	<i>desori</i> , <i>Micraster</i>	371
<i>Cyclaster platornatus</i>	383	<i>desori</i> , <i>Nenoticidaris</i>	228
		<i>desori caucasica</i> , <i>Prenaster</i>	371

<i>Desoricidaris</i>	214	<i>Discooides</i>	281
<i>Desoricidaris balli</i>	214	<i>Discooides cordobai</i>	281
<i>Desoricidaris bonolai</i>	214	Discooididae	281, 286
<i>Desoricidaris pouyannei</i>	214	<i>disparradiata</i> , forma aegra	398
<i>Desoricidaris salviensis</i>	214	<i>dissimilariis</i> , <i>Stomechinus</i> ?	261
<i>Desoricidaris sanctaecrucis</i>	214	<i>Ditremaster</i>	361
<i>Desoricidaris schweinfurthi</i>	214	<i>Ditremaster aslaniani</i>	361
<i>Desoricidaris subvenulosa</i>	214	<i>Ditremaster mereirensis</i>	361
<i>Desoricidaris venulosa</i>	214	<i>divinata</i> , <i>Leodia</i>	302
<i>devlinensis</i> , <i>Porterpygus</i>	330	<i>djabarovi</i> , <i>Parahemiaster</i>	362
<i>devonicus</i> , <i>Albertechinus</i>	206	<i>djakonovi</i> , <i>Marginoproctus</i>	294
<i>dhofarensis</i> , <i>Trochodiadema</i> ?	256	<i>djalilovi</i> , <i>Parahemiaster</i>	362
<i>dhrumaensis</i> , <i>Acrosalenia</i>	242	<i>djanelidzei</i> , <i>Conulus</i>	282
<i>dhrumaensis</i> , <i>Bothryopneustes</i>	309	<i>doederleini</i> , <i>Paraster</i>	374
<i>dhrumaensis</i> , <i>Heterosalenia</i>	242	<i>dolosus</i> , <i>Hemiaster (Bolbaster)</i>	359
<i>Diadema</i>	237	<i>Domechinus</i>	319
<i>Diadema depressus</i>	252	<i>Domechinus sinensis</i>	319
<i>Diadema setosumdepressa</i>	237	<i>Domechinus vialovi</i>	319
Diadematacea	234	<i>dondolii</i> , <i>Clypeaster</i>	289
Diadematidae	236	<i>dorsata</i> , <i>Braunechinus</i>	277
Diadematoida	236	<i>dorsata</i> , <i>Cidaris</i>	277
<i>diadematooides</i> , <i>Archaeocidaris</i> ?	209	<i>dorsata alata</i> , <i>Cidaris</i>	219
<i>Dickesicidaris</i>	230	<i>dorsata fusiformis</i> , <i>Cidaris</i>	219
<i>Dickesicidaris maximus</i>	230	<i>Douvillaster</i>	351
<i>Dicoptella bigoti</i>	271	<i>Douvillaster subtrigonalis</i>	351
<i>dictyopleuroides</i> , <i>Zeuglopleurus</i>	272	<i>drogiacus</i> , <i>Metaporinus</i>	333
<i>dietli</i> , <i>Romanocidaris</i>	229	<i>drygalskii</i> , <i>Antrechinus</i>	344
<i>dilatatus</i> , <i>Catopygus</i>	311	<i>drygalskii</i> , <i>Urechinus</i>	344
<i>dineuri</i> , <i>Samlandaster</i>	305	<i>drygalskii perfidus</i> , <i>Urechinus</i>	345
Diplocidaridae	227	<i>duncani</i> , <i>Echinolampas</i>	316
<i>Diplocidaris jacquemeti</i>	227	<i>duncani</i> , <i>Hemipneustes</i>	338
<i>Diplodetus</i>	377	<i>duplicatum</i> , <i>Cyphosoma</i>	258
<i>Diplodetus (Protobrissus)</i>	377	<i>duplicatum</i> , <i>Pomeliosoma</i>	258
<i>Diplodetus (Protobrissus) kurkurensis</i>	377	<i>durhami</i> , <i>Almucidaris</i>	216
<i>Diplopodia</i>	252, 259	<i>durhami</i> , <i>Fernandezaster</i>	385
<i>Diplopodia atlasensis</i>	252	<i>durhami</i> , <i>Fossulaster</i>	289
<i>Diplopodia bipuncata</i>	252	<i>dux</i> , <i>Platysalenia</i>	247
<i>Diplopodia thieulini</i>	252	<i>dux</i> , <i>Salenia</i>	247
Diplopodiidae	259	<i>Dysaster loryi</i>	332
<i>Diploporaster</i>	369	<i>Dysaster ovulum</i>	333
<i>Diploporaster flemingi</i>	369	<i>dyscritus</i> , <i>Caribbaster</i>	369
<i>Diplosalenia</i>	275	<i>dyscritus</i> , <i>Schizaster</i>	369
<i>Diplosalenia faurei</i>	275	Echinacea	241
<i>Diplotagma</i>	260	<i>Echinadesma</i>	302
<i>Diplotagma snellingi</i>	260	<i>Echinanthus</i>	331
<i>Dipneustes</i>	374	<i>Echinanthus reguanti</i>	331
Disasteridae	333	<i>Echinanthus schweinfurthi devriesi</i>	332

Echinarachniidae	291, 298	<i>Echinocyamus insularis macroproctus</i>	292
<i>Echinarachnius</i>	299	<i>Echinocyamus jaisalmerensis</i>	293
<i>Echinarachnius astrodapsoides</i>	299	<i>Echinocyamus nummulicus bernaniensis</i>	292
<i>Echinarachnius plafkeri</i>	299	<i>Echinocyamus pannonicus</i>	293
<i>Echinarachnius ungaensis</i>	299	<i>Echinocyamus raoi</i>	292
Echinidae	274	<i>Echinocyamus wilsoni</i>	292
<i>Echinobrissus</i>	312	<i>Echinocyamus (Lepidocyamus)</i>	293
<i>Echinobrissus faasi</i>	312	<i>Echinocyamus (Lepidocyamus) crispus</i>	293
<i>Echinobrissus gultschensis</i>	312	<i>Echinocyphus</i>	268, 273
<i>Echinobrissus haimeii</i>	311	<i>Echinocyphus intermedius</i>	269
<i>Echinobrissus vincentinus</i>	330	Echinocystitidae	205
<i>Echinobrissus volgensis</i>	312	Echinocystitioida	205
<i>Echinobrissus (Nucleolites) pannonicus</i>	312	<i>Echinodiscus</i>	304
<i>Echinocardium</i>	391	<i>Echinodiscus colchesterensis</i>	304
<i>Echinocardium biaense</i>	392	<i>Echinodiscus hsianglanensis</i>	304
<i>Echinocardium kelloggi</i>	392	<i>Echinodiscus tiliensis</i>	304
<i>Echinocardium leopolitanum</i>	392	<i>Echinodiscus yeliuensis</i>	304
<i>Echinocardium marylandiense</i>	392	<i>Echinogalerus</i>	287
<i>Echinocardium meteorensis</i>	391	<i>Echinogalerus raingeardi</i>	287
<i>Echinocardium olisiponensis</i>	392	<i>Echinogutta</i>	349
<i>Echinocardium peroni</i>	392	<i>Echinogutta amphora amphora</i>	349
<i>Echinocorys</i>	335	<i>Echinogutta amphora indica</i>	349
<i>Echinocorys ancileformis</i>	335	<i>Echinogutta antartica</i>	349
<i>Echinocorys australis</i>	335	<i>Echinogutta fabrefacta</i>	349
<i>Echinocorys balcanicus</i>	335	<i>Echinogutta valvaedentata</i>	349
<i>Echinocorys elongatus</i>	335	Echinoid Traces	396
<i>Echinocorys euxinus</i>	336	Echinoida	274
<i>Echinocorys kharagoulensis</i>	335	Echinoidea	204
<i>Echinocorys kongieli</i>	335	Echinolampadidae	315
<i>Echinocorys schafferi</i>	335	<i>Echinolampas</i>	316
<i>Echinocorys schwetzerovi</i>	335	<i>Echinolampas acuminata</i>	316
<i>Echinocorys scutata forma ovata</i>	335	<i>Echinolampas alexandri forma</i>	
<i>Echinocorys scutata forma pyrenaica</i>	337	<i>arctambulacrum</i>	316
<i>Echinocorys sphaericus</i>	334	<i>Echinolampas antunesi</i>	316
<i>Echinocorys stomias</i>	336	<i>Echinolampas bastai</i>	317
<i>Echinocorys sulcata cognatus</i>	335	<i>Echinolampas chipionkari</i>	317
<i>Echinocorys sulcata orbiculata</i>	335	<i>Echinolampas cookei</i>	317
<i>Echinocorys terminata</i>	336	<i>Echinolampas cooki</i>	317
<i>Echinocorys turanica</i>	335	<i>Echinolampas daralagezensis</i>	317
<i>Echinocorys zejszneri</i>	336	<i>Echinolampas duncani</i>	316
<i>Echinocrepis</i>	346	<i>Echinolampas garoensis</i>	317
<i>Echinocrepis rostrata</i>	346	<i>Echinolampas granti</i>	318
<i>Echinocutia</i> [sic!]	349	<i>Echinolampas gregoryi corrugata</i>	316
<i>Echinocyamus</i>	292	<i>Echinolampas gregoryi gregoryi</i>	316
<i>Echinocyamus gurnahensis</i>	292	<i>Echinolampas guvarensis</i>	317
<i>Echinocyamus hungaricus</i>	292	<i>Echinolampas jhadwensis</i>	318
<i>Echinocyamus insularis</i>	292	<i>Echinolampas jigniensis</i>	318

<i>Echinolampas khariensis</i>	318	<i>elegans, Nudicidaris</i>	228
<i>Echinolampas kieri</i>	317	<i>elegans, Remycidaris</i>	229
<i>Echinolampas lakiensis</i>	318	<i>elegantula, Kewia</i>	300
<i>Echinolampas laubei</i>	316	<i>elevatus, Ovulaster</i>	379
<i>Echinolampas laurillardi acuminata</i>	316	<i>Elliptechinus</i>	206
<i>Echinolampas lipiformis</i>	318	<i>Elliptechinus kiwiaster</i>	206
<i>Echinolampas merhi</i>	318	<i>ellipticus, Amplaster</i>	301
<i>Echinolampas pipurensis</i>	318	<i>elongatus, Echinocorys</i>	335
<i>Echinolampas posterocrassa curtata</i>	317	<i>elongatus, Nucleolites</i>	312
<i>Echinolampas qattamiaensis</i>	318	<i>eltaniana, Aporocidaris</i>	211
<i>Echinolampas rotundus</i>	318	<i>eluvialis, Hemiaster (Leymeriaster)</i>	360
<i>Echinolampas schultzi</i>	316	<i>emarginata, Encope</i>	302
<i>Echinolampas tachanabatensis</i>	318	<i>emiratescus, Noetlingaster</i>	261
<i>Echinolampas tandoni</i>	317	<i>Emiratia</i>	255
<i>Echinolampas tanypetalis</i>	318	<i>Emiratia raskhaimahensis</i>	255
<i>Echinolampas tumulus</i>	316	Emiratidae	255
<i>Echinolampas yadongensis</i>	319	Emiratiidae	255
<i>Echinolampas (Hypsoclypus) liae</i>	316	<i>Emiratus</i>	255
<i>Echinolampas (Isolampas) bononii</i>	316	<i>emiratus, Circopeltis?</i>	259
<i>Echinometra</i>	274	<i>Emiratus raskhaimahensis</i>	255
<i>Echinometra lucunter polypora</i>	274	<i>emmae, Arbacina</i>	268
Echinometridae	274	<i>Enallaster</i>	354
Echinoneidae	286	<i>Enallopneustes meslei</i>	339
Echinoneina	281	<i>Encope</i>	302
Echinoneoida	286	<i>Encope emarginata</i>	302
<i>Echinopedina</i>	240	<i>Encope gatunensis</i>	302
<i>Echinopedina paucituberculata</i>	240	<i>Encope michoacanensis</i>	302
<i>Echinosigra</i>	347	<i>Encope micropora</i>	302
<i>Echinosigra amphora</i>	347	<i>Encope perspective</i>	302
<i>Echinosigra amphora amphora</i>	347	<i>Encope secoensis</i>	302
<i>Echinosigra amphora antarctica</i>	348	<i>Encope wetmorei</i>	302
<i>Echinosigra amphora fabrefacta</i>	348	<i>Encope (Echinadesma)</i>	302
<i>Echinosigra amphora indica</i>	348	<i>ennisi, Lampadocorys</i>	339
<i>Echinosigra amphora valvaedentata</i>	348	<i>eocenicus, Prophylacanthus</i>	216
<i>Echinosigra mortenseni</i>	348	<i>Eodiadema</i>	239
<i>Echinosigra phiale partita</i>	348	<i>Eodiadema lacostei</i>	233
<i>Echinosigra porrecta</i>	348	<i>Eodiadema minutum</i>	233
<i>Echinosigra vityazi</i>	349	Eodiadematida	239
Echinothuriacea	234	Eodiadematidae	239
Echinothuriidae	234	<i>Eogaleropygus</i>	307
Echinothuriinae	234	<i>Eogaleropygus microstoma</i>	307
Echinothurioida	234	Eognathostomata	204
<i>ederlei, Clypeus</i>	309	<i>Eoholaster</i>	336
<i>ederlei, Hemicidaris</i>	250	<i>Eoholaster laffittei</i>	336
<i>eduardobarrosoi, Mellita</i>	302	<i>Eoholaster poslavskae</i>	336
<i>elegans, Astrictypeus</i>	303	<i>Eopericosmus</i>	366
<i>elegans, Cidarites</i>	228, 229	<i>Eopericosmus alta</i>	366

<i>Eopericosmus sveshnikovi</i>	366	<i>Eupatagus singhi</i>	384
<i>Eopericosmus typicus</i>	366	<i>Eupatagus tandoni</i>	384
<i>Eoscutella</i>	297, 298	<i>Eupatagus texanus</i>	384
<i>Eoscutella mirandae</i>	297	<i>Eupatagus wilsoni</i>	385
<i>Eoscutella (Tigilella)</i>	298	<i>Eupatagus (Gymnopatagus)</i>	385
<i>Eoscutella (Tigilella) kamtschatica</i>	298	<i>Eurhodia</i>	328
Eoscutellidae	297	<i>Eurhodia baumi</i>	328
<i>Epiaster</i>	352	<i>Eurhodia relictata</i>	328
<i>Epiaster angulosus</i>	352	<i>Eurhodia rugosa depressa</i>	328
<i>Epiaster baconicus</i>	352	<i>Eurhodia rugosa ideali</i>	328
<i>Epiaster brevipetalus</i>	352	Eurypataginae	390
<i>Epiaster hemiasteriformis</i>	352	<i>Eurypatagus</i>	390
<i>Epiaster hungaricus</i>	352	<i>euxinus, Echinocorys</i>	336
<i>Epiaster hungaricus carinatus</i>	353	<i>exiguus, Fossulaster</i>	290
<i>Epiaster hungaricus intermedius</i>	353	<i>exoletus, Psammechinus</i>	274
<i>Epiaster hungaricus rotundatus</i>	353	<i>exquisitus, Coelopleurus</i>	264
<i>Epiaster miyakoanus</i>	352	<i>extumidus, Schizaster (Paraster)</i>	375
<i>Epiaster nobilis</i>	377	<i>faasi, Echinobrissus</i>	312
<i>Epiaster pseudodistinctus</i>	353	<i>faasi, Nucleolites</i>	312
<i>Epiaster pseudodistinctus oblongus</i>	353	<i>faasi, Pygaulus</i>	313
<i>Epiaster pseudodistinctus rotundiformis</i>	353	<i>Faasia globosa</i>	299
<i>Epiaster romani</i>	352	<i>faassi, Archaeocidaris</i>	210
<i>Epiaster zonarius</i>	352	<i>Faassia</i>	299
<i>Ereipichnus</i>	397	<i>fabrefacta, Echinogutta</i>	349
<i>Ereipichnus geladensis</i>	397	<i>fangchui, Cidaris</i>	221
<i>erinaceus, Heterobrissus</i>	394	<i>faringdonensis, Leptosalenia</i>	246
<i>ernsti, Adelopneustes</i>	282	<i>Farquharsonia</i>	241
<i>ernsti, Galerites (Pironaster)</i>	284	<i>Farquharsonia crenulata</i>	241
<i>estepi, Hemiaster</i>	358	<i>Farquharsonia pineaui</i>	241
<i>estermanni, Lampadocorys?</i>	339	<i>farresi, Pericosmus</i>	365
<i>etalloni, Cidaris</i>	227	<i>Faujasia</i>	320, 321
<i>etalloni, Rolliericidaris</i>	227	<i>Faujasia praeacutus</i>	320
<i>etolonica, Kewia</i>	300	Faujasiidae	319
<i>Eucidaris</i>	225	<i>faurei, Diplosalenia</i>	275
Euechinoidea	232	<i>fecunda, Guettaria</i>	337
<i>eugeniae, Agassizia</i>	367	<i>Fellius</i>	214
<i>Eupatagus</i>	383	<i>Fellius foveata</i>	214
<i>Eupatagus batequensis</i>	384	<i>Fernandezaster</i>	385
<i>Eupatagus cetus</i>	383	<i>Fernandezaster durhami</i>	385
<i>Eupatagus collabus</i>	384	<i>Fibularia</i>	293, 295
<i>Eupatagus coronalis</i>	383	<i>Fibularia damensis</i>	293
<i>Eupatagus flindersi</i>	383	<i>Fibularia depressa</i>	294
<i>Eupatagus lawsonae</i>	384	<i>Fibularia gassusensis</i>	293
<i>Eupatagus ludbrookae</i>	384	<i>Fibularia guavarensis</i>	294
<i>Eupatagus planulatus</i>	384	<i>Fibularia guvarensis</i>	294
<i>Eupatagus rajasthanensis</i>	384	<i>Fibularia japonica</i>	293
<i>Eupatagus rostratus zitteli</i>	384	<i>Fibularia khargahensis</i>	294

<i>Fibularia wizrensis</i>	293	<i>Galerites vulgaris minor</i>	284
Fibulariidae	291, 306	<i>Galerites (Pironaster) ernsti</i>	284
Fibularina	295	<i>Galerites (Pironaster) schmidobtusa</i>	284
<i>fifei, Porosoma</i>	258	Galeritidae	283, 286
<i>flemingi, Diploporaster</i>	369	<i>Galerolampas</i>	327
<i>flexuosa, Cidaris</i>	276	Galeropygidae	307
<i>flexuosa, Zardinitiaris</i>	276	<i>Galeropygus</i>	307
<i>flexuosa brevicostata, Cidaris</i>	219	<i>Galeropygus parviphyllus</i>	307
<i>flindersi, Eupatagus</i>	383	<i>gallemii, Salenocidaris</i>	247
<i>foradadaensis, Cardioichnus</i>	396	<i>gangbensis, Nucleopygus</i>	325
forma aegra <i>disparradiata</i>	398	<i>garoensis, Echinolampas</i>	317
forma aegra <i>inconstans</i>	398	<i>Garumnaster</i>	337, 351
forma aegra <i>insecta</i>	398	<i>Garumnaster lamberti</i>	337
<i>formelli, Schizaster</i>	373	<i>gassusensis, Fibularia</i>	293
<i>forminensis, Cidaris</i>	219	<i>gatunensis, Encope</i>	302
<i>formosus, Hemipatagus</i>	384	<i>gaudensis, Apatopygus</i>	330
<i>forinensis, Cidaris</i>	219	<i>Gauthieria</i>	256
<i>Fossulaster</i>	289	<i>Gauthieria alternus</i>	256
<i>Fossulaster changi</i>	289	<i>Gauthieria? mosae</i>	256
<i>Fossulaster durhami</i>	289	<i>Gauthiosoma</i>	256
<i>Fossulaster exiguus</i>	290	<i>Gauthiosoma princeps</i>	256
Fossulasteridae	289	<i>geladensis, Ereipichnus</i>	397
<i>fosteri, Brissus</i>	383	<i>gemmata, Plegiocidaris?</i>	223
<i>fosteri, Schizaster (Dipneustes)</i>	374	<i>Genocidaris</i>	269
<i>fourmarieri, Tholaster</i>	343	<i>Genocidaris maculata</i>	269
<i>foveata, Cidaris</i>	214	<i>Genus? thieli</i>	278
<i>foveata, Fellius</i>	214	<i>geometricus, Sphaeropleurus</i>	273
<i>fowleri, Callaudicidaris</i>	230	<i>geryvillensis, Kieripygurus</i>	309
<i>fowleri, Cidaris</i>	230	<i>geryvillensis, Pygurus</i>	309
<i>franciscanus, Mesocentrotus</i>	275	<i>gharamulensis, Trochalosoma</i>	262
<i>franciscanus, Strongylocentrotus</i>	275	<i>ghiroboensis, Linthia</i>	370
<i>frankei, Cravechinus?</i>	208	<i>giauensis, Cidaris</i>	219
<i>frescoensis, Studeria?</i>	329	<i>Gibbaster</i>	379
<i>friasi, Calzadaster</i>	369	<i>gibbercula, Parascutella</i>	297
<i>fulva, Araeolampas</i>	390	<i>gibberiensis, Plesiaster</i>	379
<i>fulva, Homolampas</i>	390	<i>gigas, Heterobrissus</i>	394
<i>furlani, Levicidaris</i>	225	<i>gigasei, Rachiosoma</i>	258
<i>fusiformis, Balanocidaris</i>	217	<i>Giraliaster</i>	337, 343
<i>Gagara</i>	273	<i>Giraliaster bellissae</i>	337
<i>Gagara crenularis</i>	273	<i>Giraliaster jubileensis</i>	337
<i>Galeaster</i>	351	<i>Girardema</i>	252
<i>galei, Cyclaster</i>	383	<i>Girardema bipuncata</i>	252
<i>Galeola</i>	336	<i>Girardema depressus</i>	252
<i>Galeola papillosa basiplana</i>	336	<i>gissarensis, Hemiaster</i>	357
<i>Galeraster</i>	284	<i>Gitolampas</i>	329
<i>Galerites</i>	284	<i>Gitolampas mcnamarae</i>	329
<i>Galerites chovaresmicus</i>	284	<i>giulinii, Zardinechinus</i>	224

<i>glabra</i> , <i>Cidaris</i>	219	<i>grandaevus</i> , <i>Cidarites</i>	233
<i>gladius</i> , <i>Cidaris</i>	221	<i>grandaevus</i> , <i>Triadotiaris</i>	233
<i>Globator</i>	283	<i>grandituberculatus</i> , <i>Spatangus</i> (<i>Phymapatagus</i>)	390
<i>Globator minimus</i>	283	<i>granosus kouensis</i> , <i>Toxaster</i>	355
<i>Globator ravni</i>	284	<i>granti</i> , <i>Echinolampas</i>	318
<i>Globoholactypus</i>	280	<i>granti</i> , <i>Proraster</i>	372
<i>Globoholactypus globus</i>	280	<i>granulosus</i> , <i>Offaster</i>	339, 340
<i>globosa</i> , <i>Faasia</i>	299	<i>granulosus</i> , <i>Paradoxechinus</i>	269
<i>globulorostratus</i> , <i>Oolopygus</i>	312	<i>gratilla elatensis</i> , <i>Tripneustes</i>	273
<i>globus</i> , <i>Globoholactypus</i>	280	<i>gravieri suezensis</i> , <i>Nudechinus</i>	272
<i>gloriae</i> , <i>Tetragramma</i>	254	<i>gregoryi corrugata</i> , <i>Echinolampas</i>	316
Glyphocyphidae	267	<i>gregoryi gregoryi</i> , <i>Echinolampas</i>	316
<i>Glyphopneustes</i>	249, 267	<i>greifatensis</i> , <i>Brissus</i>	382
<i>Glyphopneustes hattaensis</i>	249	<i>greifatensis</i> , <i>Temnotrema</i>	270
Glyphopneustidae	267	<i>grossouveri productus</i> , <i>Gualtieria</i> (<i>Claviaster</i>)	385
Glypticidae	267	Growth Anomalies	398
<i>Glypticus</i>	265, 267	<i>gu</i> , <i>Cidaris</i>	221
<i>Glypticus icaunense</i>	265	<i>gualichensis</i> , <i>Abertella</i>	305
Glyptocidaridae	233	<i>Gualtieria</i>	385
<i>Glyptocidaris</i>	233	<i>Gualtieria alicantina</i>	385
Glyptocidaroida	233	<i>Gualtieria</i> (<i>Claviaster</i>)	385
<i>Glyptocyphus</i>	273	<i>Gualtieria</i> (<i>Claviaster</i>) <i>grossouveri</i> <i>productus</i>	385
<i>Glyptopneustes</i>	249	<i>guangxiensis</i> , <i>Melonechinus</i>	207
<i>Gnathichnus</i>	397	<i>guavarensis</i> , <i>Fibularia</i>	294
<i>Gnathichnus pentax</i>	397	<i>Guettaria</i>	337
<i>Gnathichnus stellarum</i>	397	<i>Guettaria fecunda</i>	337
Gnathostomata	278	<i>Guettaria schamchorensis</i>	337
<i>gombosae</i> , <i>Clypeaster</i>	289	<i>guittoni</i> , <i>Milnia</i>	243
<i>Gongrochanus</i>	320, 323	<i>Guittonicidaris</i>	230
<i>Gongrochanus ariyalurensis</i>	320	<i>Guittonicidaris bigoti</i>	230
<i>Gongrochanus chipionkari</i>	320	<i>gultschensis</i> , <i>Echinobrissus</i>	312
<i>Gongrochanus circularis</i>	320	<i>gultschensis</i> , <i>Nucleolites</i>	312
<i>Gongrochanus kieri</i>	320	<i>gurnahensis</i> , <i>Echinocyamus</i>	292
<i>Gongrochanus ottakoviensi</i>	320	<i>gutingkengensis</i> , <i>Taiwanaster</i>	306
<i>Gongrochanus stoliczkai</i>	320	<i>Guttonicidaris</i>	230
<i>Gongrochanus tamilnaduensis</i>	320	<i>guvarensis</i> , <i>Echinolampas</i>	317
<i>Gongrochanus? acutus</i>	320	<i>guvarensis</i> , <i>Fibularia</i>	294
Goniocidarinae	213	<i>Gymnocidaris</i>	222
<i>Goniocidaris</i>	213	<i>Gymnocidaris boutini</i>	222
<i>Goniocidaris comptoni</i>	217	<i>Gymnopatagus</i>	385
<i>Goniocidaris noetlingi</i>	213	<i>Gymnopatagus parvipetalus</i>	385
Goniophorini	249	<i>Gymnopatagus tandoni</i>	384
<i>Goniophorus</i>	249	<i>Gymnotiara</i>	238, 249
<i>Goniopygus</i>	265	<i>Gymnotiara kuhni</i>	238, 249
<i>Goniopygus arabicus</i>	265		
<i>gouldii newbernensis</i> , <i>Rhyncholampas</i>	327		
<i>gracilis</i> , <i>Kina</i>	369		

<i>Hagenowia</i>	337, 343	<i>Hemiaster nucleus</i>	362
<i>Hagenowia blackmorei anterior</i>	337	<i>Hemiaster pecten</i>	357
<i>Haimea</i>	285	<i>Hemiaster planus</i>	357
<i>Haimea bajasurensis</i>	285	<i>Hemiaster pulcher</i>	358
<i>Haimea rojasi</i>	285	<i>Hemiaster rochatensis</i>	357
<i>haimei, Baudhuinipygus</i>	311	<i>Hemiaster rollandi oblongus</i>	358
<i>haimei, Echinobrissus</i>	311	<i>Hemiaster rostratus</i>	357
<i>haleifensis, Schizaster</i>	373	<i>Hemiaster sibiricus</i>	357
<i>halli, Schizaster</i>	373	<i>Hemiaster simakovi</i>	357
<i>hamata, Archaeocidaris</i>	209	<i>Hemiaster stella</i>	362
<i>Hapalosoma</i>	235	<i>Hemiaster subameliae</i>	357
<i>Hapalosoma pulchrum</i>	235	<i>Hemiaster subbibansensis</i>	357
<i>Hardouinia</i>	321, 323	<i>Hemiaster subconicus</i>	358
<i>Hardouinia nuratensis</i>	321	<i>Hemiaster submirabilis</i>	358
<i>harmatuki, Linthia</i>	370	<i>Hemiaster turkestanensis</i>	358
<i>harshadae, Metalia</i>	388	<i>Hemiaster wayensis</i>	358
<i>hattaensis, Glyphopneustes</i>	249	<i>Hemiaster (Bolbaster)</i>	358
<i>hattaensis, Hemiaster (Bolbaster)</i>	359	<i>Hemiaster (Bolbaster) callidus</i>	359
<i>Hattopsis</i>	265	<i>Hemiaster (Bolbaster) dolosus</i>	359
<i>Hattopsis paucituberculatus</i>	260, 265	<i>Hemiaster (Bolbaster) hattaensis</i>	359
<i>Hattopsis sphericus</i>	265	<i>Hemiaster (Bolbaster) subidus</i>	359
<i>hausmanni, Mepygurus</i>	309	<i>Hemiaster (Bolbaster) verecundus</i>	359
<i>hausmanni, Cidaris</i>	277	<i>Hemiaster (Leymeriaster)</i>	360
<i>hausmanni, Wissmannechinus</i>	277	<i>Hemiaster (Leymeriaster) eluvialis</i>	360
<i>hawkinsi, Jesionekechinus</i>	241	<i>Hemiaster (Leymeriaster) polygonalis</i>	360
<i>hawkinsi, Plesiechinus</i>	241	<i>Hemiaster (Malwaster)</i>	360
<i>heckeri, Heteraster</i>	354	<i>Hemiaster (Malwaster) subsimilis</i>	360
<i>Heliocidaris</i>	274	<i>Hemiaster (Mecaster) majungensis</i>	360
<i>Heliocidaris robertsi</i>	274	<i>Hemiaster (Mecaster) mikasaensis</i>	360
<i>Hemiaster</i>	356	<i>Hemiaster (Mecaster) rarus</i>	360
<i>Hemiaster aktaschensis</i>	356	Hemiasteridae	356
<i>Hemiaster amudariensis</i>	356	<i>hemiasteriformis, Epiaster</i>	352
<i>Hemiaster anisopetalus</i>	356	Hemiasterina	356
<i>Hemiaster arachnius</i>	356	Hemicidaridae	249
<i>Hemiaster arcticus</i>	356	<i>Hemicidaris</i>	249
<i>Hemiaster baconicus</i>	358	<i>Hemicidaris bandeli</i>	249
<i>Hemiaster bedakensis</i>	356	<i>Hemicidaris depressus</i>	250
<i>Hemiaster cearensis</i>	356	<i>Hemicidaris ederlei</i>	250
<i>Hemiaster costatus</i>	356	<i>Hemicidaris psiloses</i>	250
<i>Hemiaster cotteaui</i>	361	<i>Hemicidaris rhabdoses</i>	250
<i>Hemiaster curukensis</i>	356	<i>Hemicidaris sidibouzidense</i>	250
<i>Hemiaster estepi</i>	358	<i>Hemicidaris teraniensis</i>	250
<i>Hemiaster gissarensis</i>	357	Hemicidaroida	249
<i>Hemiaster himalayensis</i>	357	<i>hemicidaroides lenoiri, Perisalenia</i>	247
<i>Hemiaster javanicus</i>	357	<i>Hemifaorina</i>	369
<i>Hemiaster kafirniganensis</i>	357	<i>Hemifaorina? rex</i>	369
<i>Hemiaster kunlunensis</i>	357	<i>hemiheterus, Heteraster (Enallaster)</i>	354

<i>Hemipatagus formosus</i>	384	<i>Histocidaris longicollis</i>	211
<i>Hemipedina</i>	240	<i>Histocidaris sicula</i>	211
<i>Hemipedina hudsoni</i>	240	<i>Holaster</i>	338
<i>Hemipedina incipiens</i>	276	<i>Holaster bischoffi</i>	334
<i>Hemipneustes</i>	338, 343	<i>Holaster hungaricus</i>	338
<i>Hemipneustes arabicus</i>	338	<i>Holaster pseudonodulosus</i>	338
<i>Hemipneustes duncani</i>	338	<i>Holaster revestensis</i>	340
<i>Hemipneustes indicus</i>	338	<i>Holaster sanglakensis</i>	338
<i>hemispinifera, Archaeocidaris</i>	210	<i>Holaster subquadratus</i>	338
<i>Hemithylus</i>	256	<i>Holaster tanamensis</i>	338
<i>Hemithylus alternus</i>	256	Holasteridae	334
<i>hemmoorensis, Tylocidaris</i>	226	Holasteroidea	332
<i>heptneri, Pourtalesia</i>	346	Holectypoida	278
<i>Heteraster</i>	352, 353	Holectypoidae	278
<i>Heteraster bungoensis</i>	353	Holectypoina	278
<i>Heteraster debensis</i>	353	<i>Holectypus</i>	280
<i>Heteraster heckeri</i>	354	<i>Holectypus depressoides</i>	280
<i>Heteraster magnus</i>	354	<i>Holectypus khelilense</i>	280
<i>Heteraster renngarteni</i>	354	<i>Holectypus phelani</i>	280
<i>Heteraster zircensis</i>	353	<i>Holectypus pictaviense</i>	280
<i>Heteraster (Enallaster)</i>	354	<i>Holectypus sarthacensis</i>	280
<i>Heteraster (Enallaster) hemiheterus</i>	354	<i>holmesii, Plagionotus</i>	386
<i>Heterobrissus</i>	394	<i>Holosalenia</i>	248
<i>Heterobrissus erinaceus</i>	394	<i>Holosalenia bahiensis</i>	248
<i>Heterobrissus gigas</i>	394	<i>Holosalenia batnensis</i>	248
<i>Heterodiadema</i>	250, 259	<i>Homoeaster</i>	363
<i>Heterodiadema buhaysensis</i>	250	<i>Homoeaster conicus</i>	363
Heterodiademataidae	259	<i>Homolampas fulva</i>	390
<i>Heterosalenia</i>	242	<i>hothiensis, Clypeaster</i>	289
<i>Heterosalenia brocki</i>	242	<i>houphoueti, Amblypygus</i>	
<i>Heterosalenia dhrumaensis</i>	242	(<i>Paramblypygus</i>)	279
<i>Heterosalenia ornata</i>	242	<i>houzeau, Neoglobator</i>	284
<i>hideshimensis, Coenholectypus</i>	279	<i>houzeau, Pyrina</i>	284
<i>Himalayechinus</i>	321	<i>hsianglanensis, Echinodiscus</i>	304
<i>Himalayechinus longus</i>	321	<i>hsui, Pericosmus</i>	365
<i>Himalayechinus minimus</i>	321	<i>hudsoni, Hemipedina</i>	240
<i>Himalayechinus typicus</i>	321	<i>hugueti, Arbacina</i>	268
<i>himalayensis, Hemiaster</i>	357	<i>humei, Leptosalenia?</i>	246
<i>hirudo, Cidaris</i>	213	<i>humei, Salenia</i>	246
<i>hirudo, Hirudocidaris</i>	213	<i>humei, Schizaster</i>	373, 375
<i>Hirudocidaris</i>	213	<i>Hungaresia</i>	311
<i>Hirudocidaris hirudo</i>	213	<i>Hungaresia hungarica</i>	311
<i>Hirudocidaris silesiaca</i>	213	<i>Hungaresia minor</i>	311
<i>Hirudocidaris uniformis</i>	213	<i>hungarica, Archiacia</i>	323
Histocidarinae	210	<i>hungarica, Hungaresia</i>	311
Histocidarini	210	<i>hungarica, Rispolia</i>	342
<i>Histocidaris</i>	211	<i>hungarica, Triadocidaris</i>	210

<i>hungaricus, Echinocyamus</i>	292	<i>Infulaster praecursor</i>	339
<i>hungaricus, Epiaster</i>	352	<i>insecta, forma aegra</i>	398
<i>hungaricus, Holaster</i>	338	<i>insolitus, Neoglobator</i>	285
<i>hungaricus, Pholiocidaris</i>	207	<i>insularis, Echinocyamus</i>	292
<i>hungaricus carinatus, Epiaster</i>	353	<i>insularis macroproctus, Echinocyamus</i>	292
<i>hungaricus intermedius, Epiaster</i>	353	<i>intermedium, Aspidodiadema</i>	237
<i>hungaricus rotundatus, Epiaster</i>	353	<i>intermedius, Echinocyphus</i>	269
<i>husseini, Jordaniaster</i>	377	<i>interpunctata, Cidarites</i>	247
<i>Huttonechinus</i>	343	<i>interpunctata, Prandinia</i>	247
<i>Huttonechinus antarctica</i>	344	<i>Iranianster</i>	350
<i>Huttonechinus spatangiformis</i>	343	<i>Iranianster affinidouvillei</i>	350
<i>Hyboclypeus</i>	308	<i>Iranianster affinimorgani</i>	350
<i>Hyboclypeus? pavyi</i>	308	<i>Iranianster bowersi</i>	350
<i>Hygrosoma</i>	236	<i>Iranianster omanensis</i>	350
Hygrosomatinae	236	Irregularia	287
<i>Hypodiadema</i>	252	<i>isolatus, Clypeaster</i>	288
<i>Hypodiadema nanituberculata</i>	252	<i>isometra, Mellita</i>	302
<i>Hyposalenia</i>	248, 249	<i>Isomicraster</i>	377
<i>Hyposalenia radians</i>	248	<i>Isomicraster babatakensis</i>	377
Hyposaleniinae	248	<i>jaccardi, Collyrites</i>	333
Hyposaleniini	249	<i>jaccardi, Tetraromania</i>	333
<i>Hypsopygaster</i>	324	<i>jacksoni, Palaechinus</i>	208
<i>Hypsopygaster bajarunasi</i>	324	<i>jacquemeti, Diplocidaris</i>	227
<i>Hypsopygaster rostriformis</i>	324	<i>jacquemeti, Smithicidaris</i>	227
<i>Hysteraster</i>	385	<i>jaisalmerensis, Echinocyamus</i>	293
<i>Hysteraster paragrapsimus</i>	385	<i>jaisalmerensis, Megapneustes</i>	388
<i>icaunense, Glypticus</i>	265	<i>janchrisorum, Rhynchopygus?</i>	327
Ichnotaxa	396	<i>japonica, Fibularia</i>	293
<i>iglahensis, Microcyphus</i>	269	<i>japonicus, Pseudowashitaster</i>	363
<i>iljakensis, Parahemiaster</i>	362	<i>japonicus, Washitaster</i>	363
<i>ilyinensis, pseudoastrodaspis</i>	299	<i>jasenecki, Retzneiosoma</i>	235
<i>imitans, Solenocystis</i>	349	<i>javanicus, Hemiaster</i>	357
<i>impar, Laurenticidaris</i>	231	<i>jeffreysi gibbosa, Pourtalesia</i>	346
<i>impar, Rhabdocidaris</i>	231	<i>jeffreysi lata, Pourtalesia</i>	347
<i>incipiens, Batheritarius</i>	276	<i>jensenae, Rictocystis</i>	349
<i>incipiens, Hemipedina</i>	276	<i>Jesionekechinus</i>	241
<i>inconstans, forma aegra</i>	398	<i>Jesionekechinus hawkinsi</i>	241
<i>incorrectus, Schizaster (Paraster)</i>	375	<i>jhadwensis, Echinolampas</i>	318
<i>incus, Spatangobrissus</i>	389	<i>jhadwensis, Schizaster (Paraster)</i>	374
<i>indicus, Hemipneustes</i>	338	<i>jigniensis, Echinolampas</i>	318
<i>indolensis, Protobrissus</i>	380	<i>Jordaniaster</i>	377
<i>indolensis, Pseudogibbaster</i>	380	<i>Jordaniaster husseini</i>	377
<i>inexpectata, Tylocidaris</i>	226	<i>jourdyi, Aliaster</i>	361
<i>inflata, Bothryopneustes</i>	308	<i>jourdyi, Opissaster</i>	361
Infraclypeidae	331	<i>jubileensis, Giraliaaster</i>	337
<i>Infraclypeus</i>	331	<i>jugamis, Toxaster</i>	355
<i>Infulaster</i>	339, 343	<i>jurensis, Kieripygurus</i>	309

<i>jurensis</i> , <i>Pygurus</i>	309	<i>Kieripygurus geryvillensis</i>	309
<i>Kachchhia</i>	256	<i>Kieripygurus jurensis</i>	309
<i>Kachchhia krohi</i>	256	<i>Kieripygurus rostratus</i>	309
<i>kafirniganensis</i> , <i>Hemiaster</i>	357	<i>Kieripygurus tenuis</i>	309
<i>kairukuensis</i> , <i>Palaeostoma</i>	364	Kieritiaridae	275
<i>kalksburgensis</i> , <i>Scutella</i>	297	<i>Kieritiaris</i>	275, 276
<i>kalon</i> , <i>Allomma</i>	255	<i>Kieritiaris thieli</i>	276
<i>Kamptosoma</i>	235	<i>kiiensis</i> , <i>Trochotiara</i>	254
<i>Kamptosoma abyssale</i>	235	<i>Kina</i>	369
Kamptosomatinae	234	<i>Kina gracilis</i>	369
<i>kamtschatica</i> , <i>Eoscutella</i> (<i>Tigilella</i>)	298	<i>kiwiaster</i> , <i>Elliptechinus</i>	206
<i>kamtschatica</i> , <i>Remondella</i>	301	<i>klairisauzei</i> , <i>Pyrgorhytis</i>	333
<i>kannoi</i> , <i>Kewia</i>	300	<i>klydonos</i> , <i>Psephoaster</i>	363
<i>kaspicum</i> , <i>Porosoma</i>	258	<i>kolatai</i> , <i>Bothriocidaris</i>	204
<i>katherinae</i> , <i>Prionocidaris</i>	215	<i>kollmanni</i> , <i>Cottreaucorys</i>	351
<i>kauffmani</i> , <i>Bothryopneustes</i>	308	<i>Kongielechinus</i>	207
<i>kehoei</i> , <i>Kewia</i>	299	<i>Kongielechinus magnituberculatus</i>	207
<i>kelatensis</i> , <i>Proacrolusia</i>	334	<i>kongieli</i> , <i>Echinocorys</i>	335
<i>kelloggi</i> , <i>Echinocardium</i>	392	<i>kongieli</i> , <i>Trochotiara</i>	254
<i>Kermabrissoides</i>	393, 394	<i>korotkovi</i> , <i>Polydiadema</i>	253
<i>Kermabrissoides siculum</i>	394	<i>krohi</i> , <i>Kachchhia</i>	256
<i>kermadecensis</i> , <i>Metalia</i>	388	<i>kubatliensis</i> , <i>Conulus</i>	282
<i>Kewia</i>	299	<i>kuhni</i> , <i>Gymnotiara</i>	238, 249
<i>Kewia elegantula</i>	300	<i>kunlunensis</i> , <i>Hemiaster</i>	357
<i>Kewia etolonica</i>	300	<i>Kupeia</i>	361
<i>Kewia kannoi</i>	300	<i>Kupeia toi</i>	361
<i>Kewia kehoei</i>	299	<i>kurangaensis</i> , <i>Clypeaster</i>	289
<i>Kewia lituyaensis</i>	300	<i>kurkurensis</i> , <i>Diplodetus</i> (<i>Protobrissus</i>)	377
<i>Kewia marquamensis</i>	300	Kutchypygidae	286
<i>Kewia minima</i>	300	<i>Kutchypygus</i>	286, 287
<i>Kewia tachilniensis</i>	299	<i>Kutchypygus bermotensis</i>	287
<i>kharagoulensis</i> , <i>Echinocorys</i>	335	<i>lacostei</i> , <i>Eodiadema</i>	233
<i>khargahensis</i> , <i>Fibularia</i>	294	<i>lacostei</i> , <i>Nicolleuma</i>	233
<i>khariensis</i> , <i>Echinolampas</i>	318	<i>laffittei</i> , <i>Eoholaster</i>	336
<i>khelilense</i> , <i>Holactypus</i>	280	<i>laffittei</i> , <i>Toxaster</i>	336
<i>kieri</i> , <i>Abatus</i>	366	Laganidae	295
<i>kieri</i> , <i>Cassidulus</i>	324	Laganina	291
<i>kieri</i> , <i>Clypeaster</i>	288	Laganoida	291
<i>kieri</i> , <i>Echinolampas</i>	317	<i>lahuseni</i> , <i>Rhabdocidaris</i>	231
<i>kieri</i> , <i>Gongrochanus</i>	320	<i>lakiensis</i> , <i>Echinolampas</i>	318
<i>kieri</i> , <i>Porterpygus</i>	330	<i>lamberti</i> , <i>Conulus</i>	282
<i>kieri</i> , <i>Tridium</i>	296	<i>lamberti</i> , <i>Garumnaster</i>	337
<i>kieri</i> , <i>Unibothriocidaris</i>	205	<i>Lambertona</i>	365
<i>kieri</i> , <i>Vinchuscanchaia</i>	276, 277	<i>Lambertona perdita</i>	366
<i>Kieria</i>	304	<i>Lambertona perplexa</i>	366
<i>Kieria semseyana</i>	304	<i>lammersmaxi</i> , <i>Plagiochasma</i>	313
<i>Kieripygurus</i>	309	<i>Lampadocorys</i>	339

<i>Lampadocorys ennisi</i>	339	<i>Leptosalenia qishnensis</i>	246
<i>Lampadocorys? estermanni</i>	339	<i>Leptosalenia texana</i>	246
<i>Lamprechinus</i>	271	<i>Leptosalenia volana</i>	246
<i>lancedelli, Cidaris</i>	219, 224	<i>Leptosalenia? humei</i>	246
<i>lancedelli, Zardinechinus</i>	224	<i>Leptosalenia? mathuri</i>	246
<i>lanceolata, Notocidaris</i>	212	<i>lesueuri augusta, Peronella</i>	295
<i>Lanieria</i>	280	<i>Leurocidaris</i>	223, 231
<i>largiporus, Remycidaris</i>	229	<i>Leurocidaris montanaro</i>	223
<i>larumbensis, Sphenaster</i>	376	<i>Levicidaris</i>	225, 276
<i>lata, Lovenia</i>	392	<i>Levicidaris furlani</i>	225
<i>latipetalus, Botriopygus</i>	314	<i>Levicidaris pfaifferi</i>	225
<i>latus, Brisaster</i>	368	<i>Levicidaris zardinia</i>	225
<i>laubei, Echinolampas</i>	316	<i>Leymeriaster</i>	359
<i>Laurenticidaris</i>	230, 231	<i>liae, Echinolampas (Hypsoclypus)</i>	316
<i>Laurenticidaris impar</i>	231	<i>Limpasiaster</i>	324
<i>Laurenticidaris major</i>	231	<i>Limpasiaster ariyalurensis</i>	324
<i>laurillardi acuminata, Echinolampas</i>	316	<i>Limpasiaster pentagonalis</i>	324, 325
<i>lawsonae, Eupatagus</i>	384	<i>Limpasiaster quadralis</i>	324, 325
<i>leberti, Anisocidaris</i>	218	<i>lineola campanulata, Cidaris</i>	219
<i>leberti, Cidaris</i>	218	<i>Linthia</i>	370
<i>legayi, Cyphosoma</i>	258	<i>Linthia africana</i>	370
<i>legayi, Pomeliosoma</i>	258	<i>Linthia bajsarensis</i>	370
<i>lehmannae, Codiopsis</i>	264	<i>Linthia brevipetala</i>	370
<i>Leiocidaris balli</i>	214	<i>Linthia ghiroboensis</i>	370
<i>Leiocidaris leoni</i>	216	<i>Linthia harmatuki</i>	370
<i>Leioechinus</i>	260	<i>Linthia madagascariensis</i>	371
<i>Leioechinus amplus</i>	260	<i>Linthia maliensis</i>	370
<i>Leioechinus namus</i>	260	<i>Linthia monteroeae</i>	370
<i>Leiopedina</i>	240	<i>Linthia pulchra</i>	370
<i>Leiopedina molinai</i>	240	<i>Linthia romani</i>	371
<i>Leiostomaster</i>	364	<i>Linthia? summesbergeri</i>	370
<i>Leiostomaster angolanus</i>	364	<i>lipiformis, Echinolampas</i>	318
<i>Lenticidaris</i>	232	<i>Lissocidaris</i>	223
<i>Leodia</i>	302	<i>Lissocidaris xanthe</i>	223
<i>Leodia divinata</i>	302	<i>lissos, Psephoaster</i>	363
<i>leonardi, Cidaris</i>	219	<i>lituyaensis, Kewia</i>	300
<i>leoni, Leiocidaris</i>	216	<i>loewisiformis, Catopygus</i>	311
<i>leoni, Prophyllacanthus</i>	216	<i>longicollis, Histocidaris</i>	211
<i>leopolitanum, Echinocardium</i>	392	<i>longus, Himalayechinus</i>	321
Lepidocentridae	206	<i>lorioli, Miocidaris</i>	230
<i>Lepidocyamus</i>	293	<i>lorioli, Panniericidaris</i>	230
<i>Leptosalenia</i>	246	<i>loryi, Dysaster</i>	332
<i>Leptosalenia arabica</i>	246	<i>loryi, Smithiaster</i>	332
<i>Leptosalenia faringdonensis</i>	246	<i>Lovenechinus</i>	207
<i>Leptosalenia mexicana</i>	246	<i>Lovenechinus zunyii</i>	207
<i>Leptosalenia plana</i>	246	<i>loveni, Caribbaster</i>	369
<i>Leptosalenia pretensis</i>	246	<i>loveni, Prenaster</i>	369

<i>Lovenia</i>	392	<i>malogostiana, Pseudosalenia</i>	243
<i>Lovenia bagheerae</i>	392	<i>Malwaster</i>	360
<i>Lovenia lata</i>	392	<i>mangyschlakensis, Pseudholaster</i>	341
Loveniidae	380, 390	<i>mangyschlakensis, Pseudoholaster</i> [sic!]	341
<i>lovisatoi, Aliaster</i>	361	<i>mannumensis, Apatopygus</i>	330
<i>lovisatoi, Opissaster</i>	361	<i>maoadentroensis, Clypeaster</i>	288
<i>lowraliensis, Mortonia</i>	294	<i>maquoketensis, Bothriocidaris</i>	204
<i>lucentinus, Brissopatagus</i>	382	<i>marchettii, Scutella</i>	297
<i>lucundus, Parahemiaster</i>	362	<i>marcoui, Acrosalenia</i>	243
<i>lucunter polypora, Echinometra</i>	274	<i>marcoui, Prosalenia</i>	243
<i>ludbrookae, Eupatagus</i>	384	<i>Maretia</i>	387
<i>Lunulaster</i>	301	<i>Maretia carolinensis</i>	388
<i>Lunulaster alatus</i>	301	<i>Maretia planulata abbassi</i>	387
<i>Lytechinus</i>	272	<i>Maretia ranjipurensis</i>	387
<i>Lytechinus baldwini</i>	272	<i>Marginoproctus</i>	294
<i>maanliaoensis, Clypeaster</i>	288	<i>Marginoproctus djakonovi</i>	294
<i>maastrichtensis, Circopeltis</i>	257	<i>mariana, Megaporocidaris</i>	225
<i>maastrichtensis, Phymosoma</i>	257	<i>marisrubis, Asthenosoma</i>	234
<i>Macropneustes</i>	386	<i>marquamensis, Kewia</i>	300
<i>Macropneustes spatangiformis</i>	343	<i>marratensis, Acrosalenia</i>	242
<i>Macropneustes (Deakia) xinjiangensis</i>	386	<i>marylandiense, Echinocardium</i>	392
<i>maculata pliorecens, Genocidaris</i>	269	<i>Masrouraster</i>	263
<i>madagascariensis, Linthia</i>	371	<i>Masrouraster ouhouissi</i>	263
<i>mafengpoensis, Cidaris</i>	221	<i>mathesoni, Spatangus</i>	389
<i>magna, Archiacia</i>	323	<i>mathuri, Leptosalenia?</i>	246
<i>magniprocta, Pseudosalenia</i>	243	<i>mathuri, Salenia</i>	246
<i>magniproctus, Coptodiscus</i>	279	<i>maximus, Cidarites</i>	230
<i>magnituberculatus, Kongielechinus</i>	207	<i>maximus, Dickesicidaris</i>	230
<i>Magnosia</i>	266	<i>mcnamarae, Gitolampas</i>	329
<i>Magnosia densituberculata</i>	266	<i>Mecaster</i>	360
<i>Magnosia suessi</i>	266	<i>Mecaster arnonensis</i>	360
<i>Magnosiopsis</i>	266	<i>Medjesia</i>	339
<i>Magnosiopsis ornata</i>	266	<i>Medjesia meslei</i>	339
<i>Magnosiopsis suessi</i>	266	<i>Megapneustes</i>	388
<i>magnus, Heteraster</i>	354	<i>Megapneustes jaisalmerensis</i>	388
<i>magnus, Nucleopygus</i>	325	<i>Megaporocidaris</i>	225, 276
<i>magnus, Proraster</i>	372	<i>Megaporocidaris mariana</i>	225
<i>magnus, Pyrgorhytis</i>	333	<i>Mellita</i>	302
<i>mai, Peronella</i>	295	<i>Mellita eduardobarrosoi</i>	302
<i>mai, Taiwanaster</i>	306	<i>Mellita isometra</i>	302
<i>major, Cidaris</i>	231	Mellitidae	291, 301
<i>major, Laurenticidaris</i>	231	<i>Melonechinus</i>	207
<i>major, Prowillungaster</i>	290	<i>Melonechinus guangxiensis</i>	207
<i>majungensis, Hemiaster (Mecaster)</i>	360	<i>mendelssohni, Pygurus</i>	310
<i>malavoyi, Pseudopygaulus</i>	329	<i>Mennerella</i>	296
<i>maleckii, Micraster</i>	378	<i>Mennerella ovata</i>	296
<i>maliensis, Linthia</i>	370	<i>Menopygoida</i>	307

<i>Meoma</i>	386	<i>Micraster (Gibbaster) sonorensis</i>	379
<i>Meoma (Schizobrissus)</i>	386	Micrasteridae	376
<i>Meoma (Schizobrissus) chiplonkari</i>	386	Micrasterina	376
<i>Meoma (Schizobrissus) cooperi</i>	386	<i>microapicalis, Parvicidaris</i>	277
<i>Meoma (Schizobrissus) misrai</i>	386	<i>Microcyphus</i>	269
<i>Meoma (Schizobrissus) singhi</i>	386	<i>Microcyphus iglahensis</i>	269
<i>Mepygurus hausmanni</i>	309	<i>micropetalus, Brisaster</i>	368
<i>Mercieripygurus</i>	309	<i>micropora, Encope</i>	302
<i>mereirensis, Ditremaster</i>	361	<i>microprocta, Salenia</i>	244
<i>merhi, Echinolampas</i>	318	<i>Micropsis</i>	257
<i>Merocidaris</i>	231	<i>Micropsis? caementum</i>	257
<i>Merriamaster</i>	298	Micropygidae	239
<i>Merriamaster weaveri</i>	298	Micropygoida	239
<i>meslei, Enallopneustes</i>	339	<i>microstoma, Eogaleropygus</i>	307
<i>meslei, Medjesia</i>	339	Microstomata	307
<i>Mesocentrotus</i>	275	<i>mikasaensis, Hemiaster (Mecaster)</i>	360
<i>Mesocentrotus franciscanus</i>	275	<i>Mikrocidaris</i>	276, 277
<i>Mesocentrotus nudus</i>	275	<i>milierensis, Cidaris</i>	219
<i>Mesodiadema</i>	232	<i>Milnia</i>	243
Mesostomata	287	<i>Milnia guittoni</i>	243
<i>Metalia</i>	388	<i>Mimiosalenia</i>	266
<i>Metalia angustus</i>	388	<i>Mimiosalenia quinquetuberculata</i>	266
<i>Metalia harshadae</i>	388	<i>minima, Kewia</i>	300
<i>Metalia kermadecensis</i>	388	<i>minimus, Globator</i>	283
<i>Metalia sternalis jousseaumei</i>	388	<i>minimus, Himalayechinus</i>	321
<i>Metalia sternalis lata</i>	388	<i>minor, Hungaresia</i>	311
<i>Metaporinus</i>	333	<i>minor, Prowillungaster</i>	290
<i>Metaporinus drogiacus</i>	333	<i>minutum, Eodiadema</i>	233
<i>meteorensis, Echinocardium</i>	391	<i>minutum, Nicolleauma</i>	233
<i>mexicana, Leptosalenia</i>	246	<i>miocenica, Amphiope</i>	303
<i>mexicana, Salenia</i>	246	Miocidaridae	210
<i>miaoliensis, Astrictlypeus</i>	303	<i>Miocidaris</i>	210
<i>miaoliensis, Peronella</i>	295	<i>Miocidaris adrianae</i>	210, 276
<i>michoacanensis, Encope</i>	302	<i>Miocidaris ampezzana</i>	210
<i>Micraster</i>	378	<i>Miocidaris cortinensis</i>	210
<i>Micraster amnicus</i>	378	<i>Miocidaris lorioli</i>	230
<i>Micraster coranguinum simpsoni</i>	378	<i>mirabilis, Pentechinus</i>	270
<i>Micraster coravium</i>	378	<i>mirandae, Eoscutella</i>	297
<i>Micraster depressus</i>	380	<i>mironovi, Amphipneustes</i>	368
<i>Micraster desori</i>	371	<i>misrai, Meoma (Schizobrissus)</i>	386
<i>Micraster maleckii</i>	378	<i>misrai, Tripneustes</i>	273
<i>Micraster praerogalae</i>	378	<i>mississippiensis rosehillensis,</i>	
<i>Micraster solignaci</i>	378	<i>Protoscutella</i>	297
<i>Micraster tercensis</i>	380	<i>mittereri, Conoclypus</i>	285
<i>Micraster turkestanensis</i>	378, 380	<i>miyakoanus, Epiaster</i>	352
<i>Micraster westlakei</i>	378	<i>Mizunamia</i>	395
<i>Micraster (Gibbaster)</i>	379	<i>Mizunamia tokiensis</i>	395

<i>moabitorum</i> , <i>Trochotiara</i>	254	<i>nehalae</i> , <i>Trachypatagus</i>	390
<i>moeschi</i> , <i>Cidaris</i>	227	<i>nelsoni</i> , <i>Pericosmus</i>	376
<i>moeschi</i> , <i>Nenoticdaris</i>	227	<i>nelsoni</i> , <i>Waurnia</i>	376
<i>molinaei</i> , <i>Leiopedina</i>	240	<i>Nenoticdaris</i>	227
<i>monastiriensis</i> , <i>Bichordites</i>	396	<i>Nenoticdaris blumenbachii</i>	227
<i>mongraensis</i> , <i>Phyrosoma</i>	257	<i>Nenoticdaris desori</i>	228
<i>monotuberculatus</i> , <i>Noetlingaster</i>	260, 265	<i>Nenoticdaris moeschi</i>	227
Monphorasteridae	301	<i>Nenoticdaris parandieri</i>	227
<i>mountainvillensis</i> , <i>Adelopneustes</i>	285	<i>Nenoticdaris pichaueriensis</i>	228
<i>mountainvillensis</i> , <i>Neoglobator</i>	284	<i>Nenoticdaris smithii</i>	227
<i>mountainvillensis</i> , <i>Pyrina</i>	284	<i>Nenoticdaris zschokkei</i>	228
<i>montanaro</i> , <i>Cidaris</i>	219, 223	Nenoticidaridae	227
<i>montanaro</i> , <i>Leurocidaris</i>	223	<i>Nenoticdaris</i>	227
<i>montanum</i> , <i>Aspidodiadema</i>	237	<i>neoataxensis</i> , <i>Botriopygus</i>	314
<i>monteroae</i> , <i>Linthia</i>	370	<i>Neobothriocidaris</i>	205
<i>moorei</i> , <i>Cidaris</i>	228	<i>Neobothriocidaris templetoni</i>	205
<i>moorei</i> , <i>Couvelardicidaris</i>	228	<i>neocylindricus</i> , <i>Catopygus</i>	311
<i>moraldina</i> , <i>Cidaris</i>	230	<i>Neoglobator</i>	284
<i>moraldina</i> , <i>Panniericidaris</i>	230	<i>Neoglobator akkajensis</i>	284
<i>moravica</i> , <i>Desorella</i>	395	<i>Neoglobator danicus</i>	285
<i>morgani</i> , <i>Philipaster</i>	290	<i>Neoglobator houzeaui</i>	284
<i>morgani</i> , <i>Scutellina</i>	290	<i>Neoglobator insolitus</i>	285
<i>moriensi</i> , <i>Amblypygus</i>	279	<i>Neoglobator mountainvillensis</i>	284
<i>moriensis</i> , <i>Amblypygus</i>	279	<i>Neoglobator ovalis</i>	284
<i>morierei</i> , <i>Roseicidaris</i>	232	<i>Neoglobator pantelevi</i>	284, 285
<i>mortenseni</i> , <i>Antrechinus</i>	344, 345	<i>Neoglobator ravni</i>	284
<i>mortenseni</i> , <i>Echinosigra</i>	348	<i>Neoglobator subcircularis</i>	284
<i>mortenseni</i> , <i>Urechinus</i>	344, 345	<i>Neoglobator subovalis</i>	284
<i>Mortonia</i>	294	Neoglobatoridae	284
<i>Mortonia lowraliensis</i>	294	Neognathostomata	287
<i>mosae</i> , <i>Gauthieria?</i>	256	<i>neolaevis</i> , <i>Pyrina</i>	283
<i>moskvini</i> , <i>Brisaster</i>	368	Neolaganidae	291
<i>moskvini</i> , <i>Parahemiaster</i>	362	Neolampadidae	394
<i>muelleri</i> , <i>Scutella</i>	297	Neolampadoida	394
<i>mulleri</i> , <i>Angusticlypeus</i>	308	<i>Neoproraster</i>	371, 376
<i>mulleri</i> , <i>Clypeus</i>	308	<i>Neoproraster bajarunasi</i>	371
<i>Murraypneustes</i>	392	<i>Neoproraster usakensis</i>	371
<i>Murraypneustes biannulatus</i>	393	<i>Neoscutella</i>	299
<i>myorensis</i> , <i>Schizaster (Ova)</i>	374	<i>nettoanus</i> , <i>Colliclypeus</i>	309
<i>mysticus</i> , <i>Pseudowashitaster</i>	363	<i>nettoanus</i> , <i>Conoclypus</i>	309
<i>nadorensis</i> , <i>Plagiochasma</i>	313	<i>nicolleaui</i> , <i>Acrocidaris</i>	252
<i>namus</i> , <i>Leioechinus</i>	260	<i>Nicolleauma</i>	233
<i>nanituberculata</i> , <i>Hypodiadema</i>	252	<i>Nicolleauma chartroni</i>	233
<i>Nannoglyphus</i>	269	<i>Nicolleauma collenoti</i>	233
<i>Nannoglyphus wehrlii</i>	269	<i>Nicolleauma lacostei</i>	233
<i>neanika</i> , <i>Breynia</i>	391	<i>Nicolleauma minutum</i>	233
<i>neglecta</i> , <i>Prionocidaris</i>	216	Nicolleaumatidae	233

<i>nitidus nascaensis</i> , <i>Trigonocidaris</i>		<i>Nudicidaris elegans</i>	228
(<i>Lamprechinus</i>)	271	<i>Nudicidaris sublaevis</i>	228
<i>nobilis</i> , <i>Epiaster</i>	377	<i>nudispina</i> , <i>Salenia</i>	244
<i>Noetlingaster</i>	260	<i>nudum</i> , <i>Sperosoma</i>	235
<i>Noetlingaster emiratescus</i>	261	<i>nudus</i> , <i>Mesocentrotus</i>	275
<i>Noetlingaster monotuberculatus</i>	260, 265	<i>nudus</i> , <i>Strongylocentrotus</i>	275
<i>Noetlingaster sphericus</i>	265	<i>nummulicus bernaniensis</i> ,	
<i>noetlingi</i> , <i>Goniocidaris</i>	213	<i>Echinocyamus</i>	292
<i>Nordenskoeldaster</i>	351	<i>nuratensis</i> , <i>Hardouinia</i>	321
<i>Nordenskoeldaster? australis</i>	351	<i>nusplingensis</i> , <i>Polycidaris</i>	211
<i>nordenskjoldi</i> , <i>Antrechinus</i>	344	<i>occultoforma</i> , <i>Orbispala</i>	290
<i>nordenskjoldi</i> , <i>Plexechinus</i>	344	<i>Ochetes</i>	326
<i>norris</i> , <i>Scutella</i>	301	<i>Ochetus</i>	323, 326
<i>Notocidaris</i>	212	<i>Ochetus pauli</i>	326
<i>Notocidaris bakeri</i>	212	<i>Oedematocidaris</i>	226, 227
<i>Notocidaris lanceolata</i>	212	<i>Oedematocidaris asperula</i>	226
<i>Notopygus</i>	313	<i>Oedematocidaris baltica</i>	227
<i>novaezealandiae</i> , <i>Caenopedina</i>	239	<i>Oedematocidaris pleracantha</i>	227
<i>Novasalenia</i>	246	<i>oedumi</i> , <i>Proraster</i>	371
<i>Novasalenia plananyensis</i>	246	<i>Offaster</i>	339
<i>Novasalenia predbojensis</i>	246	<i>Offaster granulatus</i>	339, 340
<i>nuciformis</i> , <i>Offaster</i>	340	<i>Offaster nuciformis</i>	340
<i>Nucleolites</i>	312	<i>Offaster rugius</i>	339
<i>Nucleolites brodiei</i>	311	<i>okhliensis</i> , <i>Orthaster</i>	364
<i>Nucleolites elongatus</i>	312	<i>oldhaminus</i> , <i>Cassidulus</i>	322
<i>Nucleolites faasi</i>	312	<i>oldhaminus</i> , <i>Progongrochanus</i>	322
<i>Nucleolites gultschensis</i>	312	Oligopygidae	285
<i>Nucleolites richardsoni</i>	313	<i>olisiponensis</i> , <i>Echinocardium</i>	392
<i>Nucleolites strictiporus</i>	312	<i>omanensis</i> , <i>Iranianster</i>	350
<i>Nucleolites volgensis</i>	312	<i>oolithicum</i> , <i>Pelanodiadema</i>	236
<i>Nucleolites (Pygorhynchus) plantus</i>	322	<i>Oolopygus</i>	312
<i>Nucleolites (Pygorhynchus) testudo</i>	322	<i>Oolopygus globulorostratus</i>	312
Nucleolitidae	310	<i>Opissaster</i>	361
<i>Nucleopygus</i>	325	<i>Opissaster almerai</i>	361
<i>Nucleopygus bashibulakensis</i>	325	<i>Opissaster cotteri</i>	361
<i>Nucleopygus calcalosus</i>	325	<i>Opissaster jourdyi</i>	361
<i>Nucleopygus gangbensis</i>	325	<i>Opissaster lovisatoi</i>	361
<i>Nucleopygus magnus</i>	325	<i>Opissaster tainui</i>	361
<i>Nucleopygus parvus</i>	326	<i>Opissater subsimilis</i>	360
<i>Nucleopygus peltitipus</i>	326	<i>orbicularis</i> , <i>Clypeus</i>	307
<i>Nucleopygus platypetalus</i>	325	<i>orbicularis</i> , <i>Pronucleolites</i>	307
<i>Nucleopygus salgadoi</i>	325	<i>Orbignyana</i>	333
<i>nucleus</i> , <i>Hemiaster</i>	362	<i>Orbignyana cordiformis</i>	333
<i>nucleus</i> , <i>Parahemiaster</i>	362	<i>Orbispala</i>	290
<i>Nudechinus</i>	272	<i>Orbispala occultoforma</i>	290
<i>Nudechinus gravieri suzezensis</i>	272	<i>Oriolampas</i>	331
<i>Nudicidaris</i>	227, 228	<i>ornata</i> , <i>Heterosalenia</i>	242

<i>ornata</i> , <i>Magnosiopsis</i>	266	<i>Palaeostoma</i>	364
<i>Orthaster</i>	364	<i>Palaeostoma kairukuensis</i>	364
<i>Orthaster alievi</i>	364	Palaeostomatidae	363
<i>Orthaster dagestanensis</i>	364	<i>Palaeotropus</i>	394
<i>Orthaster okhliensis</i>	364	<i>Palaeotropus uniporum</i>	394
Orthopsida	278	<i>palaformis</i> , <i>Cidaris</i>	222
Orthopsidae	278	<i>palamoensis</i> , <i>Brissopatagus</i>	382
<i>Orthopsis</i>	278	Paleopeustidae	366
<i>Orthopsis willei</i>	278	Paleopneustidae	366
Orthosternata	351	Paleopneustoidea	366
<i>otagoensis</i> , <i>Caenopedina</i>	239	<i>palmeri</i> , <i>Calocidaris</i>	217
<i>ottakoviensi</i> , <i>Gongrochanus</i>	320	<i>Palmerius</i>	223
<i>ouhouissi</i> , <i>Masrouraster</i>	263	<i>Palmerius roberti</i>	223
<i>Ova</i>	374	<i>Panniericidaris</i>	229, 230
<i>ova</i> , <i>Peronella</i>	295	<i>Panniericidaris lorioli</i>	230
<i>ovalis</i> , <i>Botriopygus</i>	314	<i>Panniericidaris moralina</i>	230
<i>ovalis</i> , <i>Cardioichnus</i>	396	<i>pannonicus</i> , <i>Echinobrissus</i>	
<i>ovalis</i> , <i>Neoglobator</i>	284	(<i>Nucleolites</i>)	312
<i>ovalis</i> , <i>Pygorhynchus</i>	284	<i>pannonicus</i> , <i>Echinocyamus</i>	293
<i>ovata</i> , <i>Cidaris</i>	219	<i>panteleevi</i> , <i>Neoglobator</i>	284, 285
<i>ovata</i> , <i>Mennerella</i>	296	<i>papilionensis</i> , <i>Clypeaster</i>	289
<i>ovatus</i> , <i>Schizaster (Paraster)</i>	374	<i>papillosa basiplana</i> , <i>Galeola</i>	336
<i>ovoides</i> , <i>Collyropsis</i>	332	<i>pappi</i> , <i>Botriopygus</i>	314
<i>Ovulaster</i>	379	<i>pappi elongatus</i> , <i>Botriopygus</i>	315
<i>Ovulaster elevatus</i>	379	<i>pappi kutassyi</i> , <i>Botriopygus</i>	315
<i>Ovulaster protodecimae</i>	379	<i>paquettei</i> , <i>Cottaldia</i>	264
<i>Ovulaster reticulatus</i>	379	<i>Paracidaris</i>	227
<i>Ovulechinus</i>	286	<i>Paradoxechinus</i>	269
<i>Ovulechinus yadongensis</i>	286	<i>Paradoxechinus granulosus</i>	269
<i>Ovulechinus (Pseudovulechinus)</i>	286	<i>Paradoxechinus profundus</i>	270
<i>Ovulechinus (Pseudovulechinus)</i>		<i>Paradoxechinus stellatus</i>	270
<i>rotundatus</i>	286	<i>paraensis</i> , <i>Clypeaster</i>	289
<i>ovulum</i> , <i>Dysaster</i>	333	<i>paragrapsimus</i> , <i>Hysteraster</i>	385
<i>ovulum</i> , <i>Tetraromania</i>	333	<i>Parahemiaster</i>	362
<i>Pachycentrotus</i>	275	<i>Parahemiaster akkapschigensis</i>	362
<i>Pachycentrotus bajulus</i>	275	<i>Parahemiaster bobkovae</i>	362
<i>Pachyclypus</i>	287, 331	<i>Parahemiaster djabarovi</i>	362
<i>paishatunensis</i> , <i>Clypeaster</i>	288	<i>Parahemiaster djalilovi</i>	362
Palaechinidae	207	<i>Parahemiaster iljakensis</i>	362
Palaechinoida	207	<i>Parahemiaster lucundus</i>	362
<i>Palaechinus</i>	208	<i>Parahemiaster moskvini</i>	362
<i>Palaechinus jacksoni</i>	208	<i>Parahemiaster nucleus</i>	362
<i>Palaechinus praematurus</i>	208	<i>Parahemiaster schmidtae</i>	362
<i>Palaechinus yangi</i>	208	<i>Parahemiaster soluni</i>	362
<i>palaecenicus</i> , <i>Stegaster</i>	342	<i>Parahemiaster stella</i>	362
<i>Palaeodiadema</i>	237	<i>Parahemiaster subrotundus</i>	362
<i>Palaeodiadema reingardae</i>	237	<i>Paralampas</i>	326

<i>Paralampas rancureli</i>	326	<i>Pedinopsis sphaerica</i>	252
<i>Paramaretia</i>	390	<i>Pedinopsis (Sinaiopsis) sinaica</i>	253
<i>Paramblypygus</i>	279	<i>Pelanodiadema</i>	236
<i>parandieri, Cidaris</i>	227	<i>Pelanodiadema oolithicum</i>	236
<i>parandieri, Nenoticdaris</i>	227	<i>Pelanodiademinae</i>	236
<i>Paraphormosoma</i>	236	<i>pellaensis, Praeopholidocidaris</i>	205
<i>Paraphormosomatinae</i>	236	<i>peltitipos, Nucleopygus</i>	326
<i>Parapygus</i>	319	<i>pentagonalis, Limpasiaster</i>	324, 324
<i>Parapygus annae</i>	319	<i>pentagonalis, Salenia</i>	245
<i>Parasaleniiidae</i>	275	<i>pentagonus, Schizechinus</i>	272
<i>Parascutella gibbercula</i>	297	<i>pentax, Gnathichnus</i>	397
<i>Paraster</i>	374	<i>Pentechinus</i>	270
<i>Paraster charginensis</i>	375	<i>Pentechinus mirabilis</i>	270
<i>Paraster doederleini</i>	374	<i>Peraspatangus</i>	388
<i>pareyni, Arbacina</i>	268	<i>Peraspatangus brevis</i>	389
<i>parrasae, Pygopistes</i>	314	<i>Peraspatangus depressus</i>	389
<i>Parvicidaris</i>	276, 277	<i>perdita, Lambertona</i>	366
<i>Parvicidaris microapicalis</i>	277	<i>Pericosmidae</i>	364
<i>parvipetalus, Gymnopatagus</i>	385	<i>Pericosmus</i>	365
<i>parviphyllus, Galeropygus</i>	307	<i>Pericosmus abatooides</i>	375
<i>parvituberculatus, Polycyphus</i>	261	<i>Pericosmus annosus</i>	365
<i>parvus, Brissopatagus</i>	382	<i>Pericosmus borraeus</i>	365
<i>parvus, Nucleopygus</i>	326	<i>Pericosmus celsus</i>	365
<i>parvus, Urechinus</i>	345	<i>Pericosmus farresi</i>	365
<i>paucituberculata, Amoraster</i>	380	<i>Pericosmus hsui</i>	365
<i>paucituberculata, Echinopedina</i>	240	<i>Pericosmus nelsoni</i>	376
<i>paucituberculatus, Hattopsis</i>	260, 265	<i>Pericosmus porphyrocardius</i>	365
<i>pauli, Ochetus</i>	326	<i>Pericosmus quasimodo</i>	365
<i>Paurocidaridae</i>	277, 278	<i>Pericosmus scaevus</i>	365
<i>Paurocidaris</i>	223, 276, 278	<i>Pericosmus torus</i>	365
<i>Paurocidaris rinbianchi</i>	223	<i>Pericosmus (Lambertona)</i>	365
<i>Paurocidaroida</i>	275, 277	<i>Perisalenia</i>	247
<i>pavyi, Hyboclypeus?</i>	308	<i>Perisalenia hemicidaroides lenoiri</i>	247
<i>pavyi, Psephechinus</i>	261	<i>Perischoechinoidea</i>	204
<i>pawsoni, Asterostoma</i>	393	<i>Peronella</i>	295
<i>pawsoni, Austrocidaris</i>	212	<i>Peronella changchihkengensis</i>	295
<i>paynei, Phymosoma</i>	257	<i>Peronella lesueuri augusta</i>	295
<i>pectenis, Hemiaster</i>	357	<i>Peronella mai</i>	295
<i>Pedina</i>	240	<i>Peronella miaoliensis</i>	295
<i>Pedina antiqua</i>	240	<i>Peronella ova</i>	295
<i>Pedina davoustiana mercieri</i>	240	<i>Peronella toukoshanensis</i>	295
<i>Pedina resecta</i>	240	<i>Peronella tschimanica</i>	295
<i>Pedina sinaica</i>	253	<i>peroni, Echinocardium</i>	392
<i>Pedinacea</i>	239	<i>perplexa, Lambertona</i>	366
<i>Pedinidae</i>	239	<i>perplexus, Phymechinus?</i>	261
<i>Pedinoida</i>	239	<i>perspective, Encope</i>	302
<i>Pedinopsis</i>	252	<i>peruviensis, Triassicidaris</i>	224

<i>Petalobrissus</i>	321	<i>Pironaster</i>	284
<i>Petalobrissus rawdahensis</i>	321	<i>pitouensis, Astriclypeus</i>	303
<i>Petalobrissus tadjikistanensis</i>	322	<i>pitouensis, Clypeaster</i>	288
<i>Petalobrissus (Paralampas)</i>		<i>pitouensis, Taiwanaster</i>	307
<i>platisternus</i>	322	<i>placenta, Prionocidaris</i>	215
<i>Petalobrissus (Paralampas)</i>		<i>plafkeri, Echinarachnius</i>	299
<i>rawdahensis</i>	321	<i>Plagiobrissus</i>	386
<i>petalodes planus, Botriopygus</i>	315	<i>Plagiochasma</i>	313
<i>pfaifferi, Levicidaris</i>	225	<i>Plagiochasma lammersmaxi</i>	313
<i>Phalacrocidaris</i>	213	<i>Plagiochasma nadorense</i>	313
<i>Phalacrocidaris catherinae</i>	213	<i>Plagionotus</i>	386
<i>phelani, Holoctypus</i>	280	<i>Plagionotus holmesii</i>	386
<i>phiale partita, Echinosisgra</i>	348	<i>Plagionotus ravenelianus</i>	387
<i>Philicidaris</i>	227, 229	<i>plana, Cidaris</i>	220
<i>Philicidaris blainvillei</i>	229	<i>plana, Leptosalenia</i>	246
<i>Philicidaris desnoyersi</i>	229	<i>plana, Salenia</i>	246
<i>Philipaster</i>	290	<i>plananyensis, Novasalenia</i>	246
<i>Philipaster morgani</i>	290	<i>plantus, Nucleolites (Pygorhynchus)</i>	322
<i>philipi, Protenaster</i>	372	<i>plantus, Progongrochanus</i>	322
<i>Philolectypus</i>	280	<i>planulata abbassi, Maretia</i>	387
<i>Philolectypus sarthacensis</i>	280	<i>planulatus, Eupatagus</i>	384
<i>Pholiocidaris</i>	207	<i>planus, Cardioichnus</i>	396
<i>Pholiocidaris hungaricus</i>	207	<i>planus, Hemiaster</i>	357
Phormosomatidae	234	<i>planus, Sinaechinocyamus</i>	306
<i>Phyllacanthus</i>	215	<i>planus, Urechinus</i>	345
<i>Phyllacanthus priscus</i>	215	<i>platisternus, Petalobrissus</i>	
<i>Phyllacanthus suleimani</i>	216	<i>(Paralampas)</i>	322
<i>Phyllacanthus</i>	225	<i>platornatus, Cyclaster</i>	383
<i>Phymapatagus</i>	390	<i>Platybrissus</i>	390
<i>Phymechinus</i>	258, 261	<i>platypetalus, Nucleopygus</i>	325
<i>Phymechinus? perplexus</i>	261	<i>Platypygus</i>	323
<i>Phymosoma</i>	257	<i>Platysalenia</i>	247
<i>Phymosoma dalmiapurensis</i>	257	<i>Platysalenia dux</i>	247
<i>Phymosoma maastrichtensis</i>	257	<i>Plegiocidaris</i>	211, 223, 227
<i>Phymosoma mongraensis</i>	257	<i>Plegiocidaris pichaueriensis</i>	228
<i>Phymosoma paynei</i>	257	<i>Plegiocidaris? gemmata</i>	223
<i>Phymosoma princeps</i>	256	<i>pleracantha, Cidaris</i>	227
Phymosomatidae	256	<i>pleracantha, Oedematocidaris</i>	227
Phymosomatoida	256	<i>Plesiaster</i>	379
<i>Phymotaxis</i>	258	<i>Plesiaster gibberiensis</i>	379
<i>pichaueriensis, Nenoticidaris</i>	228	<i>Plesiaster tschaltauensis</i>	379
<i>pichaueriensis, Plegiocidaris</i>	228	<i>Plesiaster vancouverensis</i>	379
<i>pictaviense, Holoctypus</i>	280	<i>Plesiechinus</i>	241
<i>Pilematechinus</i>	344	<i>Plesiechinus altus</i>	241
<i>Pilematechinus belyaevi</i>	344	<i>Plesiechinus hawkinsi</i>	241
<i>pineau, Farquharsonia</i>	241	<i>Plesiocidaroida</i>	275
<i>pipurensis, Echinolampas</i>	318	<i>Plesiocorys</i>	340

<i>Plesiocorys basiprocta</i>	340	<i>Porterpygus kieri</i>	330
<i>Plesiocorys transiens</i>	340	<i>portjacksonensis</i> , <i>Schizaster</i> (<i>Ova</i>)	374
<i>Plesiohemipneustes</i>	340	<i>poslavskae</i> , <i>Eoholaster</i>	336
<i>Plesiohemipneustes revestensis</i>	340	<i>posteroocrassa curtata</i> , <i>Echinolampas</i>	317
Plesiolampadidae	331	<i>posteroexpansus</i> , <i>Pygurus</i>	310
<i>Plesiolampas</i>	331	<i>Pourtalesia</i>	346
<i>Plesiosalenia</i>	247	<i>Pourtalesia heptneri</i>	346
<i>Plesiosalenia depressa</i>	247	<i>Pourtalesia jeffreysi gibbosa</i>	346
<i>Plesiozonus</i>	394	<i>Pourtalesia jeffreysi lata</i>	347
<i>Plesiozonus tenuis</i>	394	<i>Pourtalesia thomsoni</i>	347
<i>Pleurodiadema</i>	257	<i>Pourtalesia vinogradovae</i>	347
<i>Pleurodiadema raboefi</i>	257	Pourtalesiidae	345
<i>Pleurosalenia</i>	246	<i>pouyannei</i> , <i>Desoricidaris</i>	214
Plexechinidae	344	<i>pouyannei</i> , <i>Rhabdocidaris</i>	214
<i>Plexechinus</i>	344	<i>powersi</i> , <i>Agassizia</i>	367
<i>Plexechinus nordenskjoldi</i>	344	<i>praeacutus</i> , <i>Faujasia</i>	320
<i>Plexechinus sulcatus</i>	344	<i>praecursor</i> , <i>Infulaster</i>	339
<i>pliocenicus</i> , <i>Brochopleurus</i>	268	<i>praematurus</i> , <i>Palaechinus</i>	208
Pliolampadidae	328	<i>Praepholidocidaris pellaensis</i>	205
<i>Pliotoxaster</i>	355	<i>praerogalae</i> , <i>Micraster</i>	378
<i>Pliotoxaster angustisulcus</i>	355	<i>Prandinia</i>	247
Polycidaridae	229	<i>Prandinia interpunctata</i>	247
<i>Polycidaris</i>	211, 229	<i>preaustralis</i> , <i>Protenaster</i>	372
<i>Polycidaris nusplingensis</i>	211	<i>predbojensis</i> , <i>Novasalenia</i>	246
<i>Polycyphus</i>	261	<i>pregratilla</i> , <i>Tripneustes</i>	273
<i>Polycyphus arabicus</i>	261	<i>Prenaster</i>	371
<i>Polycyphus parvituberculatus</i>	261	<i>Prenaster desori caucasica</i>	371
<i>Polydiadema</i>	253, 254, 255, 259	<i>Prenaster loveni</i>	369
<i>Polydiadema ambiguum</i>	253	<i>Prenaster synapticus</i>	371
<i>Polydiadema bosei</i>	253	Prenasteridae	366
<i>Polydiadema korotkovi</i>	253	<i>pretensis</i> , <i>Leptosalenia</i>	246
Polydiademataidae	259	<i>pretensis</i> , <i>Salenia</i>	246
Polydiadematinae	254	<i>Primipygus</i>	313
<i>polygonalis</i> , <i>Hemiaster</i> (<i>Leymeriaster</i>)	360	<i>Primipygus richardsoni</i>	313
<i>Pomeliosoma</i>	258	<i>princeps</i> , <i>Gauthiosoma</i>	256
<i>Pomeliosoma duplicatum</i>	258	<i>princeps</i> , <i>Phymosoma</i>	256
<i>Pomeliosoma legayi</i>	258	<i>Prionocidaris</i>	215, 225
<i>popeae</i> , <i>Prionocidaris</i>	215	<i>Prionocidaris callista</i>	215
Poriocidarini	211	<i>Prionocidaris cookei</i>	215
<i>Poriocidaris</i>	211	<i>Prionocidaris katherinae</i>	215
<i>Porosoma</i>	258	<i>Prionocidaris neglecta</i>	216
<i>Porosoma fifei</i>	258	<i>Prionocidaris placenta</i>	215
<i>Porosoma kaspicum</i>	258	<i>Prionocidaris popeae</i>	215
<i>porphyrocardius</i> , <i>Pericosmus</i>	365	<i>Prionocidaris? emiratus</i>	215
<i>porrecta</i> , <i>Echinostigma</i>	348	<i>prisca</i> , <i>Taimanawa</i>	387
<i>Porterpygus</i>	330	<i>priscus</i> , <i>Phyllacanthus</i>	215
<i>Porterpygus devlinensis</i>	330	<i>priscus</i> , <i>Toxaster</i>	355

<i>Proacroechinoidea</i>	275	<i>Protoscutella</i>	297
<i>Proacrolusia</i>	334	<i>Protoscutella mississippiensis</i>	
<i>Proacrolusia kelatensis</i>	334	<i>rosehillensis</i>	297
<i>Procassidulus</i>	323	Protoscutellidae	297
<i>Procidaris</i>	229	<i>Prowillungaster</i>	290
<i>profundi brevispina, Salenia</i>	244	<i>Prowillungaster major</i>	290
<i>profundi intermedius, Salenia</i>	244	<i>Prowillungaster minor</i>	290
<i>profundi megalospina, Salenia</i>	244	<i>Psammechinus</i>	274
<i>profundus, Paradoxechinus</i>	270	<i>Psammechinus carolinensis</i>	274
<i>Progongrochanus</i>	322	<i>Psammechinus exoletus</i>	274
<i>Progongrochanus ariyalurensis</i>	322	<i>Psephechinus</i>	261
<i>Progongrochanus bellus</i>	322	<i>Psephechinus pavyi</i>	261
<i>Progongrochanus crassus</i>	322	<i>Psephoaster</i>	363
<i>Progongrochanus oldhaminus</i>	322	<i>Psephoaster apokryphos</i>	363
<i>Progongrochanus plantus</i>	322	<i>Psephoaster klydonos</i>	363
<i>Progongrochanus testudo</i>	322	<i>Psephoaster lissos</i>	363
<i>Proisaster</i>	354	<i>Pseudanachys</i>	341
<i>Proisaster coramandeli</i>	354	<i>Pseudanachys alpina</i>	341
<i>Pronechinus</i>	206	<i>Pseudanachys tumida</i>	341
<i>Pronechinus cretensis, Pronechinus</i>	206	<i>Pseudholaster</i>	341, 343
<i>Pronucleolites</i>	307	<i>Pseudholaster baconicus</i>	341
<i>Pronucleolites orbicularis</i>	307	<i>Pseudholaster blackdownensis</i>	341
<i>Pronucleolites? crickleyensis</i>	307	<i>Pseudholaster caseyi</i>	341
<i>Prophyllacanthus</i>	216	<i>Pseudholaster depressus</i>	341
<i>Prophyllacanthus eocenicus</i>	216	<i>Pseudholaster mangyschlakensis</i>	341
<i>Prophyllacanthus leoni</i>	216	Pseudholasterinae	343
<i>Proraster</i>	371, 376	<i>Pseudoastrodaspis ilyinensis</i>	299
<i>Proraster granti</i>	372	<i>Pseudocatopygus</i>	319
<i>Proraster magnus</i>	372	<i>Pseudocatopygus bucharensis</i>	319
<i>Proraster oedumi</i>	371	Pseudocidaridae	251
<i>Prosalenia</i>	243	<i>Pseudocidaris</i>	251
<i>Prosalenia marcouii</i>	243	<i>Pseudocidaris depressa</i>	251
<i>prosorovskyi, Codechinus</i>	260	<i>Pseudocidaris raratuberculata</i>	251
<i>Protenaster</i>	372	<i>Pseudocidaris romani</i>	251
<i>Protenaster philipi</i>	372	<i>Pseudocidaris santacruzensis</i>	251
<i>Protenaster preaustralis</i>	372	<i>Pseudodiadema</i>	253
Proterocidaridae	208	<i>Pseudodiadema bakonyense</i>	253
<i>Proterocidaris</i>	208	<i>Pseudodiadema collenoti</i>	233
<i>Protobrissus</i>	377	<i>Pseudodiadema vadeti</i>	253
<i>Protobrissus akkajensis</i>	380	Pseudodiadematidae	251
<i>Protobrissus indolensis</i>	380	<i>pseudodistinctus, Epiaster</i>	353
<i>Protobrissus rionensis</i>	377, 380	<i>pseudodistinctus oblongus, Epiaster</i>	353
<i>Protocardia</i>	340	<i>pseudodistinctus rotundiformis, Epiaster</i>	353
<i>Protocardia cotteauanus</i>	340	<i>pseudofilograna, Romanocidaris</i>	229
<i>Protocardia truncates</i>	340	<i>Pseudogibbaster</i>	380
<i>protodecimae, Ovulaster</i>	379	<i>Pseudogibbaster akkajensis</i>	380
<i>protoplanus, Cardia</i>	334	<i>Pseudogibbaster depressus</i>	380

<i>Pseudogibbaster indolensis</i>	380	<i>Pygorhynchus debensis</i>	314
<i>Pseudogibbaster rionensis</i>	377, 380	<i>Pygorhynchus ovalis</i>	284
<i>Pseudogibbaster tercensis</i>	380	Pyguridae	310
<i>Pseudoholaster</i> [sic!] <i>mangyschlakensis</i>	341	<i>Pygurus</i>	310
<i>Pseudolinthia</i>	372, 376	<i>Pygurus andinus</i>	310
<i>Pseudolinthia triporata</i>	372	<i>Pygurus arabicus</i>	310
<i>pseudonodulosus</i> , <i>Holaster</i>	338	<i>Pygurus beineuensis</i>	310
<i>pseudoplanus</i> , <i>Cardiaster</i>	334	<i>Pygurus geryvillensis</i>	309
<i>Pseudopygaulus</i>	329, 331	<i>Pygurus jurensis</i>	309
<i>Pseudopygaulus malavoyi</i>	329	<i>Pygurus mendelsohni</i>	310
<i>Pseudopygaulus romani</i>	329	<i>Pygurus posteroexpansus</i>	310
<i>Pseudopyrina subcircularis</i>	284	<i>Pygurus rostratus</i>	309
<i>Pseudopyrina subovalis</i>	284	<i>Pygurus tenuis</i>	309
<i>Pseudosalenia</i>	243	<i>Pygurus yamamaensis</i>	310
<i>Pseudosalenia magniprocta</i>	243	<i>Pyrgorhytis</i>	333
<i>Pseudosalenia malogostiana</i>	243	<i>Pyrgorhytis klairisauzei</i>	333
Pseudosaleniidae	243	<i>Pyrgorhytis magnus</i>	333
<i>Pseudosorella</i>	313	<i>Pyrina</i>	283
<i>Pseudovulechinus</i>	286	<i>Pyrina azemati</i>	283
<i>Pseudowashitaster</i>	363	<i>Pyrina houzeaui</i>	284
<i>Pseudowashitaster japonicus</i>	363	<i>Pyrina montainvillensis</i>	284
<i>Pseudowashitaster mysticus</i>	363	<i>Pyrina neolaevis</i>	283
<i>psiloses</i> , <i>Hemicidaris</i>	250	<i>Pyrina sphaerica</i>	283
Psychocidaridae	225	<i>Pyrinodia</i>	287, 331
<i>pulchellus</i> , <i>Brissopatagus</i>	382	<i>qattamiaensis</i> , <i>Echinolampas</i>	318
<i>pulchellus</i> , <i>Stigmatopygus?</i>	322	<i>qenaensis</i> , <i>Bandelicyphus</i>	268
<i>pulchellus</i> , <i>Zardinechinus</i>	224	<i>qishnensis</i> , <i>Leptosalenia</i>	246
<i>pulcher</i> , <i>Hemiaster</i>	358	<i>quadralis</i> , <i>Limpasiaster</i>	324, 325
<i>pulchra</i> , <i>Linthia</i>	370	<i>quadrialata</i> , <i>Cidaris</i>	220
<i>pulchrum</i> , <i>Hapalosoma</i>	235	<i>quadriserrata</i> , <i>Cidaris</i>	220
<i>Pygaster microstoma</i>	307	<i>quasimodo</i> , <i>Pericosmus</i>	365
Pygasteridae	241	<i>quaylei</i> , <i>Ambipleurus?</i>	267
Pygasteroida	239, 241	<i>quinqueradialis</i> , <i>Roderosignus</i>	397
<i>Pygaulus</i>	313	<i>quinquetuberculata</i> , <i>Mimiosalenia</i>	266
<i>Pygaulus aksuensis</i>	313	<i>raboeufi</i> , <i>Pleurodiadema</i>	257
<i>Pygaulus cyclotus</i>	313	<i>Rachiosoma</i>	258
<i>Pygaulus faasi</i>	313	<i>Rachiosoma gigasei</i>	258
<i>Pygaulus richardi</i>	313	<i>radeti</i> , <i>Romanocidaris</i>	229
<i>Pygomalus</i>	308	<i>radians</i> , <i>Hyposalenia</i>	248
<i>Pygomalus azemati</i>	308	<i>raingeardi</i> , <i>Echinogalerus</i>	287
<i>Pygopistes</i>	314	<i>rajasthanensis</i> , <i>Eupatagus</i>	384
<i>Pygopistes parrasae</i>	314	<i>rancureli</i> , <i>Paralampas</i>	326
<i>Pygopyrina</i>	283	<i>ranjitpurensis</i> , <i>Maretia</i>	387
<i>Pygopyrina thibaudi</i>	283	<i>raoi</i> , <i>Echinocyamus</i>	292
<i>Pygorhynchus</i>	314	<i>raratuberculata</i> , <i>Pseudocidaris</i>	251
<i>Pygorhynchus acutus</i>	314	<i>rarus</i> , <i>Hemiaster (Mecaster)</i>	360
<i>Pygorhynchus clypeiformis</i>	314	<i>raskhaimahensis</i> , <i>Emiratia</i>	255

<i>raskhaimahensis</i> , <i>Emiratus</i>	255	<i>Rhabdocidaris reginae</i>	231
<i>rathieri</i> , <i>Angusticypeus</i>	308	<i>Rhabdocidaris remus</i>	231
<i>rathieri</i> , <i>Clypeus</i>	308	<i>Rhabdocidaris schneideri</i>	231
<i>ratiparensis</i> , <i>Amblypygus</i>	279	<i>Rhabdocidaris varusensis</i>	230
<i>ravenelianus</i> , <i>Plagionotus</i>	387	<i>rhabdoses</i> , <i>Hemicidaris</i>	250
<i>ravni</i> , <i>Globator</i>	284	<i>Rhyncholampas</i>	326
<i>ravni</i> , <i>Neoglobator</i>	284	<i>Rhyncholampas candidoi</i>	326
<i>rawdahensis</i> , <i>Petalobrissus</i>	321	<i>Rhyncholampas chipolanus</i>	326
<i>rawdahensis</i> , <i>Petalobrissus</i> (<i>Paralampas</i>)	321	<i>Rhyncholampas gouldii newbernensis</i>	327
<i>rebouli</i> , <i>Reboulicidaris</i>	226	<i>Rhyncholampas smithi</i>	327
<i>Reboulicidaris</i>	226	<i>Rhyncholampas strougoi</i>	327
<i>Reboulicidaris rebouli</i>	226	<i>Rhyncholampas (Galerolampas)</i>	327
<i>reginae</i> , <i>Rhabdocidaris</i>	231	<i>Rhyncholampas (Galerolampas)</i> <i>tinrhertensis</i>	327
<i>reguanti</i> , <i>Echinanthus</i>	331	<i>Rhynchopygus</i>	327
<i>regularis</i> , <i>Zardinechinus</i>	224	<i>Rhynchopygus arumaensis</i>	327
<i>reingardae</i> , <i>Palaeodiadema</i>	237	<i>Rhynchopygus? janchrisorum</i>	327
<i>relicta</i> , <i>Eurhodia</i>	328	<i>Rhynobrissus</i>	387
<i>relictus</i> , <i>Brissopatagus</i>	381	<i>Rhynobrissus tumulus</i>	387
<i>remanei</i> , <i>Toxaster</i>	355	<i>richardi</i> , <i>Pygaulus</i>	313
<i>Remondella</i>	300	<i>richardsoni</i> , <i>Nucleolites</i>	313
<i>Remondella asiatica</i>	300	<i>richardsoni</i> , <i>Primipygus</i>	313
<i>Remondella kamtschatica</i>	301	<i>Rictocystis</i>	349
<i>Remondella waldroni</i>	300	<i>Rictocystis jensenae</i>	349
<i>remus</i> , <i>Rhabdocidaris</i>	231	<i>rimaporus Bromidechinus</i>	204
<i>Remycidaris</i>	229	<i>rinbianchi</i> , <i>Cidaris</i>	220, 223
<i>Remycidaris elegans</i>	229	<i>rinbianchi</i> , <i>Paurocidaris</i>	223
<i>Remycidaris largiporus</i>	229	<i>rindensis</i> , <i>Schizaster</i>	373
<i>reniformis</i> , <i>Cardioichnus</i>	396	<i>rionensis</i> , <i>Protobrissus</i>	377, 380
<i>renngarteni</i> , <i>Heteraster</i>	354	<i>rionensis</i> , <i>Pseudogibbaster</i>	377, 380
<i>resecta</i> , <i>Pedina</i>	240	<i>Rispolia</i>	342
<i>resecta</i> , <i>Thieulinipedina</i>	240	<i>Rispolia hungarica</i>	342
<i>reticulata</i> , <i>Cidaris</i>	220	<i>roberti</i> , <i>Palmerius</i>	223
<i>reticulatus</i> , <i>Ovulaster</i>	379	<i>robertsi</i> , <i>Heliocidaris</i>	274
<i>Retzneiosoma</i>	235	<i>rochatensis</i> , <i>Hemiaster</i>	357
<i>Retzneiosoma jasenecki</i>	235	<i>Roderosignus</i>	397
<i>revestensis</i> , <i>Holaster</i>	340	<i>Roderosignus quinqueradialis</i>	397
<i>revestensis</i> , <i>Plesiohemipneustes</i>	340	<i>rojasi</i> , <i>Haimea</i>	285
<i>rex</i> , <i>Hemifaorina?</i>	369	<i>rollandi oblongus</i> , <i>Hemiaster</i>	358
<i>Rhabdocidaridae</i>	230	<i>Rolliericidaris</i>	227
<i>Rhabdocidarinae</i>	214	<i>Rolliericidaris alternans</i>	227
<i>Rhabdocidaris</i>	230, 231	<i>Rolliericidaris etalloni</i>	227
<i>Rhabdocidaris bigoti</i>	230	<i>romani</i> , <i>Epiaster</i>	352
<i>Rhabdocidaris blainvillei</i>	229	<i>romani</i> , <i>Linthia</i>	371
<i>Rhabdocidaris impar</i>	231	<i>romani</i> , <i>Pseudocidaris</i>	251
<i>Rhabdocidaris lahuseni</i>	231	<i>romani</i> , <i>Pseudopygaulus</i>	329
<i>Rhabdocidaris pouyannei</i>	214	<i>romani</i> , <i>Scutella</i>	297

<i>romani</i> , <i>Tremaster</i>	271	<i>Salenia dux</i>	247
<i>Romanocidaris</i>	227, 229	<i>Salenia humei</i>	246
<i>Romanocidaris dietli</i>	229	<i>Salenia mathuri</i>	246
<i>Romanocidaris pseudofilograna</i>	229	<i>Salenia mexicana</i>	246
<i>Romanocidaris radeti</i>	229	<i>Salenia microprocta</i>	244
Roseicidaridae	231	<i>Salenia nudispina</i>	244
<i>Roseicidaris</i>	231, 232	<i>Salenia pentagonalis</i>	245
<i>Roseicidaris morieri</i>	232	<i>Salenia plana</i>	246
Roseicidaroida	231	<i>Salenia pretensis</i>	246
<i>rostrata</i> , <i>Echinocrepis</i>	346	<i>Salenia profundi brevispina</i>	244
<i>rostrata</i> , <i>Taimanawa</i>	387	<i>Salenia profundi intermedius</i>	244
<i>rostratus</i> , <i>Hemiaster</i>	357	<i>Salenia profundi megalospina</i>	244
<i>rostratus</i> , <i>Kieripygurus</i>	309	<i>Salenia scutigera hungarica</i>	245
<i>rostratus</i> , <i>Pygurus</i>	309	<i>Salenia sigillata pozaryskae</i>	245
<i>rostratus zitteli</i> , <i>Eupatagus</i>	384	<i>Salenia tandoni</i>	244
<i>rostriformis</i> , <i>Hypsopygaster</i>	324	<i>Salenia texana</i>	246
<i>Rostropygus</i>	319	<i>Salenia trisuranalis</i>	244
<i>Rostropygus annae</i>	319	<i>Salenia volana</i>	246
Rotulidae	291	<i>Salenia wilmingtonensis</i>	245
<i>rotundatus</i> , <i>Ovulechinus</i>		<i>Salenia (Pleurosalenia) vanbirgeleni</i>	246
(<i>Pseudovulechinus</i>)	286	<i>Salenidia</i>	247
<i>rotundus</i> , <i>Echinolampas</i>	318	<i>Salenidia sanctipetri</i>	247
<i>rougeriei</i> , <i>Schizaster (Paraster)</i>	375	Saleniidae	243
<i>rowdahensis</i> , <i>Vologesia</i>	328	Salenioida	241
<i>rowei anglicus</i> , <i>Boletechinus</i>	262	<i>Salenocidaris</i>	247, 248
<i>ruegensis</i> , <i>Brissopneustes</i>	377	<i>Salenocidaris gallemii</i>	247
<i>rügensis</i> , <i>Brissopneustes</i>	377	<i>salgadoi</i> , <i>Nucleopygus</i>	325
<i>rugius</i> , <i>Offaster</i>	339	<i>salviensis</i> , <i>Desoricidaris</i>	214
<i>rugosa depressa</i> , <i>Eurhodia</i>	328	<i>Samlandaster</i>	305
<i>rugosa ideali</i> , <i>Eurhodia</i>	328	<i>Samlandaster dineuri</i>	305
<i>rupellensis</i> , <i>Codiopsis</i>	264	<i>sanctaecrucis</i> , <i>Desoricidaris</i>	214
<i>ryukyuensis</i> , <i>Stylocidaris</i>	224	<i>sanctipetri</i> , <i>Salenidia</i>	247
<i>saemanni</i> , <i>Abelleicidaris</i>	228	<i>sanglakensis</i> , <i>Holaster</i>	338
<i>saemanni</i> , <i>Cidaris</i>	228	<i>santacruzensis</i> , <i>Pseudocidaris</i>	251
<i>safiense</i> , <i>Tylocidaris</i>	226, 228	<i>santolayae</i> , <i>Cassidulus</i>	324
<i>Salenia</i>	244	<i>Sardocidaris</i>	227
<i>Salenia ammonitorum</i>	245	<i>sarthacensis</i> , <i>Holectypus</i>	280
<i>Salenia arabica</i>	246	<i>sarthacensis</i> , <i>Philolectypus</i>	280
<i>Salenia banatica</i>	245	Sarthostomata	287
<i>Salenia batnensis</i>	248	<i>scaevus</i> , <i>Pericosmus</i>	365
<i>Salenia baylissi</i>	245	Scaphechinidae	305
<i>Salenia bella</i>	245	<i>Scaphechinus</i>	305
<i>Salenia bella parva</i>	245	<i>Scaphechinus brykovi</i>	305
<i>Salenia cascadenensis</i>	244	<i>schafferi</i> , <i>Echinocorys</i>	335
<i>Salenia cherei</i>	245	<i>schamchorensis</i> , <i>Guettaria</i>	337
<i>Salenia danubiana</i>	245	<i>scherzeri</i> , <i>Schizaster</i>	373
<i>Salenia depressa</i>	247	<i>schickleri</i> , <i>Aguayoaster</i>	367

<i>Schizaster</i>	372	<i>Scutella aegyptiaca</i>	296
<i>Schizaster alphonsei</i>	372	<i>Scutella conica</i>	297
<i>Schizaster archiaci complanatus</i>	373	<i>Scutella kalksburgensis</i>	297
<i>Schizaster caddoensis</i>	373	<i>Scutella marchettii</i>	297
<i>Schizaster dyscritus</i>	369	<i>Scutella muelleri</i>	297
<i>Schizaster formelli</i>	373	<i>Scutella norris</i>	301
<i>Schizaster halei</i>	373	<i>Scutella romani</i>	297
<i>Schizaster halli</i>	373	<i>Scutella vindobonensis altus</i>	297
<i>Schizaster humei</i>	373, 375	<i>scutellaris, Willungaster</i>	290, 291
<i>Schizaster rindensis</i>	373	Scutellidae	296
<i>Schizaster scherzeri</i>	373	Scutellidea	296
<i>Schizaster stenzeli</i>	373	<i>Scutellina</i> (Genus)	294
<i>Schizaster tarimensis</i>	373	<i>Scutellina</i> (Suborder)	296
<i>Schizaster</i> (<i>Dipneustes</i>)	374	<i>Scutellina morgani</i>	290
<i>Schizaster</i> (<i>Dipneustes</i>) <i>fosteri</i>	374	<i>Scutellina supramarginalis</i>	294
<i>Schizaster</i> (<i>Ova</i>)	374	<i>Scutellinoides</i>	289, 291
<i>Schizaster</i> (<i>Ova</i>) <i>myorensis</i>	374	Scutellinoididae	291
<i>Schizaster</i> (<i>Ova</i>) <i>portjacksonensis</i>	374	<i>scutigera hungarica, Salenia</i>	245
<i>Schizaster</i> (<i>Paraster</i>) <i>ashrafi</i>	374	<i>secoensis, Encope</i>	302
<i>Schizaster</i> (<i>Paraster</i>) <i>carinatus</i>	375	<i>seelandica, Cidaris</i>	220
<i>Schizaster</i> (<i>Paraster</i>) <i>extumidus</i>	375	<i>seelandica misurinensis, Cidaris</i>	220
<i>Schizaster</i> (<i>Paraster</i>) <i>incorrectus</i>	375	<i>semseyana, Kieria</i>	304
<i>Schizaster</i> (<i>Paraster</i>) <i>jhadwensis</i>	374	<i>sergipensis, Acriaster</i>	323
<i>Schizaster</i> (<i>Paraster</i>) <i>ovatus</i>	374	Serpianotiaridae	233, 276
<i>Schizaster</i> (<i>Paraster</i>) <i>rougeriei</i>	375	<i>Serpianotiaris</i>	233
<i>Schizaster</i> (<i>Paraster</i>) <i>susana</i>	375	Serpianotiaroida	233
<i>Schizaster</i> (<i>Paraster</i>) <i>tatei</i>	375	<i>serradentata, Cidaris</i>	222
Schizasteridae	366	<i>setosumdepressa, Diadema</i>	237
<i>Schizechinus</i>	272	<i>Seunaster</i>	342
<i>Schizechinus pentagonus</i>	272	<i>Seunaster bulgaricus</i>	342
<i>Schizobrissus</i>	386	<i>Sexpyga</i>	266
<i>Schizocosmus</i>	375, 376	<i>Sexpyga soyoae</i>	266
<i>Schizocosmus abatoides</i>	375	<i>seymourensis, Austrocidaris</i>	212
<i>schlueteri, Temnocidaris</i>	214	<i>shaimae, Brissus</i>	382
<i>schmidiobtusa, Galerites</i> (<i>Pironaster</i>)	284	<i>sibiricus, Hemiaster</i>	357
<i>schmidtae, Parahemiaster</i>	362	<i>sicula, Histocidaris</i>	211
<i>schmidti, Stegaster</i> (<i>Seunaster</i>)	342	<i>siculum, Acanthotrema</i>	393, 394
<i>schneideri, Rhabdocidaris</i>	231	<i>siculum, Kermabrissoides</i>	394
<i>schréteri, Archaeocidaris</i>	209	<i>sidibouzidense, Hemicidaris</i>	250
<i>schultzi, Echinolampas</i>	316	<i>sigillata pozaryskae, Salenia</i>	245
<i>schweinfurthi, Desoricidaris</i>	214	<i>silesiaca, Hirudocidaris</i>	213
<i>schweinfurthi devriesi, Echinanthus</i>	332	<i>silesiaca, Stereocidaris</i>	213
<i>schwetzovi, Echinocorys</i>	335	<i>silleroi, Brissopatagus</i>	382
<i>scillae eythraea, Temnotrema</i>	270	<i>simakovi, Hemiaster</i>	357
<i>scutata forma pyrenaica, Echinocorys</i>	337	<i>similis, Braunechinus</i>	277
<i>scutataa forma ovata, Echinocorys</i>	335	<i>similis, Cidaris</i>	277
<i>Scutella</i>	296	<i>Sinaechinocyamus</i>	306

<i>Sinaechinocyamus planus</i>	306	<i>sphaerica</i> , <i>Pyrina</i>	283
<i>sinaica</i> , <i>Pedina</i>	253	<i>sphaericus</i> , <i>Crassiholaster</i>	334
<i>sinaica</i> , <i>Pedinopsis</i> (<i>Sinaiopsis</i>)	253	<i>sphaericus</i> , <i>Echinocorys</i>	334
<i>Sinaiopsis</i>	253	<i>Sphaeropleurus</i>	273
<i>sinensis</i> , <i>Conulus</i>	282	<i>Sphaeropleurus geometricus</i>	273
<i>sinensis</i> , <i>Domechinus</i>	319	<i>Sphenaster</i>	376
<i>singhi</i> , <i>Eupatagus</i>	384	<i>Sphenaster larumbensis</i>	376
<i>singhi</i> , <i>Meoma</i> (<i>Schizobrissus</i>)	386	<i>sphericus</i> , <i>Hattopsis</i>	265
<i>sinica</i> , <i>Tetradia</i>	291	<i>spongiosa</i> , <i>Cidaris</i>	220, 221
<i>sinuosum</i> , <i>Aspidodiadema</i>	237	<i>stanleyi</i> , <i>Zardinechinus</i>	225
<i>smiseri</i> , <i>Codiopsis</i>	264	<i>Stegaster</i>	342
<i>smithi</i> , <i>Rhyncholampas</i>	327	<i>Stegaster palaeocenicus</i>	342
<i>Smithiaster</i>	332	<i>Stegaster</i> (<i>Seunaster</i>) <i>schmidti</i>	342
<i>Smithiaster loryi</i>	332	<i>stella</i> , <i>Hemiaster</i>	362
<i>Smithicidaris</i>	227	<i>stella</i> , <i>Parahemiaster</i>	362
<i>Smithicidaris jacquemeti</i>	227	<i>stellarum</i> , <i>Gnathichnus</i>	397
<i>smithii</i> , <i>Cidaris</i>	227	<i>stellatus</i> , <i>Paradoxechinus</i>	270
<i>smithii</i> , <i>Nenoticidaris</i>	227	<i>stenzeli</i> , <i>Schizaster</i>	373
<i>snellingsi</i> , <i>Diplotagma</i>	260	<i>Sterechinus</i>	274
<i>Sogdiaster</i>	380	<i>Sterechinus bernasconiae</i>	274
<i>Sogdiaster turkestanensis</i>	378, 380	Stereocidarinae	213
<i>solemi</i> , <i>Bothriocidaris</i>	204	<i>Stereocidaris</i>	213, 214
<i>Solenocystis</i>	349	<i>Stereocidaris silesiaca</i>	213
<i>Solenocystis imitans</i>	349	<i>Stereocidaris trigonodus</i>	213
<i>solignaci</i> , <i>Micraster</i>	378	<i>sternalis jousseaumei</i> , <i>Metalia</i>	388
<i>soluni</i> , <i>Parahemiaster</i>	362	<i>sternalis lata</i> , <i>Metalia</i>	388
Somaliasteridae	350	Stigmatopyginae	323
<i>sonorensis</i> , <i>Micraster</i> (<i>Gibbaster</i>)	379	<i>Stigmatopygus</i>	322, 323
<i>soyuae</i> , <i>Sexpyga</i>	266	<i>Stigmatopygus?</i> <i>pulchellus</i>	322
Spatangidae	380, 387	<i>stoliczkai</i> , <i>Gongrochanus</i>	320
<i>spatangiformis</i> , <i>Huttonechinus</i>	343	Stomechinidae	259, 262
<i>spatangiformis</i> , <i>Macropneustes</i>	343	Stomechinoida	262
<i>Spatangobrissus</i>	389	<i>Stomechinus</i>	261
<i>Spatangobrissus dermodyorum</i>	389	<i>Stomechinus tinginatus</i>	262
<i>Spatangobrissus incus</i>	389	<i>Stomechinus?</i> <i>dissimilaris</i>	261
Spatangoida	351	<i>stomias</i> , <i>Echinocorys</i>	336
<i>Spatangus</i>	389	<i>strawberryensis</i> , <i>Aarchaeocidaris</i>	209
<i>Spatangus mathesoni</i>	389	<i>strictiporus</i> , <i>Nucleolites</i>	312
<i>Spatangus subglobosus</i>	334	Strongylocentridae	275
<i>Spatangus truncates</i>	340	<i>Strongylocentrotus franciscanus</i>	275
<i>Spatangus</i> (<i>Phymapatagus</i>)	390	<i>Strongylocentrotus nudus</i>	275
<i>Spatangus</i> (<i>Phymapatagus</i>)		<i>strougoi</i> , <i>Rhyncholampas</i>	327
<i>grandituberculatus</i>	390	<i>Studeria</i>	329
<i>Sperosoma</i>	235, 236	<i>Studeria?</i> <i>frescoensis</i>	329
<i>Sperosoma nudum</i>	235	<i>Stylocidaris</i>	224
Sperosomatinae	236	<i>Stylocidaris ryukyuensis</i>	224
<i>sphaerica</i> , <i>Pedinopsis</i>	252	<i>subameliae</i> , <i>Hemiaster</i>	357

<i>subbibansensis</i> , <i>Hemiaster</i>	357	<i>tainui</i> , <i>Opissaster</i>	361
<i>subcircularis</i> , <i>Neoglobator</i>	284	<i>Taiwanaster</i>	306
<i>subcircularis</i> , <i>Pseudopyrina</i>	284	<i>Taiwanaster gutingkengensis</i>	306
<i>subcircularis sulcatus</i> , <i>Catopygus</i>	311	<i>Taiwanaster mai</i>	306
<i>subconicus</i> , <i>Hemiaster</i>	358	<i>Taiwanaster pitouensis</i>	307
<i>subglobosus</i> , <i>Crassiholaster</i>	334	Taiwanasteridae	306
<i>subglobosus</i> , <i>Spatangus</i>	334	Taiwanasteritida	306
<i>subglobosus</i> forma <i>norfolkensis</i> ,		<i>Tamililampus</i>	329
<i>Crassiholaster</i>	334	<i>Tamililampus tumidus</i>	329
<i>subidus</i> , <i>Hemiaster</i> (<i>Bolbaster</i>)	359	<i>tamilnaduensis</i> , <i>Gongrochanus</i>	320
<i>sublaevis</i> , <i>Cidaris</i>	228	<i>tanamensis</i> , <i>Holaster</i>	338
<i>sublaevis</i> , <i>Nudicidaris</i>	228	<i>tandoni</i> , <i>Echinolampas</i>	317
<i>submirabilis</i> , <i>Hemiaster</i>	358	<i>tandoni</i> , <i>Eupatagus</i>	384
<i>subovalis</i> , <i>Neoglobator</i>	284	<i>tandoni</i> , <i>Gymnopatagus</i>	384
<i>subovalis</i> , <i>Pseudopyrina</i>	284	<i>tandoni</i> , <i>Salenia</i>	244
<i>subquadratus</i> , <i>Botriopygus</i>	315	<i>tanypetalis</i> , <i>Echinolampas</i>	318
<i>subquadratus</i> , <i>Holaster</i>	338	<i>tarimensis</i> , <i>Schizaster</i>	373
<i>subrotundus</i> , <i>Parahemiaster</i>	362	<i>tasmanicus</i> , <i>Brisaster</i>	368
<i>subsimilis</i> , <i>Hemiaster</i> (<i>Malwaster</i>)	360	<i>tatei</i> , <i>Schizaster</i> (<i>Paraster</i>)	375
<i>subsimilis</i> , <i>Opissaster</i>	360	<i>tauricus elatior</i> , <i>Clypeaster</i>	289
<i>subtrigonalis</i> , <i>Douvillaster</i>	351	<i>teetotumensi</i> , <i>Bahariya</i>	391
<i>subvenulosa</i> , <i>Desoricidaris</i>	214	<i>Temnocidaris</i>	214
<i>suemegensis</i> , <i>Antiquibrissus</i>	380, 381	<i>Temnocidaris schlueteri</i>	214
<i>suemegensis</i> , <i>Botriopygus</i>	315	Temnopleuridae	268
<i>suessi</i> , <i>Magnosia</i>	266	Temnopleuroida	267
<i>suessi</i> , <i>Magnosiopsis</i>	266	<i>Temnotrema</i>	270
<i>sulcata</i> , <i>Cidaris</i>	220	<i>Temnotrema greifatensis</i>	270
<i>sulcata cognatus</i> , <i>Echinocorys</i>	335	<i>Temnotrema scillae eythraea</i>	270
<i>sulcata orbiculata</i> , <i>Echinocorys</i>	335	<i>Temnotrema wizrensis</i>	270
<i>sulcatus</i> , <i>Plexechinus</i>	344	<i>Temnotrema xishaensis</i>	270
<i>sulciproctus</i> , <i>Crassiholaster</i>	334, 335	<i>templetoni</i> , <i>Neobothriocidaris</i>	205
<i>suleimani</i> , <i>Phyllacanthus</i>	216	<i>tenuicostata</i> , <i>Cidaris</i>	220
<i>suleiovense</i> , <i>Trochotiara</i>	254	<i>tenuis</i> , <i>Kieripygurus</i>	309
<i>sümegensis</i> , <i>Antiquibrissus</i>	381	<i>tenuis</i> , <i>Plesiozonus</i>	394
<i>sümegensis</i> , <i>Botriopygus</i>	315	<i>tenuis</i> , <i>Pygurus</i>	309
<i>summesbergeri</i> , <i>Linthia?</i>	370	<i>teraniensis</i> , <i>Hemicidaris</i>	250
<i>supramarginalis</i> , <i>Scutellina</i>	294	<i>tercensis</i> , <i>Micraster</i>	380
<i>susana</i> , <i>Schizaster</i> (<i>Paraster</i>)	375	<i>tercensis</i> , <i>Pseudogibbaster</i>	380
<i>sveshnikovii</i> , <i>Eopericosmus</i>	366	<i>Termieria</i>	331
<i>sylviae</i> , <i>Centrostephanus</i>	237	<i>terminata</i> , <i>Echinocorys</i>	336
<i>synapticus</i> , <i>Prenaster</i>	371	<i>tessieri</i> , <i>Thylechinus</i>	258
<i>tachanabatensis</i> , <i>Echinolampas</i>	318	<i>testudo</i> , <i>Nucleolites</i> (<i>Pygorhynchus</i>)	322
<i>tachilniensis</i> , <i>Kewia</i>	299	<i>testudo</i> , <i>Progongrochanus</i>	322
<i>tadjikistanensis</i> , <i>Petalobrissus</i>	322	<i>Tetradiella</i>	291
<i>Taimanawa</i>	387	<i>Tetradiella sinica</i>	291
<i>Taimanawa prisca</i>	387	<i>tetraedrica</i> , <i>Cidaris</i>	220
<i>Taimanawa rostrata</i>	387	<i>tetraedrica appendiculata</i> , <i>Cidaris</i>	220

<i>Tetragramma</i>	254, 259	<i>Toxaster jugamis</i>	355
<i>Tetragramma gloriae</i>	254	<i>Toxaster laffittei</i>	336
<i>Tetragramma variolare baconicum</i>	254	<i>Toxaster priscus</i>	355
<i>Tetraromania</i>	333	<i>Toxaster remanei</i>	355
<i>Tetraromania jaccardi</i>	333	Toxasteridae	351
<i>Tetraromania ovulum</i>	333	Toxasterina	351
<i>texana, Leptosalenia</i>	246	Toxopneustidae	272
<i>texana, Salenia</i>	246	<i>Trachyaster aichinoi</i>	361
<i>texanus, Eupatagus</i>	384	<i>Trachypatagus</i>	390
<i>thibaudi, Pygopyrina</i>	283	<i>Trachypatagus nehalaie</i>	390
<i>thieli, Genus?</i>	278	<i>transiens, Plesiocorys</i>	340
<i>thieli, Kieritiaris</i>	276	<i>trapezoidalis, Cidaris</i>	221
<i>thieli, Thielicidaris</i>	232	<i>Tremaster</i>	271
Thielicidaridae	232	<i>Tremaster romani</i>	271
<i>Thielicidaris</i>	232	<i>trempinus, Tylocidaris (Sardocidaris)</i>	227
<i>Thielicidaris thieli</i>	232	<i>Tretocidaris</i>	224
Thielicidaroida	232	<i>Tretocidaris anguillensis</i>	224
<i>thieulini, Diplopodia</i>	252	Triadocidaridae	276
<i>Thieulinipedita</i>	240	<i>Triadocidaris</i>	210, 276
<i>Thieulinipedita antiqua</i>	240	<i>Triadocidaris hungarica</i>	210
<i>Thieulinipedita resecta</i>	240	Triadocidaroida	275, 276
<i>Tholaster</i>	342	Triadotiaridae	232
<i>Tholaster carvalhoi</i>	342	<i>Triadotiaris</i>	232, 233
<i>Tholaster fourmarieri</i>	343	<i>Triadotiaris grandaevus</i>	233
<i>thomsoni, Pourtalesia</i>	347	Triadotiaroida	232
<i>Thylechinus</i>	258	<i>Triassicidaris</i>	224
<i>Thylechinus tessieri</i>	258	<i>Triassicidaris peruviansis</i>	224
<i>Thylechinus vanderhami</i>	258	<i>Tridium</i>	296
<i>Tiaromma</i>	259	<i>Tridium kieri</i>	296
<i>Tigilella</i>	298	<i>Trigonocidaris</i>	271
<i>tiliensis, Echinodiscus</i>	304	<i>Trigonocidaris (Lamprechinus) nitidus</i>	
<i>tinginatus, Stomechinus</i>	262	<i>nascaensis</i>	271
<i>tinrhertensis, Rhyncholampas</i>		<i>Trigonocidaris (Tuberculocidaris)</i>	271
<i>(Galerolampas)</i>	327	<i>Trigonocidaris (Tuberculocidaris)</i>	
<i>toi, Kupeia</i>	361	<i>tuberculata</i>	271
<i>tokiensis, Mizunamia</i>	395	<i>trigonodus, Stereocidaris</i>	213
<i>torus, Pericosmus</i>	365	<i>Tripneustes</i>	273
<i>toucasi, Cidaris</i>	230	<i>Tripneustes gratilla elatensis</i>	273
<i>toucasi, Zbindenicidaris</i>	230	<i>Tripneustes misrai</i>	273
<i>toukoshanensis, Peronella</i>	295	<i>Tripneustes pregratilla</i>	273
<i>Toxaster</i>	354	<i>triporata, Pseudolinthia</i>	372
<i>Toxaster bajarunasi</i>	355	<i>Tripylus</i>	375
<i>Toxaster cauveriae</i>	354	<i>Tripylus beatriceae</i>	375
<i>Toxaster compressa</i>	354	<i>trisuranalis, Salenia</i>	244
<i>Toxaster compressa</i>	354	<i>Trochalosoma</i>	262
<i>Toxaster crassissulcatus</i>	355	<i>Trochalosoma gharamulensis</i>	262
<i>Toxaster granosus kouensis</i>	355	<i>Trochodiadema</i>	255, 259

<i>Trochodiadema? dhofarensis</i>	256	<i>Urechinus aoteanus</i>	345
<i>Trochotiara</i>	254	<i>Urechinus drygalskii</i>	344
<i>Trochotiara kiiensis</i>	254	<i>Urechinus drygalskii perfidus</i>	345
<i>Trochotiara kongieli</i>	254	<i>Urechinus mortenseni</i>	344, 345
<i>Trochotiara moabitorum</i>	254	<i>Urechinus parvus</i>	345
<i>Trochotiara suleiovensis</i>	254	<i>Urechinus planus</i>	345
<i>Tromikosoma</i>	236	<i>uruaensis, Coraster</i>	350
<i>truncatus, Protocardia</i>	340	<i>usakensis, Neoprora</i>	371
<i>truncatus, Spatangus</i>	340	<i>usarpi, Aporocidaris</i>	211
<i>tschaltauensis, Plesiaster</i>	379	<i>vadeti, Pseudodiadema</i>	253
<i>tschimanica, Peronella</i>	295	<i>valparolae, Cidaris</i>	221
<i>tuberculata, Amoraster</i>	380	<i>valvaedentata, Echinogutta</i>	349
<i>tuberculata, Trigonocidaris</i>		<i>vanbirgeleni, Salenia (Pleurosalenia)</i>	246
(<i>Tuberculocidaris</i>)	271	<i>vancouverensis, Plesiaster</i>	379
<i>Tuberculocidaris</i>	271	<i>vanderhami, Thylechinus</i>	258
<i>tumida, Pseudanachys</i>	341	<i>vannoordenburgi, Brissalius</i>	381
<i>tumidus, Tamillilampus</i>	329	<i>Vaquerosella</i>	301
<i>tumulus, Echinolampas</i>	316	<i>Vaquerosella coreyi</i>	301
<i>tumulus, Rhynobrissus</i>	387	<i>variolare baconicum, Tetragramma</i>	254
<i>turanica, Echinocorys</i>	335	<i>varusensis, Rhabdocidaris</i>	230
<i>turkestanensis, Hemiaster</i>	358	<i>varusensis, Zbindenicidaris</i>	230
<i>turkestanensis, Micra</i>	380	<i>venulosa, Desoricidaris</i>	214
<i>turkestanensis, Sogdiaster</i>	378, 380	<i>venustespinosa, Cidaris</i>	222
<i>Tylocidaris</i>	226	<i>verecundus, Hemiaster (Bolbaster)</i>	359
<i>Tylocidaris bermudezi</i>	226	<i>verticillata, Cidaris</i>	221
<i>Tylocidaris hemmoorensis</i>	226	<i>vialovi, Domechinus</i>	319
<i>Tylocidaris inexpectata</i>	226	<i>Viaudechinus</i>	271
<i>Tylocidaris safiense</i>	226, 228	<i>Viaudechinus bigoti</i>	271
<i>Tylocidaris (Sardocidaris) trempinus</i>	227	<i>viladensis, Ambipleurus</i>	267
<i>typicus, Eopericosmus</i>	366	<i>vilanovae, Coraster</i>	350
<i>typicus, Himalayechinus</i>	321	<i>vincentinus, Apatopygus</i>	330
<i>Typocidaridae</i>	231	<i>vincentinus, Echinobrissus</i>	330
<i>Typocidaris</i>	214, 231	<i>Vinchuscanchaia</i>	276
<i>Typocidaris ubaghisi</i>	214	<i>Vinchuscanchaia kieri</i>	276, 277
<i>ubaghisi, Typocidaris</i>	214	<i>vindobonensis, Brissopneustes</i>	376
<i>ultimus, Collyropsis</i>	332	<i>vindobonensis altus, Scutella</i>	297
<i>ulugqatensis, Coelopleurus</i>	264	<i>vinogradovae, Pourtalesia</i>	347
<i>ungaensis, Echinarachnius</i>	299	<i>vityazi, Echinosisgra</i>	349
<i>Unibothriocidaris</i>	205	<i>volana, Leptosalenia</i>	246
<i>Unibothriocidaris bromidensis</i>	205	<i>volana, Salenia</i>	246
<i>Unibothriocidaris kieri</i>	205	<i>volgensis, Echinobrissus</i>	312
<i>uniformis, Cidaris</i>	213	<i>volgensis, Nucleolites</i>	312
<i>uniformis, Hirudocidaris</i>	213	<i>Vologesia</i>	328
<i>uniporum, Palaeotropus</i>	394	<i>Vologesia rowdahensis</i>	328
<i>Urechinidae</i>	344	<i>vulcani, Bothriocidaris</i>	204
<i>Urechinus</i>	345	<i>vulgaris minor, Galerites</i>	284
<i>Urechinus antipodeanus</i>	345	<i>wachangpoensis, Cidaris</i>	222

<i>waechteri</i> , <i>Braunechinus</i>	277	<i>Zardinitiaris</i>	276
<i>waechteri</i> , <i>Cidaris</i>	277	<i>Zardinitiaris flexuosa</i>	276
<i>waiwulunensis</i> , <i>Astriclypeus</i>	304	<i>Zbindenicidaris</i>	229, 230
<i>waldroni</i> , <i>Remondella</i>	300	<i>Zbindenicidaris toucasi</i>	230
<i>Washitaster japonicus</i>	363	<i>Zbindenicidaris varusensis</i>	230
<i>Waurnia</i>	376	<i>zejszneri</i> , <i>Echinocorys</i>	336
<i>Waurnia nelsoni</i>	376	<i>Zeuglopleuridae</i>	273
<i>wayensis</i> , <i>Hemiaster</i>	358	<i>Zeuglopleurus</i>	272, 273
<i>weaveri</i> , <i>riamaster</i>	298	<i>Zeuglopleurus dictyopleuroides</i>	272
<i>wehrlii</i> , <i>Nannoglyphus</i>	269	<i>Zeuglopleurus wehrlii</i>	269
<i>wehrlii</i> , <i>Zeuglopleurus</i>	269	<i>zircensis</i> , <i>Heteraster</i>	353
<i>westlakei</i> , <i>Micraster</i>	378	<i>zonarius</i> , <i>Epiaster</i>	352
<i>wetmorei</i> , <i>Encope</i>	302	<i>zschokkei</i> , <i>Cidaris</i>	228
<i>whatleyensis</i> , <i>Archaeocidaris</i>	210	<i>zschokkei</i> , <i>Nenoticidaris</i>	228
<i>willeibi</i> , <i>Orthopsis</i>	278	<i>zunyii</i> , <i>Lovenechinus</i>	207
<i>Willungaster</i>	289, 290		
<i>Willungaster scutellaris</i>	290, 291		
<i>wilmingtonensis</i> , <i>Salenia</i>	245		
<i>wilmingtonica inflata</i> , <i>Agassizia</i> (<i>Anisaster</i>)	367		
<i>wilsoni</i> , <i>Echinocyamus</i>	292		
<i>wilsoni</i> , <i>Eupatagus</i>	385		
<i>Wissmannechinus</i>	277		
<i>Wissmannechinus haussmanni</i>	277		
<i>wizrensis</i> , <i>Fibularia</i>	293		
<i>wizrensis</i> , <i>Temnotrema</i>	270		
<i>wrighti</i> , <i>Allomma</i>	255		
<i>wrightii</i> , <i>Chesniericidaris</i>	217		
<i>wrightii</i> , <i>Cidaris</i>	217		
<i>xanthe</i> , <i>Lissocidaris</i>	223		
<i>xinjiangensis</i> , <i>Macropneustes</i> (<i>Deakia</i>)	386		
<i>xishaensis</i> , <i>Temnotrema</i>	270		
<i>yadongensis</i> , <i>Echinolampas</i>	319		
<i>yadongensis</i> , <i>Ovulechinus</i>	286		
<i>yamamaensis</i> , <i>Pygurus</i>	310		
<i>yangi</i> , <i>Palaechinus</i>	208		
<i>yeliuensis</i> , <i>Astriclypeus</i>	303		
<i>yeliuensis</i> , <i>Echinodiscus</i>	304		
<i>yeni</i> , <i>Clypeaster</i>	289		
<i>yingkoensis</i> , <i>Clypeaster</i>	289		
<i>Zardinechinus</i>	224, 276, 277		
<i>Zardinechinus giulini</i>	224		
<i>Zardinechinus lancedelli</i>	224		
<i>Zardinechinus pulchellus</i>	224		
<i>Zardinechinus regularis</i>	224		
<i>Zardinechinus stanleyi</i>	225		
<i>zardinia</i> , <i>Levicidaris</i>	225		

