

Complementary references: TOLLMANN (1977), SCHÖNLAUB (1979, 1980a, 2001), EBNER et al. (1989), HEINISCH & SCHÖNLAUB (1993).

Erzführende Kalke / Metalliferous Limestones

FRITZ EBNER

Validity: Invalid; general working term for not subdivided Devonian limestones in the E-GWZ.

Type area: ÖK50-UTM, map sheet 4215 Eisenerz (ÖK 50-BMN, map sheets 101 Eisenerz and 131 Kalwang), ÖK50-UTM, map sheet 4211 Neuberg an der Mürz (ÖK50-BMN, map sheet 103 Kindberg).

Type section: -

Reference section(s): -

Derivation of name: After siderite-ankerite mineralizations which are frequently hosted by mainly Devonian limestones (CZERMAK, 1931; TOLLMANN, 1977; WEBER, 1997a, b).

Synonyms: See remarks.

Lithology: Variegated bedded flaser limestones hosting irregular metasomatic stocks of siderite and ankerite mineralizations (WEBER, 1997a, b).

Fossils: See remarks.

Origin, facies: See remarks.

Chronostratigraphic age: Devonian; see remarks.

Biostratigraphy: See remarks.

Thickness: See remarks.

Lithostratigraphically higher rank unit: -

Lithostratigraphic subdivision: -

Underlying unit(s): -

Overlying unit(s): -

Lateral unit(s): -

Geographic distribution: E-GWZ; Styria, Eisenerzer Alpen.

Remarks: Since the beginning of geologic research in the E-GWZ (STUR, 1865, 1866) the general term Metalliferous Limestones was used cumulatively for mainly Devonian limestones which include irregular stocks of metasomatic iron mineralizations. Later parts of these limestones, especially in the Eisenerzer Alpen, were assigned as informal lithostratigraphic units (SCHÖNLAUB 1979, 1980a, 1982a, b; FLAJS & SCHÖNLAUB, 1976). In the Veitsch area of the E-GWZ (ÖK50-UTM, map sheet 4211 Neuberg an der Mürz; ÖK50-BMN, map sheet 103 Kindberg) isolated outcrops of Metalliferous Limestones above the Rad Schists (NIEVOLL, 1983, 1987) yielded conodont faunas representing the entire Devonian period (EBNER, 1973, 1974; NIEVOLL 1983, 1987).

Complementary references: TOLLMANN (1977), EBNER et al. (1989), HEINISCH & SCHÖNLAUB (1993).

Veitsch Nappe

In the ASC 2004 the stratigraphic sequence of the Veitsch Nappe is shown in the Tournaisian to Moskovian parts of the column "E-Grauwackenzone" (E-Greywacke Zone) right of the Eisenerz Formation.

The Veitsch Nappe is situated in the E-GWZ (Styria, Lower Austria) at the tectonic footwall of the Greywacke Zone and is composed of Carboniferous (lower Visean–Moskovian) marine shallow water sediments. They were only affected by Alpine (Cretaceous) deformation and low grade metamorphism (RATSCHBACHER, 1984, 1987; NEUBAUER et al., 1994; RANTITSCH et al., 2004; EBNER et al., 2007, 2008). NEUBAUER et al. (1994) proposed to summarize these sediments within the Veitsch Group (not shown in the ASC 2004).

In the western part of the E-GWZ the sequence of the Veitsch Nappe was formerly subdivided into three formations: Steilbachgraben Formation, Triebenstein Formation, and Sunk Formation (RATSCHBACHER, 1984). In a restricted area SE Kapfenberg the Sunk Formation is superposed by (?) Permian ochre shales and sandstones (= Graschnitz Formation; NEUBAUER, 1983; NEUBAUER et al., 1994). It is not clear if RATSCHBACHER's subdivision into formations is also applicable for the eastern parts of the Veitsch Nappe. Nevertheless, magnesite and dark clastic sediments rich in graphite and plants resembling the Sunk Formation also occur in the eastern parts of the Veitsch Nappe. Especially in the magnesite deposit of Veitsch dolomites and clastics below the magnesite are rich in fossils (corals, brachiopods, crinoids, trilobites, gastropods, agglutinated foraminifers, spicula, ostracods; KOCH, 1893; KLEBELSBERG, 1927; HERITSCH, 1928a, 1930a; METZ, 1937; FELSER & FLÜGEL, 1975; HAHN & HAHN, 1977; KRAINER, 1992). Trilobites indicate lower Visean (HAHN & HAHN, 1977) and corals upper Visean *Dibunophyllum* Zone (FELSER & FLÜGEL, 1975; FELSER, 1977), however, a correlation with the sequence of Hohentauern/Sunk is problematic.

The abundance of magnesite and graphite is also responsible to assign some parts of the Veitsch Nappe as the "Magnesite Carboniferous" (EBNER, 1997) and the "Graphite Carboniferous" (e.g., HAMMER, 1924). The Carboniferous of the Veitsch Nappe represents post-orogenic sediments (in relation to an early Carboniferous [= Bretonic] tectonic phase) deposited in a shallow marine foredeep (FLÜGEL, 1977; NEUBAUER & VOZAROVA, 1990; EBNER, 1992; EBNER et al., 1991, 2007, 2008). The metasomatic magnesite deposits, all most probably situated in the Steilbachgraben Formation or its equivalents, form the Veitsch Nappe magnesite (talc) district. The Sunk Formation hosts the Veitsch Nappe graphite district (WEBER, 1977a, b).

Steilbachgraben-Formation / Steilbachgraben Formation

FRITZ EBNER

Validity: Valid; formal description by RATSCHBACHER (1984).

Type area: Rottenmanner Tauern; ÖK50-UTM, map sheet 4214 Trieben (ÖK50-BMN, map sheet 130 Trieben).

Type section: Overlapping parts of the type section are situated around the abandoned magnesite mine NW of Hohentauern (N 47°26'53" / E 14°27'59"). Sections 8–11 (RATSCHBACHER, 1984: Figs. 1–3) are situated at the NE-ridge of Sunkmauer (altitude 1,180 m), the southern margin of the magnesite open pit, in the upper Steilbachgraben (N 47°26'22" / E 14°29'57" to N 47°26'26" / E 14°30'06") and within the magnesite open pit.

Reference section(s): -

Austrian Stratigraphic Chart 2004 - Paleozoic

(sedimentary successions)

