

Hochwipfel Formation

CLAUDIA SPALLETTA, CORRADO VENTURINI, HANS-PETER SCHÖNLAUB & MONICA PONDRELLI

Österreichische Karte 1:50.000

Blatt BMN 197 Kötschach

Blatt BMN 198 Weißbriach

Blatt BMN 199 Hermagor

Blatt UTM 3109 Oberdrauburg

Blatt UTM 3110 Kötschach-Mauthen

Blatt UTM 3116 Sonnenalpe Naßfeld

Blatt UTM 3117 Nötsch im Gailtal

Carta Topografica d'Italia 1:50.000

Foglio 018 Passo di Monte Croce Carnico

Foglio 031 Ampezzo

Foglio 032 Tolmezzo

Foglio 033 Tarvisio

Definition

Quartz-rich sandstone, siltstone and shale with at places interbedded ruditic horizons, commonly at the base of the unit, and rare acid volcanic bodies (keratophyres).

Description


The Hochwipfel Formation can be subdivided into 4 lithofacies:

a – Quartz-arenite to lithic sandstone, siltstone and shale in beds with thickness from few to 30 cm with turbiditic features. The lithology division A, B, C and D of the Bouma sequence (Bouma turbidite facies model, according to the classification of WALKER & MUTTI (1973), are frequently recognized within lithofacies a. Locally the lithic sandstone yield fossil plant remains.


b – Silicatic rudite, breccia and conglomerate made mainly by radiolarian chert clasts, and rare clasts of sandstone and siltstone, very rare basic volcanics and quartz clasts are also present. The siliciclastic rudite may reach thickness of 30-40 m.

c – Limestone rudite mainly constitutes by clasts derived from the Upper Devonian limestone, and subordinate radiolarian chert, sandstone and siltstone. The limestone rudite are sporadically interbedded within the lower part of the formation, their thickness varies from few dm to 3 m. At places limestone clasts rich in crinoid debris (encrinites) of late Visean age are present.

d – Acid volcanic and sub-volcanic (tuffs) products (Keratophyres: GORTANI, 1906; GENTILI & PELLIZZER, 1964; VAI in BRAGA et al., 1971).



Areas of outcrop of the Hochwipfel Formation with indication of the stratotype of the lower and upper boundary (asterisks): A: Rio Chianaletta Section; B: Collina-Tors Section; C: Givigliana Section. Reference section (square): D: Pic Chiadin section.



The Hochwipfel Formation type section. a) log of the Rio Chianaletta Section (modified after PERRI & SPALLETTA, 1998), and view of the section in the field (photo G.B. VAI); b) log of the Collina–Tors and Givigliana Sections (after SPALLETTA & VENTURINI, 2009).

Fossil content

Plant remains and spores rarely found within the turbiditic sequence of lithofacies a (AMEROM & SCHÖNLAUB, 1992). One of the localities richer in flora was studied by AMEROM et al. (1984). Other findings of plant remains are located at Mt. Bruca and in the Pramosio area.

Depositional environment

Pelagic, in open marine basin with local strong tectonic control (Flysch: VAI, 1963) (SPALLETTA & VENTURINI, 1995; VENTURINI & SPALLETTA 1998).

Stratotype

Rio Chianaletta Section (PERRI & SPALLETTA, 1998), at coordinates N 46°36'13.7", E 12°50'02.8" for the lower boundary. Collina–Tors and Givigliana Sections (SPALLETTA & VENTURINI, 2009), for the upper boundary, at coordinates respectively N 46°34'36", E 12°49'13" and N 46°34'19", E 12°49'21".

Reference sections

Pic Chiadin section (SPALLETTA & VENTURINI, 1988), at coordinates N 46°35'52", E 12°54'2", where the lithofacies b is well exposed and the arenite yields plant remains.

Type area

Carnic Alps.

Main outcrop areas

The Hochwipfel Formation crops out along the whole Carnic Alps from the Bordaglia Valley to the west to the Mt. Ois-ternig to the east.

Thickness

The estimated thickness of the Hochwipfel Formation is less than 1000 m; measured thickness varies from 350 to 600 m.

Boundaries

Underlying units – Zollner Formation (conformable gradual contact), Pal Grande Formation (paraconformable–unconformable contact affected by tectonic superimposition).

Overlying units – Dimon Formation (conformable gradual contact), Late- and Post-Variscan units (Permo-Carbonifero Pontebbano and Val Gardena Formation) (unconformable contact: Variscan angular unconformity).

Lateral units – Dimon Formation.

Derivation of name

After Mount Hochwipfel.

Synonymy

'complesso argilloso-scistoso carbonifero': GORTANI (1913).

Hochwipfelfazies: HERITSCH (1928).

Hochwipfelschichten: KAHLER & METZ (1955).

'Flysch ercinico' [partim]: VAI (1963).

Formazione dell'Hochwipfel: SELLI (1963); VAI in BRAGA et al. (1971).

Formazione del Hochwipfel: SPALLETTA et al. (1980, 1982); GERMANI (2007); SPALLETTA & VENTURINI (2009).

Hochwipfel-Formation: SCHÖNLAUB (1985).

Hochwipfel Formation: BRIME et al. (2008).

Hochwipfel-Formation/Hochwipfel Formation: SUTTNER (2014).



Views of the Hochwipfel Formation in the field. a) sandstones, siltstones and shales of lithofacies a, outcrop along the Valentin Valley (photo C. VENTURINI); b) silicatic breccia of lithofacies b, near Casera Plotta (photo C. SPALLETTA); c) limestone rudite of lithofacies c, near Creta di Collina (photo C. SPALLETTA); d) silicatic breccia of lithofacies b, near Casera Plotta (photo F. SGORBINO).

Chronostratigraphic age

Carboniferous: Middle Viséan (AMEROM et al., 1984) to lower Bashkirian.

Biostratigraphy -

Complementary references -

Remarks -

References

- AMEROM, H.W.J. van & SCHÖNLAUB, H.P. (1992): Pflanzenfossilien aus dem Karbon von Nötsch und der Hochwipfel-Formation der Karnischen Alpen (Österreich). – *Jahrbuch der Geologischen Bundesanstalt*, **135/1**, 195–216, Wien.
- AMEROM, H.W.J. van, FLAJS, G. & HUNGER, G. (1984): Die „Flora der Marinelli-Hütte“ (Mittleres Visé) aus dem Hochwipfelflysch der Karnischen Alpen (Italien). – *Mededelingen Rijks Geologische Dienst*, **37/1**, 1–41, Heerlen.
- BRAGA, G.P., CARLONI, G.C., COLANTONI, P., CORSI, M., CREMONINI, P., FRASCARI, F., LOCATELLI, D., MONESI, A., PISA, G., SASSI, F.P., SELLI, R., VAI, G.B. & ZIRPOLI, G. (1971): Note illustrative della Carta geologica d'Italia. Foglio 4c-13 M. Cavallino-Ampezzo. – Ministero dell'Industria, del Commercio e dell'Artigianato, Direzione Generale delle Miniere, Servizio Geologico d'Italia, 108 p., Roma.
- BRIME, C., PERRI, M.C., PONDRELLI, M., SPALLETTA, C. & VENTURINI, C. (2008): Polyphase metamorphism in the eastern Carnic Alps (N Italy-S Austria): Clay minerals and conodont Colour Alteration Index evidence. – *International Journal of Earth Sciences*, **97/6**, 1213–1229, Berlin-Heidelberg.
- GENTILI, G. & PELLIZER, R. (1964): Le rocce eruttive del Paleozoico carnico. – *Bollettino della Società Geologica Italiana*, **83/1**, 151–205, Roma.
- GERMANI, D. (2007): Formazione del Hochwipfel. – Carta geologica d'Italia 1: 50.000, Catalogo delle Formazioni, Quaderni del Servizio Geologico d'Italia, Serie III – 7(VI), Agenzia per la Protezione dell'Ambiente e per i Servizi Tecnici, 15–20, Roma.
- GORTANI, M. (1906): Studi sulle rocce eruttive delle Alpi Carniche. – *Memorie Società Toscana Scienze Naturali*, **22**, 166–198, Pisa.
- GORTANI, M. (1913): Rilevamento del nucleo centrale carnico. – *Bollettino Reale Comitato geologico Italiano*, **43/2**, 71–75, Roma.
- HERITSCH, F. (1928): Tektonische Fragen im Karbon der Karnischen Alpen. – *Sitzungsberichte der Akademie der Wissenschaften Wien*, **137/1**, 303–338, Wien.
- KAHLER, F. & METZ, K. (1955): Die Hochwipfelschichten der Karnischen Alpen. – *Verhandlungen der Geologischen Bundesanstalt*, **1955**, 253–260, Wien.
- PERRI, M.C. & SPALLETTA, C. (1998): Conodont distribution at the Tournaisian/Viséan boundary in the Carnic Alps (Southern Alps, Italy). – In: SZANIAWSKI, H. (ed.): *Proceedings of the Sixth European Conodont Symposium (ECOS VI)*. – *Palaeontologia Polonica*, **58**, 225–245, Warszawa.
- SCHÖNLAUB, H.P. (1985): Das Paläozoikum der Karnischen Alpen. – In: SCHÖNLAUB, H.P. (ed.): *Arbeitstagung der Geologischen Bundesanstalt 1985 Kötschach-Mauthen, Gailtal* – Geologische Bundesanstalt, 34–52, Wien.
- SELLI, R. (1963): Schema geologico delle Alpi Carniche e Giulie occidentali. Scala 1:100.000. – *Giornale di Geologia*, **30**, 1–136, Bologna.
- SPALLETTA, C. & VENTURINI, C. (1988): Conglomeratic Sequences in the Hochwipfel Formation: A New Palaeogeographic Hypothesis on the Hercynian Flysch Stage of the Carnic Alps. – *Jahrbuch der Geologischen Bundesanstalt*, **131/4**, 637–647, Wien.
- SPALLETTA, C. & VENTURINI, C. (1995): Late Devonian-Early Carboniferous synsedimentary tectonic evolution of the Palaeocarnic domain (Southern Alps, Italy). – *Giornale di Geologia*, **56/2**, 1994, 211–222, Bologna.
- SPALLETTA, C. & VENTURINI, C. (2009): Formazione del Hochwipfel. – In: VENTURINI, C. (ed.): *Note Illustrative del Foglio 031 Ampezzo*. – Carta Geologica d'Italia alla scala 1:50000, Istituto Superiore per la Protezione e la Ricerca Ambientale (ex-Agenzia per la Protezione dell'Ambiente e per i Servizi Tecnici, Servizio Geologico d'Italia), 48–51, Stampa A.T.I. – S.EL.CA. srl. – L.A.C. srl. – System Cart srl., Firenze.
- SPALLETTA, C., VAI, G.B. & VENTURINI, C. (1980): Il Flysch ercinico nella geologia dei Monti Paularo e Dimon (Alpi Carniche). – *Memorie della Società Geologica Italiana*, **20**, 243–265, Roma.
- SPALLETTA, C., VAI, G.B. & VENTURINI, C. (1982): La Catena Paleocarnica. – In: CASTELLARIN, A. & VAI, G.B. (eds.): *Guida alla geologia del Sudalpino centro-orientale*. – *Guide Geologiche Regionali, Società Geologica Italiana*, 281–292, Bologna.
- SUTTNER, T.J. (2014): Hochwipfel-Formation/Hochwipfel Formation. – In: PILLER, W.E. (ed.): *The lithostratigraphic units of the Austrian Stratigraphic Chart 2004 (sedimentary successions), Vol. I – The Paleozoic Era(them)*. – *Abhandlungen der Geologischen Bundesanstalt*, **66**, 80–81, Wien.
- VAI, G.B. (1963): Ricerche geologiche nel gruppo del M. Coglians e nella zona di Volaia (Alpi Carniche). – *Giornale di Geologia*, **30**, 137–198, Bologna.
- VENTURINI, C. & SPALLETTA, C. (1998): Remarks on the Palaeozoic stratigraphy and the Hercynian tectonics of the Palaeocarnic Chain (Southern Alps). – In: PERRI, M.C. & SPALLETTA, C. (eds.): *Southern Alps Field Trip Guidebook. Seventh International Conodont Symposium held in Europe*. – *Giornale di Geologia, Special Issue*, **60**, 69–88, Bologna.
- WALKER, R.G. & MUTTI, E. (1973): Turbidite facies and facies associations. – In: MIDDLETON, G.V. & BOUMA, A.H. (eds.): *Turbidites and Deep-Water Sedimentation, SEPM, Pacific Section*, 119–157, Los Angeles.