

The ammonite *Forresteria (Harleites)* from the Santonian Gosau Group of the Randograben (Rußbach am Pass Gschütt, Salzburg), Austria

HERBERT SUMMESBERGER¹, WILLIAM J. KENNEDY², MICHAEL WAGREICH³, PETER SKOUMAL⁴,
 NORBERT SCHWAIGHOFER⁵ & ALFRED LEIBLFINGER⁶

1 Text-Figure, 1 Table, 2 Plates

Österreichische Karte 1:50.000

BMN / UTM

95 Sankt Wolfgang im Salzkammergut / NL 33-01-11 Bad Ischl

NL 33-01-17 Hallstatt

Ammonites

Forresteria

Santonian

Late Cretaceous

Northern Calcareous Alps

Gosau Group

Austria

Contents

Abstract 5
 Zusammenfassung 5
 Introduction and Geological Setting 6
 Repositories of Specimens 6
 Systematic Palaeontology 7
 Discussion 7
 Stratigraphy 7
 Conclusions 8
 Acknowledgements 8
 References 8
 Plates 9

Abstract

Lower Santonian representatives of the ammonite genus *Forresteria* have been collected for the first time from the Streiteck Formation (Gosau Group) of the Randobach section near Rußbach am Pass Gschütt in the Austrian Alps. *Forresteria (Harleites) seidl* nov. sp. is described herein; its co-occurrence with *Texanites quinquenodosus* (REDTENBACHER, 1873), *Eulophoceras natalense* HYATT, 1903, *Texasia cricki* (SPATH, 1921), and *Cladoceramus undulatoplicatus* (ROEMER, 1852) indicates an Early Santonian age.

Der Ammonit *Forresteria (Harleites)* aus dem Santonium der Gosau Gruppe des Randograbens (Rußbach am Pass Gschütt, Salzburg), Österreich

Zusammenfassung

Erstmals wurden Vertreter der Gattung *Forresteria (Harleites)* aus dem unteren Santonium der Streiteck-Formation der Gosau Gruppe des Randobach-Profiles gesammelt: *Forresteria (Harleites) seidl* nov. sp. wird hier beschrieben. Die Begleitfauna von *Texanites quinquenodosus* (REDTENBACHER, 1873), *Eulophoceras natalense* HYATT, 1903, *Texasia cricki* (SPATH, 1921) und *Cladoceramus undulatoplicatus* (ROEMER, 1852) belegt frühsantonies Alter.

1 HERBERT SUMMESBERGER: Museum of Natural History, Burgring 7, 1010 Vienna, Austria. herbert.summesberger@nhm-wien.ac.at
 2 WILLIAM J. KENNEDY: Oxford University Museum of Natural History, Parks Road, Oxford OX1 3PW, UK, and Department of Earth Sciences, Parks Road, Oxford OX1 3AN, UK.
 3 MICHAEL WAGREICH: University of Vienna, Faculty of Earth Sciences, Geography and Astronomy, Institute of Geology, Althanstraße 14, 1090 Vienna, Austria.
 4 PETER SKOUMAL: Bastiengasse 56, 1180 Vienna, Austria.
 5 NORBERT SCHWAIGHOFER: Rußbach am Pass Gschütt, 5442 Salzburg, Austria.
 6 ALFRED LEIBLFINGER: Haus der Natur Salzburg, Museumsplatz 5, 5020 Salzburg, Austria.

Introduction and Geological Setting


Representatives of the Late Cretaceous ammonite genus *Forresteria* are generally considered to be of late Turonian to Coniacian age (e.g. KENNEDY et al., 1983; KENNEDY, 1984; KENNEDY & COBBAN, 1991; IFRIM et al., 2019). *Forresteria* (*Harleites*) *petrocoriensis* (COQUAND, 1859) is widespread in the French Coniacian (KENNEDY, 1984: 49) and used as zonal marker for the Early Coniacian in ammonite zonations of Europe (KENNEDY & COBBAN, 1991: 9), including probably the uppermost Turonian (KENNEDY & WALASZCZYK, 2004; IFRIM et al., 2019). *Forresteria peruana* (BRÜGGEN, 1910) and *Forresteria* (*Forresteria*) *brancoi* (SOLGER, 1904) are common in the Coniacian of South and Central America. *F. (F.) alluaudi* (BOULE, LEMOINE & THÉVENIN, 1907) is known from the Coniacian of Europe, South Africa (Zululand; Coniacian II of KENNEDY & KLINGER, 1975), Madagascar, Japan, the United States Western Interior, Peru, and Mexico (IFRIM et al., 2019).

In the Austrian Alps, *Forresteria* has been poorly known. *Forresteria* (*Forresteria*) *alluaudi* (BOULE, LEMOINE & THÉVENIN, 1907) was described from the Bad Ischl road tunnel section (SUMMESBERGER et al., 2022), where it co-occurs with the Late Coniacian *Paratexanites serratomarginatus* (REDTENBACHER, 1873). In the present study, *Forresteria* (*Harleites*) *seidli* nov. sp. is described for the first time from the Randobach section (Rußbach am Pass Gschütt, Salzburg, Austria) of the Gosau Group of the Northern Calcareous Alps, where it was

collected from the Streiteck Formation (WAGREICH, 1988). In contrast to the nearby assemblage of the Bad Ischl tunnel section, *Forresteria* (*Harleites*) *seidli* nov. sp. does not co-occur with *Paratexanites serratomarginatus*, but with an Early Santonian fauna of *Texanites quinquenodosus* (REDTENBACHER, 1873), *Eulophoceras natalense* HYATT, 1903, *Texasia cricki* (SPATH, 1921), and *Cladoceras undulatoplicatus* (ROEMER, 1852).

Repositories of Specimens

- | | |
|-----|---|
| CE | Seidl collection, Salzburg. |
| SCH | W. and N. Schwaighofer collection, Rußbach am Pass Gschütt, Salzburg. |
| LEI | Leiblfinger collection, Golling, Salzburg. |
| SK | Skoumal collection, Vienna. |



Text-Fig. 1.
Sketch map of the Randobach, with modifications after SUMMESBERGER et al. (2017: Text-Fig. 2).

Systematic Palaeontology

Subclass Ammonoidea Zittel, 1884

Order Ammonitida Hyatt, 1889

Superfamily Acanthoceratoidea de Grossouvre, 1894

Family Collignoniceratidae Wright & Wright, 1951

Subfamily Barroisiceratinae Basse, 1947

Genus *Forresteria* Reeside, 1932

Subgenus *Harleites* Reeside, 1932

The high variability and widespread, yet isolated, occurrences of the genus *Forresteria* REESIDE, 1932 led to the introduction of a number of taxa now regarded as synonyms, which are discussed at length by KENNEDY et al. (1983: 253). The type species is *Barroisicerat (Forresteria) forresteri* REESIDE, 1932: 17, Pl. 5, Figs. 2–7, by the subsequent designation of WRIGHT (1957: 432), which is a synonym of *Acanthoceras (Prionotropis) alluaudi* BOULE, LEMOINE & THÉVENIN, 1907: 12, Pl. 1, Figs. 6, 7 (KENNEDY et al., 1983: 275). The type species of *Harleites* is *Barroisicerat haberfellneri* var. *harléi* DE GROSSOUVRE, 1894: 56, Pl. 2, Figs. 7, 8, by the subsequent designation of KENNEDY et al. (1983: 259), which is a synonym of *Ammonites petrocoriensis* COQUAND, 1859: 995. *Solgerites* REESIDE, 1932 is a synonym of *Forresteria* REESIDE, 1932 (KENNEDY, 1985).

Forresteria (Harleites) seidli nov. sp.

(Pl. 1, Figs. 1–4; Pl. 2, Figs. 1–4)

Etymology: The species is named after Heinz Seidl (Salzburg), who collected the first specimen at the Randobach site (Text-Fig. 1) in 2016.

Material: Five specimens from the locality Rando 3 (Text-Fig. 1): CE.00027; LEI/GO/2019/0017; SCH/RA/2019/01, 02; SK/RA/2020/168 (Tab. 1).

Holotype: Designated herein is CE.00027 (Pl. 1, Figs. 3, 4) from the Seidl collection (Salzburg).

Description: All individuals are internal moulds with shell preserved in all specimens, often retaining the nacreous layer. They are laterally crushed to a certain degree, but measurements are still useful. Coiling is evolute. All specimens comprise both phragmocone and body chamber. Small sections of the suture are visible but impossible to

Inventory	D (mm)	Wh (mm)	Wb (mm)	U (mm)	U (%)
CE.00027	123.3	53.0	33.4	31.6	25.6 %
LEI/GO/2019/0017	98.6	35.7	15.0	30.0	30.4 %
SCH/RA/2019/01	104.4	40.8	22.4	31.9	28.7 %
SCH/RA/2019/02	111.4	45.3	--	35.7	32.0 %
SK/RA/2020/168	67.6	25.9	13.8	18.0	26.6 %

Tab. 1.

Measurements of the Randograben specimens of *Forresteria (Harleites) seidli* nov. sp.

D = diameter; Wh = whorl height; Wb = whorl breadth; U = umbilical diameter; U % = ratio umbilical diameter to specimen diameter.

describe. The whorl height increases rapidly. The umbilicus is wide, the umbilical wall vertical to slightly inclined outwards, towards the umbilical shoulder. The flanks are feebly inflated with the greatest thickness in median position. The sculpture is very coarse with 9 to 11 very strongly pinched tubercles at the umbilical shoulder that give rise to very strong ribs. Most ribs are ornamented by an additional tubercle at the internal (CE.00027) or external (SCH/RA/2019/01) third of the flank and they link to very strong clavi at the ventrolateral shoulder. Most ribs arise in pairs from the umbilical tubercles and there are some intercalatories. In adult specimens, a very feeble median ventral elevation is succeeded by a totally flat venter (Pls. 1, 2).

Discussion

Forresteria (Harleites) seidli nov. sp. is established as a new species on the basis of its wide umbilicus and it is so far the sole occurrence of the genus in the Santonian. Coniacian *Forresteria (Harleites) petrocoriensis* (COQUAND, 1859) (KENNEDY, 1984: 49, Pl. 4, Figs. 1, 2; Pl. 6, Figs. 1–9; Pl. 7, Figs. 1–20; Pl. 9, Figs. 1–4; Pl. 21, Figs. 2, 4; Text-Figs. 13d–f, h, 16, 18a, b, g) differs by its deep and narrow umbilicus. The Upper Coniacian *Forresteria (Forresteria) alluaudi* (BOULE, LEMOINE & THÉVENIN, 1907) (KENNEDY et al., 1983: 276, Text-Figs. 5–9, 10a, b, e–f, 11–14, 15a, b, 16–31, 33, 34, 35c–e, 40d–e) differs also by its narrower umbilicus and by its great variability (KENNEDY & COBBAN, 1991: Pl. 5, Figs. 1–11; SUMMESBERGER et al., 2022) from very coarsely ribbed and tuberculated to flattened and feebly ornamented. *Forresteria (Forresteria) peruana* (BRÜGGEN, 1910: 720, Pl. 27, Fig. 1) differs from *Forresteria (Harleites) seidli* nov. sp. by its tendency towards a smooth flank, by preservation of a row of mediolateral tubercles and weak ribs, and involute coiling (KENNEDY & COBBAN, 1991: 26, Pl. 2, Figs. 1, 15–17; Pl. 5, Figs. 1–11; Text-Figs. 4c, 6, 7). *Forresteria (Forresteria) brancoi* (SOLGER, 1904: 174, Pl. 5, Figs. 1, 2, 4, 5; Text-Figs. 63–66, 68) differs by its reduction of ribbing and loss of umbilical tuberculation leading to smooth flanks but retains a few but very coarse ventrolateral tubercles (KENNEDY & COBBAN, 1991: 30, Text-Figs. 8–11).

Stratigraphy

Forresteria (Harleites) seidli nov. sp. is present in the basal Santonian of the Streiteck Formation (SUMMESBERGER et al., 2017) of the Randograben (Rußbach, Salzburg), co-occurring with *Texanites quinquenodosus* (REDTENBACHER, 1873), *Eulophoceras natalense* HYATT, 1903, *Texasia cricki* (SPATH, 1921), and *Cladoceras undulatoplicatus* (ROEMER, 1852), which confirm the Early Santonian age of the species. No biostratigraphically distinct calcareous nannofossils were found in samples from the site. *Muniericeras gosauicum* (HAUER, 1858) is absent in the Streiteck Formation (Rando 3; SUMMESBERGER et al., 2017: Text-Fig. 3) but it is very common in the sites Rando 1 and Rando 2 (Text-Fig. 1). *Forresteria (Harleites) seidli* nov. sp. appears to be so far the youngest reported representative of the genus.

Conclusion

A new species, *Forresteria (Harleites) seidli* nov. sp. is described from the Cretaceous of the Austrian Gosau Group. The most surprising observation is its appearance in the Santonian in contrast to the widespread occurrence of the genus in the upper Turonian to Coniacian.

Acknowledgements

WOLFGANG SCHWAIGHOFER is gratefully acknowledged for his help in the field and for the loan of material from his collection. ALICE SCHUMACHER took the photos and ILKA WÜNSCHE created the plates.

References

- BASSE, E. (1947): Paléontologie de Madagascar – 26. Les peuplements Malgaches de Barroisiceras (Révision du genre Barroisiceras DE GROSS.). – Annales de Paléontologie, **33**, 99–190, Paris.
- BOULE, M., LEMOINE, P. & THÉVENIN, A. (1906–1907): Paléontologie de Madagascar – III. Céphalopodes crétacés des environs de Diego-Suarez. – Annales de Paléontologie, **1**, 173–192 (1–20); **2**, 1–56 (21–76), Paris.
- BRÜGGEN, H. (1910): Die Fauna des unteren Senons von Nord-Peru. Beiträge zur Geologie und Paläontologie von Südamerika. – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beilageband **30**, 717–788, Stuttgart.
- COQUAND, H. (1859): Synopsis des animaux et des végétaux fossils observés dans la formation crétacée du Sud-Ouest de la France. – Bulletin de la Société Géologique de France, **16**, 945–1023, Paris.
- GROSSOUVRE, A. DE (1894): Recherches sur la craie supérieure, 2, Paléontologie. Les ammonites de la craie supérieure. – Mémoires du Service de la Carte Géologique détaillée de la France, 264 pp. (misdated 1893), Paris.
- HAUER, F. VON (1858): Über die Cephalopoden der Gosauschichten. – Beiträge zur Palaeontographie von Oesterreich, **1**, 7–14, Wien.
- HYATT, A. (1889): Genesis of the Arietitidae. – Smithsonian Contributions to Knowledge, **673**, 238 pp., Washington, D.C.
- HYATT, A. (1903): Pseudoceratites of the Cretaceous. – Monographs of the United States Geological Survey, **44**, 351 pp., Washington, D.C.
- IFRIM, C., MÚZQUIZ, H.P. & STINNESBECK, W. (2019): Ammonooids, their biozonation and their palaeobiogeographic relation across the Turonian-Coniacian boundary in northern Coahuila, Mexico. – Cretaceous Research, **102**, 170–195, London.
- KENNEDY, W.J. (1984): Systematic Palaeontology and Stratigraphic Distribution of the Ammonite Faunas of the French Coniacian. – Special Papers in Palaeontology, **31**, 160 pp., London.
- KENNEDY, W.J. (1985): *Solgerites* REESIDE, 1932 (Cretaceous Ammonoidea) a synonym of *Forresteria* REESIDE, 1932, with a revision of *Solgerites brancoi* (SOLGER, 1904) (Cretaceous Ammonoidea) from Cameroon. – Paläontologische Zeitschrift, **59**, 211–222, Stuttgart.
- KENNEDY, W.J. & COBBAN, W. (1991): Coniacian ammonite faunas from the United States Western Interior. – Special Papers in Palaeontology, **45**, 96 pp., London.
- KENNEDY, W.J. & KLINGER, H.C. (1975): Cretaceous faunas from Zululand and Natal, South Africa. Introduction, Stratigraphy. – Bulletin of the British Museum (Natural History): Geology Series, **25**, 263–315, London.
- KENNEDY, W.J. & WALASZCZYK, I. (2004): *Forresteria (Harleites) petrocoriensis* (COQUAND, 1859), from the Upper Turonian *Mytiloides scupini* Zone of Slupia Nadbrzenna, Poland. – Acta Geologica Polonica, **54**, 55–59, Warszawa.
- KENNEDY, W.J., WRIGHT, C.W. & KLINGER, H.C. (1983): Cretaceous Faunas from Zululand and Natal, South Africa. The Ammonite Subfamily Barroisiceratinae BASSE, 1947. – Annals of the South African Museum, **90**, 241–324, Cape Town.
- REDTENBACHER, A. (1873): Die Cephalopodenfauna der Gosauschichten in den nordöstlichen Alpen. – Abhandlungen der k. k. Geologischen Reichsanstalt, **5**, 91–140, Wien.
- REESIDE, J.B. (1932): The Upper Cretaceous ammonite Genus *Barroisiceras* in the United States. – United States Geological Survey Professional Paper, **170-B**, 7–18, Washington, D.C.
- ROEMER, F.A. (1852): Die Kreidebildungen von Texas und ihre organischen Einschlüsse. – 100 pp., Bonn (Adolph Marcus).
- SOLGER, F. (1904): Die Fossilien der Mungokreide in Kamerun und ihre geologische Bedeutung, mit besonderer Berücksichtigung der Ammoniten. – In: ESCH, E., SOLGER, F., OPPENHEIM, P. & JAEKEL, O.: Beiträge zur Geologie von Kamerun, **2**, 85–242, Stuttgart.
- SPATH, L.F. (1921): On Upper Cretaceous Ammonoidea from Pondoland. – Annals of the Durban Museum, **3**, 39–56, Durban.
- SUMMESBERGER, H., KENNEDY, W.J., WAGREICH, M. & SKOUMAL, P. (2017): Early and middle Santonian Cephalopods from the Gosau Group (Upper Cretaceous, Austria). – Abhandlungen der Geologischen Bundesanstalt, **71**, 5–100, Wien.
- SUMMESBERGER, H., KENNEDY, W.J., WAGREICH, M., SKOUMAL, P. & MAHERNDL, W.P. (2022): Coniacian Cephalopods from the Gosau Group (Upper Cretaceous, Salzkammergut, Austria). – Abhandlungen der Geologischen Bundesanstalt, **75**, 15–109, Wien.
- WAGREICH, M. (1988): Sedimentologie und Beckenentwicklung des tieferen Abschnitts (Santon – Untercampan) der Gosauschichtgruppe von Gosau und Rußbach (Oberösterreich – Salzburg). – Jahrbuch der Geologischen Bundesanstalt, **131**, 663–685, Wien.
- WRIGHT, C.W. (1957): Cretaceous ammonites. – In: MOORE, R.C. (Ed.): Treatise on Invertebrate Paleontology, Part L, Mollusca 4, Cephalopoda Ammonoidea, 490 pp., New York–Lawrence (The Geological Society of America & The University of Kansas).
- WRIGHT, C.W. & WRIGHT, E.V. (1951): A survey of the fossil Cephalopoda of the Chalk of Great Britain. – Monograph of the Palaeontographical Society, 1–40, London.
- ZITTEL, K.A. V. (1884): Handbuch der Paläontologie, **1**, Abteilung 2: Cephalopoda, 329–522, München–Leipzig.

Plates

Plate 1

Figs. 1, 2: *Forresteria (Harleites) seidli* nov. sp.; SCH/RA/2019/02.

Figs. 3, 4: *Forresteria (Harleites) seidli* nov. sp.; CE.00027, the holotype.

All figures are natural size, coated with ammonium chloride, and from Randobach (Rando 3), Rußbach (Salzburg), Lower Santonian.




Plate 2

Figs. 1, 2: *Forresteria (Harleites) seidli* nov. sp.; SCH/RA/2019/01.

Figs. 3, 4: *Forresteria (Harleites) seidli* nov. sp.; LEI/GO/2019/0017.

All figures are natural size, coated with ammonium chloride, and from Randobach, Rußbach (Salzburg), Lower Santonian.

