

## SOME NEW SILURIAN NAUTILOID CEPHALOPODS FROM THE ITALIAN SIDE OF THE CARNIC ALPS

Serventi Paolo

Dipartimento di Scienze della Terra (Modena University, Italy). Email: gnolim@unimo.it

A preliminary investigation carried out on scattered blocks of "Orthoceras" Limestone from the proximity of the old mines of Mt. Cocco (Ugovizza, Udine, Italian side of the Carnic Alps) allows the taxonomic identification at the generic and specific level of many nautiloid specimens among which I can report: *Hemicosmorthoceras laterculum* Ristedt, 1968, *H. semiannulatum* (Barrande, 1870), *H.?* cf. *semilaeve* (Barrande, 1868), *Sphaerorthoceras carnicum* Ristedt, 1968, *S. curvum* Ristedt, 1968, *S.* cf. *teichertii* Ristedt, 1968, *Parasphaerorthoceras* sp. H. *sensu* Ristedt. 1968, *P.* sp. E *sensu* Ristedt. 1968, *P.* sp. C *sensu* Ristedt. 1968, *Kopaninoceras thyrsus* (Barrande, 1870), *Leurocycloceras subannulare* (Münster, 1840).

Beyond the above reported taxa I can list other taxa of nautiloid cephalopod closely resembling those already studied by Gnoli & Histon (1998) such as *Arionoceras submoniliforme* (Meneghini, 1857) with protoconch, *Kopaninoceras* cf. *jucundum* (Barrande, 1870), some specimens probably belonging to genera *Kionoceras* Hyatt, 1884 and *Parakionoceras* (Foerste, 1928) and *Plagiostomoceras?* cf. *gruenewaldti* (Barrande, 1866)

All the reported taxa are found for the first time in the Italian side of the Carnic Alps. The reported taxa indicate without doubt a close faunal relationship between Soutwestern Sardinia and Central Bohemia.

Further collections will be made from other localities in the Italian Carnic Alps including "Casera Pal Grande", "Cima Costa Alta", "Mt. Zermula".

### References

- BARRANDE, J., 1866, 1868, 1870. Système Silurien du centre de la Bohême. Recherches Paléontologiques, v. 2, Classe des Mollusques, Ordre des Céphalopodes, Prague, Paris.
- FOERSTE, A. F., 1928. A restudy of American orthoconic Silurian cephalopods. Denison University Bulletin Journal Science Laboratory, v. 23, pp. 236-320.
- GNOLI, M. & HISTON, K., 1998. Silurian Nautiloid cephalopods from the Carnic Alps: a preliminary investigation: Bollettino della Società Paleontologica Italiana, 36(3): 311-330, 4 pls., 4 text-figs., 3 tabs.
- HYATT, A., 1884. Genera of fossil cephalopods. Proceeding of Boston Society Natural History, v. 22, pp. 273-338, Boston.
- MENEGHINI, G., 1857. Paléontologie de l'île de Sardaigne. In: La Marmora A., Voiage en Sardaigne, p. 53-144, pl. C, Imprimerie Royale, Turin.
- MÜNSTER, G. G., 1840. Die Versteinerungen des Uebergangskalkes mit Clymenien und Orthoceratiten: Beiträge zur Petrefacten-Kunde, 3: 33-121, Bayreuth.
- RISTEDT, H., 1968. Zur Revision der Orthoceratidæ Abh. Der Math.-Natur. wissenschaft. Klasse, No. 4, pp. 212-287, 5 pls., 4 text-figs.