

HOW MANY TETHYAN TRIASSIC OCEANS?

Leopold KRYSTYN ¹, Richard LEIN ² & Hans-Jürgen GAWLICK ³

¹ Department of Palaeontology, University of Vienna, Althanstrasse 14, A-1090 Vienna.

² Department of Geodynamics & Sedimentology, University of Vienna, Althanstraße 14, A-1090 Vienna.

³ Department for Applied Geosciences & Geophysics, Univ. of Leoben, Peter-Tunner-Str.e 5/I, A-8700 Leoben.
leopold.krystyn@univie.ac.at richard.lein@univie.ac.at, hans-juergen.gawlick@mu-leoben.at

In recent years the number of independent oceanic basins within the Western Tethys has considerably increased. From an originally single Western Tethys ocean (LAUBSCHER 1971, BERNOULLI & LAUBSCHER 1972), especially STAMPFLI and collaborators (STAMPFLI et al. 1991, 1999, STAMPFLI & KOZUR 2006, STAMPFLI in MOIX et al. 2008) have introduced a system of at least four, more or less parallel E-W directed oceanic basins with intermediate small ribbon-continental blocks. These from Ladinian time onward existing oceans are from north to south the Meliata-, Maliac-, Pindos- and Neotethys oceans and are described to represent highly individual Triassic histories in space and time.

Starting from critical key areas for this concept we discuss its strength as well weakness and its reliability. Based on facial and tectonic considerations we see no reasons for this multiple splitting of the oceanic Western Tethys end and present arguments for combining at least the Meliata- and the Maliac ocean as well as the Pindos- and the Neotethys ocean into single oceans. Following the concept of SCHMID et al. (2008) and GAWLICK et al. (2008) for a far distance westward transport of the Pindos ophiolites in combination with their later complicated deformational history in their present place, (VAMVAKA et al. 2006), all western Tethys remnants of oceanic crust would fit in a single ocean paleogeography as classically supposed.

References

- BERNOULLI, D. & LAUBSCHER, H. (1972): The palinspastic problem of the Hellenides. - *Ecl. Geol. Helv.*, **65**: 107-118.
- GAWLICK, H.-J., FRISCH, W., HOXHA, L., DUMITRICA, P., KRYSTYN, L., LEIN, R., MISSONI, S. & SCHLAGINTWEIT, F. (2008): Mirdita Zone ophiolites and associated sediments in Albania reveal Neotethys Ocean origin. - *Int. Journ. Earth. Sci.* **97**: 865-881, (Springer) Berlin. DOI 10.1007/s00531-007-0193-z
- LAUBSCHER, H. P. (1971): Das Alpen-Dinariden-Problem und die Palin-spastik der südlichen Tethys. - *Geol. Rundschau*, **60**: 813-833.
- MOIX, P., BECCALETTO, L., KOZUR, H.W., HOCHARD, C., ROSSELET, F. & STAMPFLI, G. (2008): A new classification of the Turkish terranes and sutures and its implication for the paleotectonic history of this region. - *Tectonophys.*, **451**: 7-39.
- SCHMID, S.M., BERNOULLI, D., FÜGENSCHUH, B., MATENCO, L., SCHEFER, S., SCHUSTER, R., TISCHLER, M. & USTASZEWSKI, K. (2008): The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units. - *Swiss J. Geosci.*, **101**: 139-183, Basel.
- STAMPFLI, G. & KOZUR, H.W. (2006): Europe from Varican to the Alpine cycles. - In: GEE, D.G. & STEPHENSON, R.A. (Eds.): *Europaean lithosphere dynamics*. – Mem. Geological Society, 57-82, London.
- STAMPFLI, G., MARCOUX, J. & BAUD, A. (1991): Tethyan margins in space and time. - *PPP*, **87**: 373-409.
- STAMPFLI, G. M. & MOSAR, J. (1999): The making and becoming of Apulia. - *Mem. Sci. Geol.*, **51/1**: 141-154.
- VAMVAKA, A. et al. (2006): Geometry and structural evolution of the Mesohellenic Trough (Greece): a new approach. - *Geol. Soc. Spec. Publ.*, **260**: 521-538.