

Inoceramidae from the Campanian (Upper Cretaceous) of the Gschliefgraben (Ultrahelvetic; Austria)

by

Karl-Armin TRÖGER¹, Herbert SUMMESBERGER² & PETER Skoumal³

TRÖGER, K.-A., SUMMESBERGER, H. & SKOUMAL, P., 1999. Inoceramidae from the Campanian (Upper Cretaceous) of the Gschliefgraben (Ultrahelvetic; Austria). — Beitr. Paläont., 24: 41–61, 16 Text-figures, 15 Tables, 4 Plates, Wien.

Contents

Abstract, Zusammenfassung	41
Introduction	41
Systematic Palaeontology	42
Conclusion	50
References	52

Abstract

Twelve taxa of Inoceramidae are described from the Late Campanian through Early Maastrichtian of the Gschliefgraben (Ultrahelvetic, Upper Austria): *Cataceramus balticus balticus* (BÖHM), *Cataceramus balticus ellipticus* (GIERS), *Cataceramus balticus* aff. *haldemensis* (GIERS), *Cataceramus balticus* (BÖHM) subsp. indet., *Endocostea impressa* (D'ORBIGNY), *Inoceramus regularis* D'ORBIGNY, *Inoceramus cf. boriliensis* JOLKIČEV, *Inoceramus* aff. *bererensis* SORNAY, *Inoceramus sagensis* OWEN, *Cordiceramus* ? aff. *heberti* (FALLOT), *Cremneceramus* aff. *inconstans* Woods und *Cremneceramus sarumensis* (Woods sensu SORNAY 1982). The material is from a permanently active mudflow area. The sequence was compiled by KENNEDY & SUMMESBERGER (1984) after PREY (1983). The diverse fauna indicates a Late Campanian through (?) Early Maastrichtian age.

Zusammenfassung

Zwölf Taxa Inoceramidae werden aus dem Obercampan bis (?) tiefsten Untermaastricht des Gschliefgrabens (Ultrahelveticum; Oberösterreich) beschrieben: *Cataceramus balticus balticus* (BÖHM), *Cataceramus balticus ellipticus* (GIERS), *Cataceramus balticus* aff. *haldemensis* (GIERS), *Cataceramus balticus* (BÖHM)

subsp. indet., *Endocostea impressa* (D'ORBIGNY), *Inoceramus regularis* D'ORBIGNY, *Inoceramus cf. boriliensis* JOLKIČEV, *Inoceramus* aff. *bererensis* SORNAY, *Inoceramus sagensis* OWEN, *Cordiceramus* ? aff. *heberti* (FALLOT), *Cremneceramus* aff. *inconstans* Woods und *Cremneceramus sarumensis* (Woods sensu SORNAY 1982). Das Material stammt aus einem dauernd aktiven Murenstrom. Die kompilierte lithostratigraphische Abfolge (KENNEDY & SUMMESBERGER, 1984) beruht daher vor allem auf der Interpretation des Fossilinhaltedes (PREY 1983).




Figure 1: Sketchmap of the Gschliefgraben area.

Introduction

The only inoceramid taxon mentioned in the previous literature (PREY 1983) from the Gschliefgraben is *Inoceramus salisburgensis*. We cannot confirm this identification. *I. salisburgensis* FUGGER & KASTNER is not present in PREY's field collection, which is stored in the collections of the Museum of Natural History in Vienna. A second collection of Inoceramidae was donated (1976) to the NHMW by Prof. Norbert VAVRA (PIUW). The bulk of the fauna was collected by the authors. The description given below is a contribution

¹ Bergakademie Freiberg, Geologisches Institut, Zeunerstr. 12 – Meißen-Bau, D-09596 Freiberg, Germany
Naturhistorisches Museum Wien, Burgring 7,
A -1014 Wien, Austria
e-mail: herbert.summesberger@nhm-wien.ac.at
Bastiengasse 56, A-1180 Wien, Austria

to an integrated documentation of the fossil content of the Late Campanian of the Gschließgraben. Most of the contributions are presented in this volume.

Abbreviations

NHMW	Naturhistorisches Museum Wien
GIBF	Geologisches Institut Bergakademie Freiberg (Technische Universität)/Sachsen
ISNB	Institut Royal des Sciences Naturelles de Belgique
MNHNP	Muséum National d'Histoire Naturelle, Paris
SK	Skoumal private collection, Vienna
GSLT	Geological Survey of Libya, Tripoli
USNM	United States National Museum Washington
PIUW	Paläontologisches Institut Universität Wien

Systematic Palaeontology

Class Bivalvia

Supraorder Pteriomorphia BEURLEN 1944
Order Pteroidea NEWELL 1965
Family Inoceramidae GIEBEL 1852

Genus *Cataceramus* HEINZ 1932

Type species *Inoceramus goldfussianus* D'ORBIGNY (= *Inoceramus balticus* BÖHM, 1907) by original designation of COX, 1969: N 315.

Cataceramus balticus balticus (BÖHM) (Pl. 4, Figs. 1, 2; Text-fig. 2)

Synonymy:

- 1907 *Inoceramus balticus* J. BÖHM: 113.
- 1909 *Inoceramus balticus* J. BÖHM; J. BÖHM: 47; pl. 11, fig. 1.
- 1964 *Inoceramus balticus balticus* J. BÖHM; GIERS: 238–239; pl. 1, figs. 2–4.
- 1967 *Inoceramus (Endocostea) balticus* J. BÖHM; SEITZ: 67–72; pl. 6, figs. 1, 2; pl. 8, figs. 1, 2; pl. 10, fig. 1; pl. 12, figs. 1, 2.
- 1993 *Endocostea baltica baltica* (J. BÖHM, 1907); DHONDTE: 221, pl. 4, fig. 3.
- 1997 *Cataceramus balticus* (BÖHM, 1907); WALASZCZYK: 18; pl. 12, figs. 1–5 (with additional synonymy).

Material: Internal moulds of two right valves (NHMW 1979/2076/46, NHMW 1979/2076/37) and one left valve (GIBF 1994/4a).

Preservation Incomplete. Portions of the wings, the anterior auricles and the ventral margins are missing. NHMW 1979/2076/46 is completely flattened by compaction.

Description: Valve medium-sized, elongated subquadrate. Thick beak projecting to the hinge line. The umbonal pole is slightly turned to the anterior margin. Anterior margin at the beak concave. Geniculations with a change in the distances between the undulations (ribs) were observed at $H = 10.5$ mm (NHMW 1979/2076/37) and $H = 75.5$ mm (NHMW 1979/2076/46). Total angles: 93–105° (GIERS 1964, 238: 98°). The undulations are toprounded to sharp due to compac-

tion. The H/L- and Vo/L-ratios are similar to those of the holotype and specimen IG 6312 (MNHNP). Geographic distribution: Widespread; E-, W- and S-Europe, Asia, N- and S-America (Brasil).




Figure 2: Diagram showing H/L and Vo/L ratios of *Cataceramus balticus balticus* (J. BÖHM).

Distance from UP	1	2	3
10–30 mm	—	—	—
30–50 mm	4.73 mm	6.0 mm	5.5 mm
50–70 mm	—	1.1 mm	—

Table 1. Average undulation intervals (DU) in different distances from the umbilical pole (UP); 1 = NHMW 1979/2076/37, 2 = NHMW 1979/2076/46, 3 = IG 6312 (MNHNP).

Stratigraphic distribution: After GIERS (1964, table 1) *Cataceramus balticus balticus* is a long ranging subspecies appearing in the uppermost Santonian and ranging through the Early Campanian, disappearing in the Late Campanian *Polypliocum* Zone. According to TRÖGER (1989; Text-fig. 5) it occurs in Inoceramid-zones 29–32.

Cataceramus balticus ellipticus (GIERS) (Text-figs. 3, 4)

- 1964 *Inoceramus balticus ellipticus* GIERS: 244–245; pl. 2, figs. 3, 4
- 1993 *Endocostea baltica elliptica* (GIERS, 1964); DHONDTE: 223, pl. 6, fig. 1; text-fig. 6
- 1997 *Inoceramus ellipticus* GIERS, 1964; WALASZCZYK: 39; pl. 23, figs. 1–7; pl. 24, figs. 1–5; pl. 25, figs. 1, 4.

Material: Two internal moulds of right valves without shell (NHMW 1979/2076/34, NHMW 1979/z/160/5); additional fragments from the Gschließgraben.

P r e s e r v a t i o n : Incomplete. Portions of the wings (posterior auricle) and the ventral margins are absent. Flattened by compaction, with radial cracks.

D e s c r i p t i o n Valve small to medium-sized, shape elongated oval. Small beak slightly turned to the anterior margin and separated from the wing. Anterior auricles are missing. Total angles: 102°, 105°. Angle of





Figure 3:
Cataceramus balticus ellipticus (GIERS);
NHMW 1979/2076/34; Gschliefgraben
near Gmunden; x 1.



H/L , Vo/L and S/L at *Cataceramus balticus ellipticus* (GIERS)

Figure 4: Diagram showing H/L-, Vo/L- and S/L- ratios at *Cataceramus balticus ellipticus* (GIERS).

inclination (growth axis): 25–30°. Geniculations were observed at $H=21$ mm (NHMW 1979/2076/34) and $H=20.2$ mm (NHMW 1997/z/160/5). At the geniculations there is a sudden change in the intervals of the undulations (ribs). Distance between the undulations on the umbonal region (10–30 mm from UP): 2–4 mm.

D i s c u s s i o n WALASZCZYK (1997:40) quoted DHONDT's *Endocostea baltica elliptica* (GIERS) with some doubt concerning the ornament.

Geographic distribution: W- Europe, Münsterland (GIERS 1964), France (HANCOCK et al. 1993).

Stratigraphic distribution: According to GIERS (1964: 244) *Cataceramus balticus ellipticus* (GIERS) occurs in

	NHMW 1979/2076/34		NHMW 1997/z/160/5		
Distance from UP	H/L	Vo/L	S/L	H/L	Vo/L
10–30 mm	54.6	39.6	48.0	68.0	50.0

Table 2. Average H/L-, Vo/L - and S/L - ratios of *Cataceramus balticus ellipticus* (GIERS) from the Gschliefgraben.

the Vorhelmer Schichten, representing the upper part of the early Late Campanian. After WALASZCZYK (1997) it appears in the zone of *Cataceramus balticus beckumensis* (GIERS), which is earliest Late Campanian in age.

***Cataceramus balticus* aff. *haldemensis* (GIERS)**
(Pl. 3, Fig. 4; Text-fig. 5)

C o m p a r e :

- * 1932 *Cataceramus haldemensis* HEINZ (nomen nudum): 15
- * 1964 *Inoceramus balticus haldemensis* GIERS: 243–244; pl. 2, fig. 2
- 1967 *Inoceramus (Endocostea) balticus haldemensis* GIERS; SEITZ: 75; pl. 2, fig. 2.
- * 1982 *Inoceramus balticus haldemensis* GIERS; TRÖGER & RÖHLICH: 101–111; 9 figs., 1 pl.
- * 1992 *Inoceramus balticus haldemensis* GIERS; TRÖGER & RÖHLICH: 1363–1365; text-figs. 4–7; pl. 1, figs. 1–8.
- 1997 *Inoceramus haldemensis* GIERS; WALASZCZYK: p. 40; pl. 30, figs. 1, ?3, 5.

M a t e r i a l : Five internal moulds: 2 right valves (GIBF 1995/1, NHMW 1997/160/4), 2 left valves (GIBF 1995/2, GIBF 1995/6), 1 bivalved (SK/B/GG/1997/4).


D e s c r i p t i o n Medium-sized, equivalve, inequilateral. Shape axe-like (SK/B/GG/1997/4, GIBF 1995/1). The other specimens are higher (H/L -ratio > 100 , see text-fig 5). Beak in terminal position. Umbonal region slightly prosogyrate and thrusted over the hinge line. Anterior auricle is missing. Anterior margin convex. Undulations (concentric ribs) toprounded. The undulations are turned to the beak on the wing. Undulations sometimes bifurcated. Geniculations were observed in different growth stages. Total angles between 110–150°.

R e m a r k s The H/L - and Vo/L - ratios (text-fig. 3) are comparable with those of a specimen from the Late Campanian of Libya (GSLT/1385-III-C-5/5).

Distance from UP	1	2	3	4
10–30 mm	3.48 mm	3.12 mm	–	3.0 mm
30–50 mm	5.00 mm	10.00 mm	–	3.53 mm
50–70 mm	11.00 mm	–	–	–

Table 3. Average undulation intervals (DU) in different distances from the umbonal pole (UP). 1 GIBF 1995/4, 2 SK/B/GG/1997/4, 3 GIBF 1995/1, 4 GIBF 1995/6

Geographical distribution: W- and E-Europe, N-Africa.



Variability of H / L and Vo / L at *Cataceramus balticus* aff. *haldemensis* (GIERS) from the Gschlieffgraben

Figure 5: Diagram showing H/L and VO/L -ratios at *Cataceramus balticus* aff. *haldemensis* (GIERS).


Stratigraphic distribution: After GIERS (1994: 243) *I. balticus haldemensis* appears in the „Obere Vorhelmer Schichten“, representing the *Vari* Zone of the Late Campanian, and extends into the *Polyplocum* Zone. After WALASZCZYK (1997) it occurs in the Late Campanian *C. haldemensis* Zone, which is more or less identical with the *Polyplocum* Zone of the traditional Campanian zonation.

***Cataceramus balticus* (BÖHM) subsp. indet.
(Pl. 3, Fig. 2; Pl. 4, fig. 3; Text-fig. 6)**

M a t e r i a l : Internal moulds of four right valves (NHMW 1979/2076/24, SK/B/GG/ 1997/1, GIBF 1995/7, NHMW 1979/2077/5), two left valves (SK/B/GG/1997/2, NHMW 1979/2077/5) and one bivalved specimen (NHMW 1997/2076/28) from the Gschlieffgraben.

P r e s e r v a t i o n : The bivalved specimen is completely flattened by compaction. The other specimens are slightly deformed, showing radial cracks.

D e s c r i p t i o n : Shell medium-sized, equivalve, inequilateral. Shape elongated oval. Beak distinctly separated from the wing, erected. Umbonal pole slightly prosogyrate. Anterior margin convex. Total angle: 120–125°. Undulations (concentric ribs) staircase shaped to toprounded. Growth lines and undulations are bent to the umbonal pole on the wing. Anterior auricle is



Variability of H / L and Vo / L at *Cataceramus balticus* (BÖHM) subsp. indet. from the Gschlieffgraben

Figure 6: Diagram showing H/L- and Vo/L- ratios of *Cataceramus balticus* (J. BÖHM) al from the Gschlieffgraben.

Distance from UP	1	2	3
10–30 mm	–	2.53 mm	6.00 mm
30–50 mm	9.25 mm	–	–
50–70 mm	11.40 mm	–	–

Table 4. Average undulation intervals (DU) of *C. balticus* subsp. indet. at different distances from the umbonal pole (UP) from Gschlieffgraben specimens: 1 = NHMW 1979/2076/24, 2 = NHMW 1979/2076/28, 3 = GIBF 1995/7.

Distance from UP	1	2	3
10–30 mm	4.87 mm	2.76 mm	–
30–50 mm	5.00 mm	5.70 mm	5.10 mm
50–70 mm	–	–	–

Table 5. Average undulation intervals (DU) of *C. balticus* subsp. indet. at different distances from the umbonal pole (UP) from Wentneralm I locality in Gams (SUMMERSBERGER et al. 1999): 1 = NHMW 1997/z /158/1, 2 = NHMW 1997/z/158/3, 3 = NHMW 1997/z/158/8.

missing. Geniculations were observed in: GIBF 1995/7, SK/B/GG/1997/2 and NHMW 1979/2076/24. NHMW 1979/2076/28 agrees with the specimens from Gams concerning the undulation intervals. The intervals are greater in the other Gschlieffgraben specimens. **R e m a r k s** Concerning the shape and the H/L- ratios there are distinct differences between *C. balticus* subsp. indet. and *C. balticus balticus* (J. BÖHM) including *C. balticus haldemensis* (GIERS) (Text-figs. 2, 5).

The shape, development of the umbonal region and the total angle of *C. balticus marcki* (GIERS) are comparable. In contrast to *Cataceramus balticus* (J. BÖHM) subsp. indet. in *Cataceramus balticus marcki* (GIERS) the H/L-ratio strongly increases (170–180 %).

Stratigraphic distribution: *Cataceramus balticus* ssp. is a long-ranging species:

Cataceramus balticus marcki (GIERS) occurs from latest Santonian to the earliest Late Campanian *Polyplacum* Zone.

Genus *Endocostea* WHITFIELD 1880

Type species *Inoceramus (Endocostea) typicus* WHITFIELD.

Endocostea impressa (D'ORBIGNY)

(Pl. 2, Figs. 1, 3, 5, 6; Text-figs. 7 a, b; 8)

1845 *Inoceramus impressus* D'ORBIGNY: 515; pl. 409.

* 1957 *Inoceramus impressus* D'ORBIGNY; SORNAY: figs. 4, 5.


According to SORNAY (1957) the lectotype of *Endocostea impressa* (D'ORBIGNY) is no. 7592 A in d'ORBIGNY's collection (SORNAY, 1957, fig. 4).

M a t e r i a l Three specimens from the Gschliefgraben are available to us: One bivalved (NHMW 1979/2076/25/1,2), one right (SK/B/GG/1997/3) and one left valve (NHMW 1979/2076/30). Two internal moulds of right valves (NHMW 1997z158/0009, 0011) from




Figure 7a: *Endocostea impressa* (d'ORBIGNY) not deformed umbonal region;

NHMW 1979/2076/25/2; x 1.



7b: *Endocostea impressa* (d'ORBIGNY) with radial cracks caused by compaction; SK/B/GG/1997/3, both from Gschliefgraben; x 1.



Variability of H / L , Vo / L and S / L at *Endocostea impressa* (D'ORBIGNY) from the Gschliefgraben (no. 3-5) .

Figure 8: Diagram showing H/L- and VO/L- ratios at *Endocostea impressa* (d'ORBIGNY) from the Gschliefgraben and of *Endocostea aff. impressa* (d'ORBIGNY) from Wentnernalm, Gams/Steiermark.

the Wentnernalm I at Gams (SUMMESBERGER, WAGREICH, TRÖGER & JAGT 1999) are ancestral to the present species.

Preservation: All internal moulds are incomplete. Parts of the wings and the ventral margins are missing.

Distance from UP	1	2	3	4
10–30 mm	3.73 mm	3.16 mm	3.26 mm	3.60 mm
30–50 mm	4.55 mm	—	5.87 mm	—
50–70 mm	3.61 mm	—	5.20 mm	—

Table 6. Average undulation intervals (DU) of *E. impressa* from Gschliefgraben at different distances from UP. 1 = NHMW 1979/2076/25, 2 = NHMW 1979/2076/30, 3 = SK/B/GG/1997/3, 4 = 7592A (MNHN, collection d'ORBIGNY).

Description: Medium-sized, inequilateral. Shape elongated subquadrate. Beak vaulted, distinctly separated from the wing, prosogyrate. Hinge line straight. Anterior margin slightly concave at the umbonal region. Undulations (concentric ribs) following the shape outline, toprounded. The undulations are bent to the umbonal pole (angle 115–120°).

The course of the H/L-, Vo/L- and S/L-ratios (Text-fig. 8) of specimens from the Gschliefgraben and Gams are similar, but they do not agree completely. No. 1966 from the collection of D'ORBIGNY (lectotype) has an intermediate position. Total angle: 92–110° (lectotype: 102°).

R e m a r k s : The specimens from the Gschliefgraben and Wentneralm I (Gams) differ slightly in H/L- and Vo/L-ratios (see Text-fig. 8). The specimens from the Gschliefgraben agree with no. 1966 in D'ORBIGNY'S collection of Early Maastrichtian age.

Geographic distribution: W- and E- Europe, N-Africa.

Stratigraphic distribution: Early Late Campanian (GIERS 1964) to earliest Maastrichtian (SORNAY 1957). According to WALASZCZYK (1997) Late Campanian through Early Maastrichtian, (?) Late Maastrichtian.

Genus *Inoceramus* J. SOWERBY 1814

T y p e s p e c i e s *Inoceramus cuvieri* SOWERBY by subsequent designation of Cox (1969: N 315).

Inoceramus regularis D'ORBIGNY

(Pl. 2, Fig. 2; Text-fig. 9)

- 1845 *Inoceramus regularis* D'ORBIGNY: 517, pl. 410
- * 1962 *Inoceramus regularis* D'ORBIGNY; SORNAY: pl. 7, fig. 3
- * 1964 *Inoceramus regularis* D'ORBIGNY; GIERS: 247–248; pl. 3, fig. 3
- * 1976 *Inoceramus regularis* D'ORBIGNY; SORNAY: 7–8; pl. 2, fig. 3; pl. 3, figs. 3–4; text-fig. 4.

Following HEINZ (in sched.) SORNAY (1976: 7–8) designated no. 7594 in D'ORBIGNY's collection as lectotype (SORNAY 1962: pl. 1, fig. 3).

M a t e r i a l : A single internal mould of a right valve (NHMW 1979/2076/39) from the Gschliefgraben.


Preservation: Incomplete. Parts of the wing and the anterior margin are missing.

D e s c r i p t i o n : Medium-sized, inequilateral, shape nearly cycloid. Flat with convex anterior margin. Prosogyrate. The umbonal pole projects somewhat above the straight hinge line. Undulations (concentric ribs) toprounded. Average distance between the undulations at 10–30 mm distance from the umbonal pole: 3.5 mm. Total angle: 120°. Text-fig. 9 shows H/L-, S/L- and Vo/L-ratios.

R e m a r k s : The present specimen corresponds on all features with the lectotype of *Inoceramus regularis* D'ORBIGNY.

Geographic distribution: Europe, Madagascar (NODA & KANIE 1978 a,b), N-America (KAUFFMAN et al. 1993), N-Africa (TRÖGER & RÖHLICH 1992).

Stratigraphic distribution: Late Senonian after SORNAY (1962); Late Campanian after NODA & KANIE (1978 a, b), early Late Campanian after GIERS (1964).



Inoceramus regularis D'ORBIGNY from the Gschliefgraben.

Figure 9: Diagram showing H/L-, Vo/L- and S/L-ratios at *Inoceramus regularis* D'ORBIGNY from the Gschliefgraben.

Distances from UP

NHMW 1979/2077/3

10–30 mm	—
30–50 mm	3.5 mm
50–70 mm	5.0 mm
over 70 mm	11.5 mm

Table 7. Average undulation intervals of *Inoceramus* cf. *borealis* JOLKIČEV (NHMW 1979/2077/3) from the Gschliefgraben.

Distance from UP	H/L	Vo/L
10–30 mm	44.9 mm	32.7 mm
30–50 mm	66.3 mm	46.75 mm

Table 8. Average H/L- and Vo/L- ratios of *Inoceramus* cf. *borealis* JOLKIČEV (NHMW 1979/2077/3) from the Gschliefgraben.

Inoceramus cf. *borealis* JOLKIČEV (Text-figs. 10, 11)

C o m p a r e :

- 1962 *Inoceramus borealis* JOLKIČEV: 1952–1954; pl. 7, figs. 1a, 1b
- 1993 „*Inoceramus*“ *borealis* JOLKIČEV; DHONDT: 215; pl. 1, fig. 1; pl. 3, fig. 2a, b; text-fig. 3a, b
- 1993 „*Inoceramus*“ cf. *borealis* JOLKIČEV; DHONDT: 216; pl. 4, fig. 1; text-fig. 2a, b, c.

M a t e r i a l : Two internal moulds are at our disposal: NHMW 1979/2077/3, a left valve from the Gschliefgraben and NHMW 1998/96/1 from Tonionböden (Steiermark).




Figure 10: *Inoceramus cf. borilensis* JOLKIČEV with geniculation. NHMW 1979/2077/3; Gschliefgraben; x 1.




Figure 11: Diagram showing H/L and Vo/L- ratios at *Inoceramus cf. borilensis* JOLKIČEV.

Preservation: Highly incomplete. Only the beaks are preserved in both cases. Shell missing.

Description: The umbonal part is strongly erect and separated from the wing, slightly deformed by compaction. Strongly prosogyrate. In NHMW 1979/2076/3 there is a geniculation, which is associated with an increase in the undulation distances. Total angle: 105°.

Remarks An exact identification of the specimens from Austria is not possible. In both cases the shell is missing; this feature is very important for the examination of *Inoceramus borilensis* JOLKIČEV according to JOLKIČEV (1962). The specimens of Austria are comparable with *Inoceramus borilensis* JOLKIČEV from Houthalen determined by SORNAY (1982) and depos-

ited in the collection SORNAY under no. 10222 in Brussels (ISNB).

Occurrence: W- Europe (Belgium, France), E- Europe (Bulgaria), Gschliefgraben.

Stratigraphic distribution Maastrichtian (holotype – Bulgaria). Late Campanian *Nostoceras hyatti* Zone (Tercis/SW France). Early Maastrichtian *Pachydiscus epiplectus* Zone („*Inoceramus*“ cf. *borilensis*; Tercis/SW France).

***Inoceramus aff. bererensis* SORNAY
(Pl. 3, Fig. 1; Text-figs. 12–13)**

1975 *Inoceramus bererensis* SORNAY: 11–13; pl. 2, fig. 3; pl. 5, fig. 2; plate 6, fig. 1; Text-figs. 5–6.

M a t e r i a l Two internal moulds of one left (NHMW 1979/2076/44) and one right (SK/B/GG/1997/5) valve.

Preservation: Incomplete. Ventral margins and large parts of the wings are missing. SK/B/GG/1997/5 is highly deformed by compaction. The umbonal part is thrust over the anterior margin.

D e s c r i p t i o n Medium-sized to tall, shape elongated oval. Umbonal part is distinctly separated from the wing. Umbonal pole projecting above the hinge line. Hinge line straight to slightly convex. Geniculation at Ha = 98 mm (NHMW 1979/2076/44). At the hinge line the undulations are bent to the umbonal pole. Total angle: 100°; wing angle: 43°. Undulations flat rounded (Anwachsstreifen sensu HEINZ 1928). The growth axis is increasing from 40° to 60°– prosocline. Thickness of the shell 0.8–1.5 mm.





Figure 12: *Inoceramus aff. bererensis* SORNAY, SK/B/GG/1997/5 umbonal part thrust over the hinge line (compaction); x 1.

Remarks Shape more elongated than in *Inoceramus bererensis* SORNAY. The Na/Ha-ratio in *Inoceramus bererensis* SORNAY is between 80–90 in contrast to *I. aff. bererensis* SORNAY (text-fig. 13). *Inoceramus aff. bererensis* SORNAY (NODA & KANIE 1978 a; pl.1,



Na / Ha and Vo / Ha at *Inoceramus* sp. aff. *bererensis* SORNAY from the Gschliefgraben .

Figure 13: Diagram showing Na/Ha-, Vo/Ha -ratios and WA growth axis at *Inoceramus* aff. *bererensis* SORNAY.

Distance from UP	1	2
10–30 mm	—	—
30–50 mm	9.25 mm	6.50 mm
50–70 mm	13.00 mm	—

Table 9. Average undulation intervals (DU) of *Inoceramus* aff. *bererensis* SORNAY in different distances from the umbonal pole (UP). 1 = NHMW 1979/2076/44; 2 = SK/B /GG/1997/5.

fig. 3) is comparable but has an anterior auricle. In *Inoceramus* aff. *bererensis* SORNAY (TRÖGER & RÖHLICH 1992) the course of the undulations is more cycloid. Geographic distribution: Madagascar (SORNAY 1975), N-Africa (TRÖGER & RÖHLICH 1992). Stratigraphic distribution: Middle Campanian (SORNAY 1975).

Inoceramus sagensis OWEN 1852


(Pl. 2, Fig. 4; Text-fig. 14)

- 1852 *Inoceramus sagensis* OWEN: 582 ; pl. 7, fig. 3
- * 1876 *Inoceramus sagensis* OWEN var. *nebrascensis* OWEN-MEEK: 52; pl. 13, figs. 2a, b.
- 1959 *Inoceramus sagensis* OWEN; DOBROV & PAVLOVA :155; pl. 22, fig. 3; pl. 23, fig. 5.

M a t e r i a l Three internal moulds (one left = NHMW 1979/2076/26; two right ones = NHMW 1979/2076/33, NHMW 1979/2077/2/1) from the Gschliefgraben.

Preservation: Incomplete. Parts of the wing and of the ventral margin are missing. Slightly deformed by compaction.

D e s c r i p t i o n : Medium to tall in size, shape elongated subquadratic. The vaulted beak is distinctly separated from the wing. Umbonal pole prosogyrate. Total angle 120°. Anterior margin distinctly concave. Thick-



1 = NHMW / 1979 / 2076 / 33 (Gschliefgraben)
2 = NHMW / 1979 / 2076 / 26 (Gschliefgraben)
3 = US N.M. 485 (USA)

Variability of H / L , Vo / L and S / L at *Inoceramus sagensis* OWEN.

Figure 14: Diagram showing H/L-, Vo/L- and S/L- ratios at *Inoceramus sagensis* OWEN.

Distance from UP	1	2
10–30 mm	2.52 mm	3.33 mm
30–50 mm	4.20 mm	5.28 mm
50–70 mm	—	5.63 mm

Table 10. Average undulation intervals (DU) at different distances from the umbonal pole (UP) in NHMW 1979/2076/33 (1) and *Inoceramus sagensis* var. *nebrascensis* OWEN (2 USNM 485).

ness of shell: 1.5 mm. The interior moulds bear radial striae.

R e m a r k s The changes of the H/L- , Vo/L- and S/L- ratios are shown in text-fig. 14. They are comparable with the corresponding ratios of USNM no. 485 from the Fort Pierre Group. One part of an umbonal region was determined as *Inoceramus* aff. *sagensis* OWEN (NHMW 1979/2077/2, 1).

Geographic distribution: Caucasus, Crimea, N- America (DOBROV & PAVLOVA [in:] MOSKVIN 1959); Gschliefgraben.

Stratigraphic distribution: Late Campanian to Early Maastrichtian (DOBROV & PAVLOVA [in:] MOSKVIN 1959).

Genus *Cordiceramus* HEINZ 1932

T y p e s p e c i e s : *Inoceramus cordiformis* J. SOWERBY, 1823.

Cordiceramus ? aff. heberti (FALLOT)
 (Pl. 1, Fig. 2 ; Text-figs. 15, 16)

- * 1885 *Inoceramus heberti* FALLOT: 249; pl. 7, fig. 1
- 1968 *Inoceramus (Cordiceramus) heberti* FALLOT - SORNAY: 41; pl. 4, fig. 5.

The lectotype was designated by SORNAY (1968:41).

M a t e r i a l Internal moulds of two valves (NHMW 1997/z/160/2 , SK/B/GG/1997/6).

Preservation: Incomplete. Flattened by compaction, with radial cracks. The umbonal region of SK/B/GG/6 is missing.

D e s c r i p t i o n : Medium to large in size. Shape cycloid, flat. Umbo less prominent, not distinctly separated from the wing. Total angle: 120°. Undulations toprounded, sometimes sharp (compaction). Course of the undulations pentagonal subquadratic (e.g. SK/B/GG/1997/6). At the hinge line the undulations are bent to the umbonal pole (angle: 110–120°).

Anterior margin slightly concave. Growth axis: 45–58°, slightly prosoclinal. H/L- ratio (text-fig. 15) similiar to the lectotype. SK/B/GG/1997/6 bears an *Endocostea* scar (Text-fig. 15).




Figure 15: *Cordiceramus ? sp. aff. heberti* (FALLOT) with *Endocostea* scar- SK/B/GG/1997/6; Gschliefgraben; x 1.

Distance from UP	1	2
10–30 mm	—	—
30–50 mm	6.33 mm	5.25 mm
50–70 mm	7.03 mm	8.16 mm
70–90 mm	8.85 mm	9.00 mm

Table 11. Average undulation intervals of NHMW z/160/2 (1) in relation to the lectotype of *Cordiceramus heberti* FALLOT (2).

R e m a r k s In *Cordiceramus heberti* (FALLOT) the course of the undulations is pentagonal. The undulation distances are greater in the gerontic stages. The Gschliefgraben specimens may be compared with the Late Campanian *Inoceramus* aff. *heberti* FALLOT from Libya (TRÖGER & RÖHLICH, 1992).


Geographic distribution: W-Europe / Hautes Alpes), Spain (GALLEMI et al., 1993), N-Africa (TRÖGER & RÖHLICH, 1992).

Stratigraphic distribution: Late Campanian (SORNAY 1968, TRÖGER & RÖHLICH 1992).

Genus *Cremonoceramus* Cox 1969 (non HEINZ 1932)

T y p e s p e c i e s : *Inoceramus inconstans* Woods 1912

D i s c u s s i o n The genus *Sphaeroceramus* was established by HEINZ (1932) without formal description and illustration. According to HEINZ the type specimen of text-fig. 48 in Woods (1910–1912) should be chosen as the genotype, described as *Inoceramus inconstans* Woods (unknown locality, Upper Chalk, unknown horizon). This specimen was renamed *Sphaer-*



H/L, Vo/L and S/L at *Cordiceramus* sp. aff. *heberti* (FALLOT) from the Gschliefgraben in comparision with H/L at the lectotype (SORNAY 1980)

Figure 16: Diagram showing H/L-, Vo/L- and S/L-ratios at *Cordiceramus* ? aff. *heberti* (FALLOT)

ceramus pilula HEINZ, 1932. WALASZCZYK (1997) used the name *Sphaeroceramus*. He gave a short description of the genus. The type species according to WALASZCZYK is *Inoceramus subsarumensis* RENNGARTEN 1926 = *Sphaeroceramus pilula* HEINZ. Species of the genus *Sphaeroceramus* are *Sph. sarumensis* (Woods) and *Sph. subsarumensis* (RENNGARTEN). According to COX in MOORE & TEICHERT (1969) *Sphaeroceramus* is a synonym of *Cremonoceramus* COX 1969. We follow COX in this paper.

***Cremnoceramus sarumensis* (Woods)**
 sensu SORNAY 1982
 (Pl. 1, Fig. 1, Pl. 4, Fig. 4)

L e c t o t y p e : The original of Woods (1912, pl. 52, fig. 2a, b) designated by WALASZCZYK (1997), BMNH collection BLACKMORE.

- 1911 *Inoceramus incostans sarumensis* Woods: 293; pl. 52, figs. 2, 3
 * 1982 *Inoceramus sarumensis* Woods - SORNAY : pl. 2, fig. 2
 1997 *Sphaeroceramus sarumensis* (Woods, 1912); WALASZCZYK: 31; pl. 1, figs. 1-2, 5; pls. 21-22.

M a t e r i a l Internal mould of a single left valve (NHW 1997/z/160/1) and a bivalved but incompletely preserved specimen (NHW 1997/z/160/1). Preservation: Incomplete. Parts of the wings and of the ventral margins, highly deformed by compression, are missing.

D e s c r i p t i o n : Large, inequilateral. The beaks are weakly involute, prosogyrate. Anterior margin concave. The undulations are rounded and staircase-shaped.

Distance from UP	H / L	Vo / L	DU
10-30 mm	45.9	47.3	2.74 mm

Table 12. Average ratios H/L, Vo/L and average distances (DU) of the undulations in *Cremnoceramus sarumensis* (Woods) from the Gschliegraben. NHW 1997/z/160/1 at 10-30 mm distance from the umbonal pole (UP).

R e m a r k s : Concerning the involute character of the umbonal pole the specimens are completely comparable with *Cremnoceramus sarumensis* (Woods) from the Campanian of Belgium (charbonnage Zolder/charbonnage Limbourg), described by SORNAY (1982).

Stratigraphic distribution: Early Campanian (Woods 1911) – Smectite de Herve (Campanian – SORNAY 1982). After WALASZCZYK (1997) *S. sarumensis* occurs in the Early Campanien and early Late Campanian of Germany and does not occur in the higher Late Campanian *Polypliocum* Zone respectively *Haldemensis* Zone sensu WALASZCZYK (1997) of the Vistula valley and Westphalia.

***Cremnoceramus* ? aff. *inconstans* (Woods)**
 (Pl. 4, Fig. 5)

M a t e r i a l : A single right valve (NHW 1979/2077/4) from the Gschliegraben.

Preservation: Incomplete internal mould with adherent shell and radial cracks, flattened by compaction. The ventral margin, large parts of the wing and the umbonal pole are missing. The umbonal region is highly crushed.

D e s c r i p t i o n : Medium to tall in size, inequilateral. Prominent beak is projecting over the hinge line. Beak not distinctly separated from the wing. Anterior margin convex.

Undulations (concentric ribs) rounded. Growth lines cut the undulations at an acute angle (HEINZ, 1932 : 23, footnote – Anwachsschnittreifen). Distance between the growth lines 0.5–1.5 mm. Thickness of the shell 0.8–1 mm.

Distance from UP

	DU
10-30 mm	6.5 mm
30-50 mm	12.0 mm
50-70 mm	15.1 mm
70-90 mm	19.5 mm

Table 13. Average undulation intervals (DU) in different distances from the umbonal pole (UP); NHW 1979/2077/4 from the Gschliegraben.

Distance from UP

	H/L	Vo/L
10-30 mm	45.0 mm	45.5 mm
30-50 mm	57.9 mm	44.8 mm
50-70 mm	65.9 mm	57.6 mm

Table 14. Average ratios H/L and Vo/L at 10–70 mm distance from the umbonal pole (UP); NHW 1979/2077/4 from the Gschliegraben.

R e m a r k s : The shape, the erect beak region and the distance between the undulations indicate that the specimen is a member of the *Cremnoceramus inconstans* group. The specimen differs from all described species by the „Anwachsschnittreifen“.

Conclusion

The diverse Inoceramid fauna indicates Late Campanian through Early Maastrichtian (Tab. 15). This is in accordance with nannoplankton ages (CC22b – CC23c; WAGREICH, 1999, this volume), ammonites (KENNEDY & SUMMERSBERGER, 1999, this volume) and Echinodermata (JAGT, 1999, this volume). In terms of ammonite zones the age is from *Phaleratum* to *Polypliocum* Zone. The bulk of the Inoceramids seems to be from the *Haldemensis* Zone (= *Polypliocum* Z.). Early Maastrichtian age cannot be excluded by Inoceramids. A complete list of fossils present in the Late Cretaceous Gschliegraben locality is given by KENNEDY & SUMMERSBERGER, 1999 (this volume).

Acknowledgements

Our thanks are due to Prof. Norbert Vavra (University of Vienna, PIUW), who donated a collection of Inoceramidae to the Naturhistorisches Museum Wien in 1976. The staff of the Naturhistorisches Museum Wien assisted in collecting in the field. Photographing, drafting and preparing of the material was also done by technicians of the Museum. Herbert Summesberger is grateful for a grant from the Theodor Körner Stiftungsfonds. We are especially grateful to John W.M. Jagt (Maastricht) for careful reviewing of the manuscript. Financial support was given by the Freunde des Naturhistorischen Museums Wien for the printing of the plates. The work in

Chronostratigraphy	Biostratigraphy NW Germany	Inoceramid assemblage zones	IZ	Range of Gschlieffgraben Inoceramids
EARLY MAASTRICHTIAN	<i>Lanceolata</i> Zone			
	<i>Grimmenensis/Granulostos</i> Zone			
	<i>Langzeit</i> Zone			
	<i>Polyplacum</i> Zone	<i>Cat. haldemensis</i> Zone	33	
LATE CAMPANIAN	<i>Vulgaris</i> Zone	Unnamed Zone		
	<i>Basiplana/Spiniger</i> Zone	<i>Cat. vorhelmensis</i> Z. - <i>Inoc. adjaciensis</i> Z.	32	
	<i>Conica/Mucronata</i> Zone	<i>Cat. beckumensis</i> Zone	31	
	<i>Gracilis/Mucronata</i> Zone			
	<i>Conica/Gracilis</i> Zone	<i>Sph. strumensis</i> Zone -		
	<i>Papillosa</i> Zone		30	
EARLY CAMPANIAN	<i>Senonensis</i> Zone	<i>Cat. duriensis</i> Zone		
	<i>Pilula/Senonensis</i> Zone			
	<i>Pilula</i> Zone			
	<i>Lingua/Quadrata</i> Zone	<i>Sphe. patootensiformis</i> Zone	29	
	<i>Granulata/quadrata</i> Zone			
LATE SANTONIAN	<i>Marsupites/Granulata</i> Zone			
		<i>Sphe</i> - <i>Sphenocerasmus</i> <i>Sph.</i> - <i>Sphaerocerasmus</i> <i>Cai.</i> - <i>Cataceramus</i>		
				Biostratigraphical range of the Gschlieffgraben on the NW German biostratigraphical zonal scheme including the Subhercynian Cretaceous and the Münsterland Basin according to CHRISTENSEN (1988, 1997), ERNST (1964, 1970 b, 1972, 1974), KAUFFMAN et al. (1993), PETRASCHEK (1906), RIEDEL (1931), SCHULZ & SCHMID (1983), SCHULZ (1979), ULBRICH (1971) and unpublished data.
				<i>Inoceramus sagittensis</i> OWEN <i>Inoceramus cf. boettgeri</i> JOLIKOEVER <i>Inoceramus regularis</i> D'ORBIGNY <i>Endocostea impressa</i> (D'ORBIGNY) <i>Cataceramus ballicus</i> (BOHM) subspp. indet. <i>Cataceramus ballicus aff. haldeensis</i> (GIERAS) <i>Cataceramus ballicus ellipticus</i> (GIERAS) <i>Inoceramus cf. boettgeri</i> JOLIKOEVER <i>Inoceramus saruwamensis</i> (WOODS sensu SORKANY 1982)

Table 15. Stratigraphic range of the Gschlieffgraben Inoceramidae.

Freiberg was partly sponsored by the Deutsche Forschungsgemeinschaft. We are highly indebted to the staff of the National Museum Washington (USA), the Institut Royal des Sciences Naturelles de Belgique and the Muséum National d'Histoire Naturelle Paris for providing casts of type material. We thank Dr. Peter Baumgartner (Traunkirchen) for reviewing the sketchmap of the Gschliefgraben.

References

- AYYASAMI, K. & RAO, B. R. J., nnn in press. Dispersal of Cretaceous Inoceramids in Gondwanaland. — 9th International Symposium: 341–356, Hyderabad.
- BEYENBURG, E., 1936. Die Fauna der Halterner Sandfazies im westfälischen Unterenon. — Jb. Preuß. geol. Landesanst., **57**:284–332, 4 figs., pls. 11–13, Berlin.
- BLANK, M.J., KRIMGOLTZ, G. J., NAIDIN, D. P. & SAVTSCHINSKAJA, O.V., 1974. Atlas verchnemelovoi fauni Donbassa (russ.). — „Nedra“ Moskva, 638 pp, 128 pls., Moskva.
- BLASZKIEWICZ, A., 1980. Campanian and Maastrichtian ammonites of the Middle Vistula Valley, Poland: a stratigraphic-paleontological study. — Prace Inst. Geol. **92**: 1–63, 56 pls., Warszawa.
- BÖHM, J., 1907. Über Haenleinia n. subgen. — Z. dt. geol. Ges., **59**:317, Berlin.
- BÖHM, J., 109. Über Inoceramus crippsii auct. — Abh. Preuß. geol. Landesanst., **56**:39–58; pls., 9–14, Berlin.
- CHRISTENSEN, W.K., 1998. Upper Campanian *Belemnitella* from Austria. — Beitr. Paläont., **22**:3–21, 1 pl., Wien.
- COX, L. R., 1969 [in:] MOORE, R.C. & TEICHERT, C. Treatise on Invertebrate Paleontology, Part N. — Mollusca 6, Bivalvia, Lawrence (Univ. Kansas Press).
- DHONDT, A.V., 1983. Campanian and Maastrichtian inoceramids: A review. — Zitteliana, **10**:689–701, 6 figs., München.
- DHONDT, A.V., 1993. Upper Cretaceous bivalves from Tercis, Landes, SW France. — Bull. Inst. Roy. Sci. Nat. Belg., Sci. de la Terre, (Univ. Kansas Press) (Univ. Kansas Press) **63**:211–259, 7 pls., 17 figs., Brussels.
- DOBROV, S. A. & PAVLOVA, M. M. [in:] MOSKVIN, M. M. (ed.) 1959. Atlas verchnemelovoi fauni severnogo Kavkaza i Krima (russ.), Inocerami. — Tr. VNIGAS: 130–165, 23 pls., Moskva.
- FALLOT, J. E., 1885. Etude géologique sur les étages moyens et du Terrain Crétacé dans le Sud-est de la France. — These: 269 pp., 8 figs., Paris.
- FRAAYE, R. H. B. & SUMMESBERGER, H., 1999. New crustacean records from the Late Campanian of Austria. — Beitr. Paläont., **24**:1–6, Wien.
- GALLEMI, J., LOPEZ, G. & MARTINEZ, R. 1993. Upper Cretaceous Macrofauna from southeast Spain. 1st General Meeting Project 362 Tethyan and Boreal Cretaceous, Coimbra, 24/30 October 1993:15–118 (abstract).
- GIERS, R. 1964. Die Großfauna der Mukronatenkreide im östlichen Münsterland. Die Kreide Westfalens. Ein Symposium. — Fortschr. Geol. Rheinld. U. Westf., **7**:213–294; 8 pls., 10 figs., 3 tab., Krefeld.
- HANCOCK, J. M., PEAKE, N. B., BURNETT, J., DHONDT, A.V., KENNEDY, W. J., STOKES, R. B., 1993. High Cretaceous biostratigraphy at Tercis, southwest France. — Bull. de l' Institut Royal des Sciences Natur de Belgique, Sci. de la Terre, **63**:133–148, Brussels.
- HEINZ, R., 1928. Über die bisher wenig beachtete Skulptur der Inoceramen-Schale. — Mitt. Min. Geol. Staatsinst. Hamburg **10**: 3–39, 3 pls., 5 figs., Hamburg.
- HEINZ, R., 1932. Aus der neuen Systematik der Inoceramen. — Mitt. Min. Geol. Staatsinst. Hamburg **13**:1–26., Hamburg.
- JAGT, J.W.M., 1999. Late Campanian echinoids and crinoids from the Gschliefgraben (Ultrahelvetic, Austria). — Beitr. Paläont., **24**:7–22, Wien.
- JOLKIČEV, N., 1962. Inoceramen aus dem Maastricht Bulgariens. — Trav. sur la Géologie de Bulgarie. Sér. Paléont., **4**:134–169, 8 pls., Sofia.
- KAUFFMAN, E.G., SAGEMANN, B. B., KIRKLAND, J. I., ELDER, W.P., HARRIES, P.J. VILLAMIL, T., 1993. Molluscan biostratigraphy of the Cretaceous Western Interior basin, North America. [in:] CALDWELL, W.G.E. & KAUFFMAN, E.G. (eds.): Evolution of the Western Interior Basin. — Geol. Ass. of Canada, Spec. Paper, **39**:397–434.
- KENNEDY, W.J., 1993. Campanian and Maastrichtian ammonites from the Mons Basin and adjacent areas (Belgium). — Bull. Inst. Roy. Sci. Nat. Belgique, Sci. de la Terre, **63**:99–131, Brussels.
- KENNEDY, W. J. & SUMMESBERGER, H., 1984. Upper Campanian ammonites from the Gschliefgraben (Ultrahelvetic, Upper Austria). — Beitr. Paläont. Öst., **11**:149–206, pls. 1–14., Wien.
- KENNEDY, W. J. & SUMMESBERGER, H., 1999. New Upper Campanian Ammonites from the Gschliefgraben near Gmunden (Ultrahelvetic, Austria) — Beitr. Paläont. **24**:23–39, Wien.
- MEEK, F.B., 1876. A report on the Invertebrate and Tertiary Fossils of the Missouri country. — Report of the U.S. Geol. Surv. of the Territories (ed. HAYDEN, F.V.), **9**:1–629, 45 pls., Washington.
- NODA, M. & KANIE, Y., 1978a. Campanian *Inoceramus* from the Menabe area, southwestern Madagascar, part 1, **4**(1):1–32, 4 pls., Tokyo.
- NODA, M. & KANIE, Y., 1978b. Campanian *Inoceramus* from the Menabe area, southwestern Madagascar, part 2, **4**(2):63–72; 4 pls., Tokyo.
- D'ORBIGNY, A., 1843–47. Paléontologie Francaise. Description zoologique et géologique de tous les animaux mollusques et rayonnés fossiles de France. — Terrains crétacés, **3**: 807 pp., pls. 237–489, Paris.
- OWEN, D. D., 1852. Description of new and imperfectly known Genera and Species of organic remains, collected during the Geological Surveys of Wisconsin,

- Iowa and Minnesota. — Rep. Geol. Surv. Wisconsin, Iowa and Minnesota: 638 pp., 46 pls., geol. maps and sections, Philadelphia.
- PREY, S., 1983. Das Ultrahelvetikum — Fenster des Gschließgrabens nördlich von Gmunden (Oberösterreich). — Jahrb. Geol. B.A., **126**(1):1–95, 4 figs., 1 geol. map., Wien.
- SCHROEDER, H. & BÖHM, J., 1909. Geologie und Paläontologie der subherzynen Kreidemulde. — Abh. Preuß. Geol. Landesanst. N.F., **56**:59–64, Berlin.
- SEITZ, O., 1961. Die Inoceramen des Santons von Nordwestdeutschland. I.Teil: (Die Untergattungen *Platyceramus*, *Cladoceramus*, *Cordiceramus*) — Beih. Geol. Jahrb., **46**:186 pp, 15 pls, 39 figs., Hannover.
- SEITZ, O., 1967. Die Inoceramen des Santon und Unter-Campan von Nordwestdeutschland. III.Teil: Taxonomie und Stratigraphie der Untergattungen *Endocostea*, *Haenleinia*, *Platyceramus*, *Cladoceramus*, *Selenoceramus* und *Cordiceramus* mit besonderer Berücksichtigung des Parasitismus bei diesen Untergattungen. — Beih. Geol. Jahrb., **75**:171 pp., 27 figs, 8 tab., 27 pls., Hannover.
- SORNAY, J., 1957. *Inoceramus impressus* d'ORBIGNY. — Palaeontologia universalis, N.S. **129**, Paris.
- SORNAY, J., 1962. Etude d'une faune d'inocerames du Senonien supérieur des Charantes et description d'une espece nouvelle du Senonien de Madagascar. — Bull. soc. géol. France, **4**:118–122, Paris.
- SORNAY, J., 1968. Inocerames Senonien du Sud-Ouest de Madagascar. — Ann. Paléont. (Invertebres), **54**:25–47, pls. A-H., Paris.
- SORNAY, J., 1975. Trois especes nouvelles d'inocerames du Senonien de Madagascar. — Ann. de Paléont. (Invertébrés) **61**, fasc. 1:19–29; 6 pls., Paris.
- SORNAY, J. 1976. La Faune d'Inocerames de Dau (Region de Royan, Charente Maritime) et Remarques sur deux espèces de d'ORBIGNY *I. regularis* et *I. goldfussi*. — Ann. de Paléont., **62**, fasc., 1:1–18, Paris.
- SORNAY, J. 1982. Sur la faune d'Inocerames de la Smectite de Herve (Campanien) et sur quelques Inocerames du Campanien et Maastrichtian de la Belgique. — Bull. Inst. Royal Sci. Nat. de Belgique, Sci. de la Terre, **54**/4:1–15; 3 figs., 4 pls., Bruxelles.
- SUMMESBERGER, H., WAGREICH, M., TRÖGER, K.-A. & JAGT, J. W. M., 1999. Integrated biostratigraphy of the Santonian/Campanian Gosau Group of the Gams Area (Late Cretaceous; Styria, Austria). — Beitr. Paläont., **24**:153–203, Wien.
- TRÖGER, K.-A., 1967. Bemerkungen zur Variabilität von *Inoceramus balticus* J. BÖHM aus der subherzynen Kreide. — Freiberger Forschungsh., **C213**:1–21, 2 pls., Leipzig.
- TRÖGER, K.-A., 1989. Problems of Upper Cretaceous Inoceramid Biostratigraphy and Paleobiogeography in Europe and Western Asia.. — Cretaceous of the Western Tethys. — Proc. 3rd Int. Symp. Cret., Tübingen, **1987**: 911–930, 8 figs., Stuttgart.
- TRÖGER, K.-A., 1998. Remarks concerning morphometric parameters, biostratigraphy and palaeobiogeography of Turonian inoceramids (Bivalvia) in Europe. — Zbl. Geol. Paläont., Teil T, **1996**(11–12):1489–1499, 5 figs., Stuttgart.
- TRÖGER, K.-A. & RÖHLICH, P., 1982. Zur Variabilität von *Inoceramus balticus haldemensis* GIERS. — Freiberger Forschungsh., **C 375**:101–111, 9 figs., 1 pl., Leipzig.
- TRÖGER, K.-A. & RÖHLICH, P., 1992. Campanian-Maastrichtian Inoceramid (Bivalvia) Assemblages from NW-Libya. [in:] The Geology of Libya, V: 1357–1381, 19 figs., 5 pls., Tripoli.
- ULBRICH, H., 1971. Mitteilungen zur Biostratigraphie des Santon und Campan des mittleren Teils der Subherzynen Kreidemulde. — Freiberger Forschungsh., **C 267**:47–60, 5 pls., Leipzig.
- VOIGT, S., 996. Paläobiogeographie oberkretazischer Inoceramen und Rudisten. Ozeanographische und klimatologische Konsequenzen einer neuen Paläogeographie. — Münchner Geowissenschaftl. Abh., Reihe A, **31**:1–101, München.
- WAGREICH, M., 1999. Calcareous nannofossil assemblages from the Gschließgraben near Gmunden (Ultrahelvetic, Austria). — Beitr. Paläont., **24**:63–67, Wien.
- WALASZCZYK, I., 1997. Biostratigraphie und Inoceramen des oberen Unter-Campan und unteren Ober-Campan Norddeutschlands. — Geol. u. Paläont. in Westfalen, **49**:111 pp., 21 Text-figs., 32 pls., Münster.
- WARD, P.D. & KENNEDY, W.J., 1993. Maastrichtian Ammonites from the Biscay Region (France, Spain). — Journ. Paleont., The Paleont. Soc. Memoir, **34**:58 pp., 45 figs., Tulsa.
- WOODS, H., 1899–1913. A Monograph of the Cretaceous Lamellibranchia of England. — Monogr. Palaeontogr. Soc., **I**:232 pp., 42 pls., 7 Figs., **II**:473 pp., 62 pls., 252 figs., London.

Plate 1

Fig. 1. *Cremnoceramus sarumensis* (WOODS); NHMW 1997/z/160/1;

Fig. 2. *Cordiceramus*? aff. *heberti* (FALLOT); NHMW 1997/z/160/2, completely flattened by compaction, with radial cracks

All from Gschließgraben ; all x 1

PLATE 1




PLATE 2

Fig. 1, 3, 5, 6: *Endocostea impressa* (D'ORBIGNY);

Figs. 1, 3: NHMW 1979/2076/25/1, left valve; Figs. 5, 6: NHMW 2076/25/2; umbonal region of the right valve.

Fig. 2: *Inoceramus regularis* D'ORBIGNY; NHMW 1979/2076/39.

Fig. 4: *Inoceramus sagensis* OWEN; NHMW 1979/2076/26.

All from Gschliefgraben/Gmunden. Figs. 1–5: x 1; Fig. 6: x 2

PLATE 2




Plate 3

Fig. 1: *Inoceramus* aff. *bererensis* SORNAY; NHMW 1979/2076/44; internal mould of a left valve with adherent shell.

Fig. 2: *Cataceramus balticus* (BÖHM) subsp. indet. with *Endocostea* scar (x); NHMW 1979/2076/24.

Fig. 3, 4: *Cataceramus balticus* aff. *haldemensis* (GIERS); SK/B/GG/1997/4; straight to the growth axis; deformed by compaction.

All from Gschliefgraben. All x 1.

PLATE 3




Plate 4

- Fig. 1: *Cataceramus balticus balticus* (J. BÖHM); NHMW 2076/46 ; umbonal region flattened by compaction.
- Fig. 2: *Cataceramus balticus balticus* (J. BÖHM) with geniculation in the umbonal region NHMW 1979/2076/37.
- Fig. 3 *Cataceramus balticus* (J. BÖHM) subsp.indet.,bivalved specimen completely flattened by compaction – NHMW 1979/2076/28.
- Fig. 4 *Cremnoceramus sarumensis* (WOODS); umbonal region; NHMW 1997/z/160/1.
- Fig. 5: *Cremnoceramus* ? aff. *inconstans* Woods; NHMW 1979/2077/ 4.
- All from Gschliefgraben, all 1 x.

PLATE 4

