

in the Caribbean. Differences in coral diversity among the Pleistocene reef sites are even higher than modern differences, indicating that reef diversity in the Caribbean remained unstable after several extinction events ~1 myr ago. By contrast, high-diversity communities of Indo-Pacific Pleistocene reefs appear more stable through the Pleistocene. The reasons for that dichotomy are still unexplored but may be sought in the geographic isolation of the Caribbean.

<sup>1)</sup> Museum für Naturkunde, Leibniz Institute for Research on Evolution and Biodiversity at the Humboldt-Universität Berlin, Invalidenstr. 43, 10115 Berlin, Germany, e-mail: heike.mewis@mfn-berlin.de, wolfgang.kiessling@mfn-berlin.de

<sup>2)</sup> Centre for Marine Science, School of Biological Sciences, and Australian Research Council Centre of Excellence for Coral Reef Studies, The University of Queensland, Brisbane, Queensland 4072, Australia, e-mail: j.pandolfi@uq.edu.au

---

### Freies Thema

## Untermiozäne Biber aus Japan: Endemismen und biogeographische Verbindungen

Thomas Mörs<sup>1)</sup>, Y. Tomida<sup>2)</sup> & Irina Ruf<sup>3)</sup>

Terrestrische Kleinsäuger sind im Untermiozän von Japan extrem selten, einige Ordnungen wie z.B. Eulipotyphla und Lagomorpha sind nur durch ein oder zwei Funde belegt. Nagetiere sind mit neun Taxa aus drei oder vier Familien vertreten: ?Sciuridae, Castoridae, Eomyidae und Diatomyidae. Biber (Castoridae) sind die am besten dokumentierte Gruppe, sowohl was taxonomische Diversität als auch Anzahl der Fundstücke betrifft. Das meiste hier vorgestellte Material stammt aus der Nakamura-Formation (Mizunami-Gruppe) des Kani-Beckens in der Gifu-Prefektur, Zentraljapan. Weiteres Material stammt aus der Toki- und Akeyo-Formation (Mizunami-Gruppe) des benachbarten Mizunami-Beckens, sowie aus der Oya- und Fukazuki-Formation (Nojima-Gruppe) von Sasebo in der Nagasaki-Prefektur. Radiometrische, magnetostratigraphische und marin-biostratigraphische Untersuchungen haben für die verschiedenen Kleinsäuger-Fundstellen innerhalb der Mizunami-Gruppe numerische Alter von 19,5 bis 17,5 Ma ergeben. Daher werden die Fundstellen mit der Ostasiatischen Neogenen Säugereinheit MNU 4 korreliert, was den Europäischen Neogenen Säugereinheiten MN 3

bis MN 4 entspricht. Das fossile Bibermaterial, bestehend aus einem fast vollständigen Schädel, Unterkiefern sowie isolierten Knochen und Zähnen, repräsentiert mindestens vier Taxa sehr kleiner bis extrem grosser Castoriden. Der größte ist *Youngofiber sinensis*, ein riesiger castoroidiner Biber, der zuerst aus Sihong in der Jangsu-Provinz in Ostchina beschrieben wurde. Eine andere grosse Form belegt einen primitiven anchitheriomynen Biber, gen. et sp. nov. Er ist deutlich kleiner als der nordamerikanische *Amblycastor*, und auch kleiner als die eurasische *Anchitheriomys*. Eine dritte, mittelgroße Form dokumentiert einen fortschrittlichen palaeocastorinen Biber, gen. et sp. nov., der hochgradig an fossoriale Lebensweise angepasst ist. Die vierte Art repräsentiert einen sehr kleinen castoroidinen Biber, der möglicherweise mit *Euroxenomys* verwandt ist. Besonders interessant ist die Paläobiogeographie der japanischen Biber: mindestens zwei der vier Taxa aus der Mizunami-Gruppe – der anchitheriomynen und der palaeocastorinen Biber – sind endemisch in Japan, aber haben ihre nächsten Verwandten in Nordamerika. Der Riesenbiber *Youngofiber sinensis* ist endemisch in Ostasien und der kleinste Biber könnte eine biogeographische Verbindung mit Europa andeuten. Das paläobiogeographische Muster der Castoriden entspricht dem der anderen Kleinsäuger der Mizunami- und Nojima-Gruppe, möglicherweise ein Resultat der Trennung des japanischen Inselbogens vom ostasiatischen Festland während des Untermiozäns.

<sup>1)</sup> Swedish Museum of Natural History, P.O. Box 50007, SE-104 05 Stockholm, Sweden, e-mail: thomas.moers@nrm.se

<sup>2)</sup> National Museum of Nature and Science, Shinjuku, Tokyo 169-0073, Japan, e-mail: y-tomida@kahaku.go.jp

<sup>3)</sup> Steinmann-Institut, Universität Bonn, Bonn, Germany, e-mail: irina.ruf@uni-bonn.de

---

### Freies Thema

## A new Eocene lizard from Messel and the origin of *Amphisbaenia*

Johannes Müller<sup>1)</sup>, Christy A. Hipsley<sup>1)</sup>, Michael Wuttke<sup>2)</sup>, Jason J. Head<sup>3)</sup>, Robert R. Reisz<sup>3)</sup>, Nikolay Kardjilov<sup>4)</sup> & André Hilger<sup>4)</sup>

Molecular investigations have dramatically challenged the traditional view of squamate phylogeny. Among the most prominent changes in paradigm has been the

hypothesis of a monophyly of amphisbaenians (or worm lizards) and lacertids due its significant implications for the proposed relationship between amphisbaenians and snakes, as well as other limbless squamates. Until recently, however, there was no morphological or fossil support for an amphisbaenian-lacertid grouping, questioning the validity of this new hypothesis. A new genus and species of Eocene lizard from the famous Messel locality, consisting of an almost complete specimen, now sheds new light on this issue. The taxon is characterized by a spatula-like snout with a modified premaxilla and maxilla, notably small and anteriorly positioned orbits, the lack of a tympanic crest, a low tooth count, and reduced manus and pes. Also, it shows several features otherwise found only in amphisbaenians, such as the shape of the frontal suture and the morphology of the subolfactory processes. In a phylogenetic analysis including both morphological and molecular characters as well as extant and fossil taxa, the taxon groups as sister taxon to amphisbaenians within a lacertid-amphisbaenian clade and refutes a relationship between worm lizards and snakes, providing for the first time morphological support for the recently proposed molecular hypothesis. Morphometric analysis of body shape and ecology among squamates places the new taxon in cryptic or opportunistically burrowing habitats, and its distinctive anatomy indicates that head-first burrowing evolved prior to body elongation in *Amphisbaenia*.

The new taxon is considered to be a remnant of a non-crown group, amphisbaenian-like squamate lineage that occurred together with crown lacertids and amphisbaenians, which both are known from contemporaneous and even older Paleogene strata. Interestingly, many currently known lizards from Messel seem to be related to either Mesozoic squamate lineages or represent stemgroup taxa outside of the crown, suggesting that Europe may have acted as refuge for ancient squamate lineages during the Paleogene. Our study emphasizes that fossils are crucial for testing hypotheses of phylogenetic relationships, because extant taxa often reveal only a restricted picture of their evolutionary history.

<sup>1)</sup> Museum für Naturkunde – Leibniz-Institut für Evolutions- und Biodiversitätsforschung an der Humboldt-Universität zu Berlin, Invalidenstr. 43, D-10115 Berlin, Germany

<sup>2)</sup> Generaldirektion Kulturelles Erbe RLP, Direktion Landesarchäologie, Referat Erdgeschichte, 55116 Mainz, Germany

<sup>3)</sup> Department of Biology, University of Toronto, Mississauga, Ontario L5L 1C6,2, Canada

<sup>4)</sup> Helmholtz Centre Berlin for Materials and Energy, 14109 Berlin, Germany

---

## Zukunftspreis

### Middle Miocene freshwater gastropods from the Dinaride Lake System: punctuated morphological disparity as result of rapid climate change

Thomas A. Neubauer<sup>1)</sup>, Oleg Mandić<sup>1)</sup>, Mathias Harzhauser<sup>1)</sup>, Davor Pavelić<sup>2)</sup> & Alan Vranjković<sup>2)</sup>

This investigation focuses on evolutionary patterns and processes in Middle Miocene freshwater gastropods from the Dinaride Lake System. The point of study was a c. 100 m thick succession in the Sinj basin, comprising two limestone-coal cycles, which pass from fossil-poor limestone-dominated deposits into highly fossiliferous layers rich in organic matter. Within this section four gastropod lineages of the two unrelated genera *Melanopsis* and *Prososthenia* occur, each encompassing 3 or 4 morphotypes, which differ in size and/or sculpture. The single morphotypes are largely confined to certain intervals. Two periods of enhanced accumulation of organic matter mark distinct steps in the gastropod's shell morphology. During both intervals clear changes towards strongly sculptured and/or larger shells are observed. Both events happen within few meters (< 8 ka) and are accompanied by shifts to more arid conditions evident from the palynological record.

The synchronicity of morphological and environmental shifts suggests extrinsic factors to be responsible for the observed patterns. Probably, the tempo and intensity of the climatic changes exceeded the ecological thresholds of the discussed taxa leading to adaptive evolution. Due to increased aridity, a lowered lake level can be expected having strong influence on habitat segregation and vegetation. Enhanced photosynthetic activity led to chemosynthetically induced carbonate precipitation, which is evident from the section, and thus to over-saturation of the lake water. This might have facilitated the development of larger and sculptured shells.

<sup>1)</sup> Geological-Paleontological Department, Natural History Museum Vienna, Burgring 7, A-1010 Vienna, Austria, e-mail: thomas@indmath.uni-linz.ac.at

<sup>2)</sup> Faculty of Mining, Geology and Petroleum Engineering, University of Zagreb, Pierottijeva 6, 10000 Zagreb, Croatia