

Georgia-Augusta, wie auch Unter-Aufseher (1776), später dann (ab 1812) bis zu seinem Tode im Jahre 1840 1. Aufseher und Direktor des 1773 gegründeten „Königlich-Academischen Museums“ Aufgrund seiner zahlreichen Kontakte mit in- (z.B. Leopold v. Buch, Georg Forster, Johann Wolfgang v. Goethe, Alexander v. Humboldt, Samuel Thomas v. Soemmerring) und ausländischen Wissenschaftlern (z. B. Sir Joseph Banks, Jean-Baptiste Biot, Alexandre Brongniart, William Buckland, Georges Cuvier), aber auch wichtigen politischen Persönlichkeiten (u.a. Christian VIII. von Dänemark und Norwegen; Ludwig I. von Bayern), kommt Blumenbach eine Schlüsselrolle in der Entwicklung des Göttinger Universitätsmuseums, wie auch bei der Akquisition von Sammlungsmaterial zu. Darunter befindet sich auch zahlreiches Material verschiedenster Expeditionen jener Zeit, wie z.B. von Carsten Niebuhrs Arabienreise (1761–1767), den Cookschen Südsee-Reisen (1768–1771, 1772–1775, 1776–1780), der Billings-Saryčev Expedition (1785–1795) nach Ostasien und Alaska, oder der ersten russischen Weltumsegelung (1803–1806) unter v. Krusenstern.

Unzähliges Beleg- und Abbildungsmaterial sowie viele zoologische, botanische und mineralogische Typen, u. a. zu Blumenbachs umfangreichen Werken, wie „De generis humani varietate nativa liber“ (1776, 1781, 1795), und dem „Handbuch der Naturgeschichte“ (in 12 Auflagen zwischen 1779–1830, sowie Übersetzungen in 6 Sprachen), den „Decas collectionis suaे craniorum diversarum gentium illustrata“ (1790, 1793, 1795, 1800, 1820) und „Abbildungen naturhistorischer Gegenstände“ (1796–1800, 1802, 1804, 1810, 1827, 1830) wie auch „Specimen archaeologiae tellurio terrarumque imprimis Hannoveranarum“ (1803–1816), befindet sich ebenfalls darunter. Gut 6000 Sammlungsobjekte aus den Bereichen: (1) Anatomie, Medizin & Anthropologie, (2) Zoologie und Paläozoologie, (3) Botanik, Pharnakognosie & Paläobotanik, (4) Geologie, Mineralogie, Meteoritenkunde & Gemmologie sowie (5–6) Archäologie und Ethnologie konnten bisher lokalisiert werden. Annähernd 85% davon sind den Geowissenschaften zuzuordnen.

Das auf 15 Jahre angelegte Langzeitprojekt „Johann Friedrich Blumenbach – online“ startete im Januar 2010 und ist ein Vorhaben im Rahmen des Akademienprogramms der Union der deutschen Akademien der Wissenschaften, angesiedelt bei der Akademie der Wissenschaften zu Göttingen. Insgesamt drei institutionelle Hauptpartner, die Akademie der Wissenschaften zu Göttingen, die Staats- und Universitätsbibliothek Göttingen sowie die Georg-August-Universität als Eigentümerin eines Großteils der Sammlungen, sind beteiligt. Die Objektdigitalisierung (2D & 3D) wird hierbei vom Geowissenschaftlichen Museum der Universität Göttingen koordiniert und durchgeführt. Ziel des gesamten Projektes ist die Ermittlung, Erschließung und Vergegenwärtigung der Schriften und naturhistorischen Sammlung Johann Friedrich Blumenbachs sowie seiner internationalen Rezeption in einem Internetportal nebst

virtueller Forschungsumgebung, welches eine weltweite Kooperation bei der Nutzung, Erforschung und Kommentierung der Werke und Sammlungen Blumenbachs ermöglicht.

-
- ¹⁾ Geowissenschaftliches Museum, Georg-August-Universität Göttingen, Goldschmidtstr. 1-5, D-37077 Göttingen, e-mail: mreich@gwdg.de
- ²⁾ Geowissenschaftliches Zentrum, Abt. Geobiologie, Georg-August-Universität Göttingen, Goldschmidtstr. 3, D-37077 Göttingen
- ³⁾ Johann-Friedrich-Blumenbach-Institut für Zoologie und Anthropologie, Abt. Historische Anthropologie und Humanökologie, Georg-August-Universität Göttingen, Bürgerstr. 50, D-37073 Göttingen, e-mail: bgrossk1@gwdg.de
- ⁴⁾ Albrecht-von-Haller-Institut für Pflanzenwissenschaften, Abt. Systematische Botanik (mit Herbarium), Georg-August-Universität Göttingen, Untere Karspüle 2, D-37073 Göttingen, e-mail: jheinri@uni-goettingen.de
- ⁵⁾ Institut für Ethnologie, Ethnologische Sammlung, Georg-August-Universität Göttingen, Theaterplatz 15, D-37073 Göttingen, e-mail: gkruegel@gwdg.de
- ⁶⁾ UMG, Zentrum 1 Anatomie, Abt. Anatomie und Embryologie, Georg-August-Universität Göttingen, Kreuzbergring 36, D-37075 Göttingen, e-mail: mschult1@gwdg.de

Studentenpreisposter

Molluscs from underwater meadows – An Early Miocene seagrass mollusc community from Java, Indonesia

Sonja Reich¹⁾

A highly diverse and exceptional well preserved fossil molluscan assemblage has been found in a marine sandstone deposit near the village of Banjung-Ante (Yogyakarta, central south Java). The fauna is of Late Burdigalian (Early Miocene) age according to the associated larger benthic foraminifers. The molluscan assemblage contains predominantly small to minute gastropods. More than 3500 individuals assigned to 140 species were found. Bivalves are present with only 23 species and 325 individuals (with

one valve counting for a half individual). For a further investigation of the ecological composition the gastropods and bivalves of the Banjung Ante fauna were assigned to six different feeding guilds. The percentage of each feeding guild is illustrated in terms of species numbers and abundance (number of individuals).

In terms of abundance the fauna is dominated by grazers and detritivores that make up 69%. This group includes various cerithiid and rissoid species, as well as the genus *Bothropoma*. The abundance of small, herbivorous gastropods seems characteristic for seagrass associated assemblages with many of them grazing on the micro-algae growing on seagrass blades. Also present is the snail *Smaragdia* that appears to feed exclusively on leafs of seagrasses.

In terms of species numbers carnivores s.s. are the most common; they make up 36%. Another group of parasitic and browsing carnivores makes up an additional 24% of the species numbers. The herbivores/detritivores make up only about 25% of the fauna in species numbers. Although chemosymbiotic lucinids make up only 2% of the abundance, their presence indicates dysoxic settings in the seafloor.

Currently we are assembling data for modern and fossil seagrass faunas in order to compare to the Banjung Ante fauna. The latter has a very similar abundance/diversity composition as modern seagrass faunas found so far. The Banjung Ante fauna has a very dissimilar composition of Holocene soft bottom and coral associated mollusc faunas from the region. We intend to find data about faunas related to macroalgae in order to test whether seagrass faunas can be characterized on gross diversity and abundance of guilds.

¹⁾ NCB Naturalis, P.O. box 9517, 2300 RA Leiden, the Netherlands. e-mail: sonja.reich@ncbnaturalis.nl

already known decapod locality Cucao, on Chiloé Island, *Eriphia* sp. is added to the fauna. Specimens from Stokes Island are not sufficiently well determined yet. Chiloé and Stokes faunas are known to be of early Miocene age, while Mocha localities are supposed to be of an undetermined younger Miocene age. Specimens from Stokes Island constitute the southernmost record of Cenozoic Chilean decapod crabs. The biogeography of Miocene Chilean brachyurans will be reviewed and discussed.

¹⁾ Institut für Geowissenschaften, Christian-Albrechts-Universität zu Kiel, Ludewig-Meyn-Str. 10, 24118 Kiel, Germany; ¹ email: p.reuter2@gmx.de; ² email: nielsen@gpi.uni-kiel.de



Where are all the fossil pennatulacean octocorals? The need for more research on modern sea pens

Vanessa J. Roden¹⁾ & Michael Reich^{2,3)}

Sea pens, or Pennatulacea, are highly specialised, unbranched, colonial anthozoan octocorals with a worldwide marine distribution. There are estimated to be around 200 extant sea pen species. They are largely composed of soft tissue, the only hard parts being the axial rod and sclerites, which are not present in all modern sea pens, but are the only material that can be used to compare fossil with recent sea pens. Since fossil sea pen axes are inconspicuous and not very well known, they are frequently overlooked or misidentified. Due to the meagre fossil record, which begins in the Late Cretaceous, or perhaps earlier, and due to the lack of research that has been done on modern sea pens, not much is known on the evolution and phylogenetic relationships within this octocoral group. To make conclusions on pennatulacean systematics and evolution, 20 modern sea pen species (23 specimens in total) were studied, representing 10 of the 14 valid pennatulacean families. Using field emission scanning electron microscopy (and x-ray computed tomography in part), photographs were made of the cross-section and microstructure of the axial rods, which are important systematic characteristics, considering that soft-part morphology cannot be applied on fossil specimens.

The cross-section and microstructure changes along the axial rod and during the life of a sea pen, which can make fossil species determination difficult. With this study of the hard-part morphology of recent pennatulaceans, conclusions on fossil sea pen specimens and on the overall systematics of recent and fossil pennatulaceans can be made. In addition, by comparing recent and fossil axial

Miocene brachyuran crabs from southern Chilean islands

Philipp Reuter¹⁾ & Sven N. Nielsen²⁾

Only very few works exist about the decapod fauna of the Miocene of Chile. Here we present new records based on material from southern Chilean islands. The reported decapod genera all belong to the infraorder Brachyura and come from Mocha Island (38°S), Chiloé Island (42°S) and Stokes Island (45°S). Mocha Island localities yielded specimens of *Metacarcinus* sp., *Rochinia* sp., *Trichopeltarion* sp., and possibly a new genus; the former two being the first records of the respective genera for Chile. For the