



Modelling the background aerosol climatologies (1989-2010) for the Mediterranean basin

Pedro Jimenez-Guerrero and Sonia Jerez

Physics of the Earth, Universidad de Murcia, Spain (pedro.jimenezguerrero@um.es)

Aerosol levels and composition are influenced by multiple atmospheric physico-chemical processes that can affect them from its release point (as primary aerosol), or via gas-to-particle conversion processes that give rise to secondary aerosols. The contribution of the various aerosol sources, the role of long-range transport and the contribution of primary and secondary particulate matter to the ambient aerosol concentrations over Europe are not well known (Kulmala et al., 2009). Focusing on the Mediterranean, Querol et al. (2009) point out that there is a lack of studies on the variability of particulate matter (PM) along the Mediterranean basin, necessary for understanding the special features that differentiate aerosol processes between the western, eastern and central Mediterranean basins.

In this perspective, modelling systems based on state-of-science chemistry transport models (CTMs) are fundamental elements to investigate the transport and chemistry of pollutants behaviour at different scales and to assess the impact of emissions in aerosol levels and composition. Therefore, this study aims to summarise the results on the levels and chemical composition of aerosols along the Mediterranean basin, highlighting the marked gradient between the western-central-eastern coasts. Special attention is paid to the analysis of the seasonality of PM composition and levels. For this purpose, the regional modelling system WRF-CHIMERE-EMEP has been implemented for conducting a full transient simulation for the ERA-Interim period (1989-2010) using year-to-year changing EMEP emissions. The domain of study covers Europe with a horizontal resolution of 25 km and a vertical resolution of 23 layers in the troposphere; however the analysis focuses on the Mediterranean area. The PM levels and composition are compared to the measured values reported by the EMEP network, showing a good agreement with observations for both western and eastern Mediterranean.

The modelling results for the Mediterranean basin indicate that the aerosol levels follow a seasonal pattern with summer maximum concentrations caused by an increased secondary activity and the lower precipitation in the area, together with the contribution of Saharan dust outbreaks. The eastern Mediterranean is characterized by higher levels of sulphate, ammonium and OM+EC compared to the western part of the basin.

Nitrate presents much differentiated levels between winter (DJF) and summer (JJA) periods, with maxima during colder months in northern Italy and the Netherlands and lower levels in the warm season. This can be related to the thermal instability of the ammonium nitrate in summer ambient conditions, favouring the gas phase prevalence of nitrate. Sulphate levels at both parts of the basin increase progressively from April–May to reach maximum levels in mid-summer, due to enhanced photochemistry, low air mass renovation at regional scale, the increment of the summer mixing layer depth favouring the regional mixing of polluted air masses, and the possible higher summer contribution of marine secondary sulphate from DMS oxidation. The levels of sulphate are higher in the eastern Mediterranean and are highly correlated to ammonium levels. OM+EC concentrations are maxima in JJA in the whole basin, because of the higher formation of secondary organic aerosols (SOA) from different natural and anthropogenic sources. Secondary peaks are usually observed in spring (MAM) and autumn (SON) and are associated with winter anticyclonic pollution episodes. For sea-salt aerosols the modelling results indicate a non-uniform behaviour in the Mediterranean basin, showing a strong seasonality and gradient. A higher concentration of marine aerosol is estimated for the western Mediterranean (Alboran Sea and Gulf of Lion) during summertime, related to the increasing sea breeze circulation over the coast, which intensifies in the mid-summer. On the eastern Mediterranean, no large differences are found seasonally; here the sea spray clearly follows the wind speed variation.

The results confirm the capability of the modelling strategies to reproduce the particulate matter levels, composition and variation in the Mediterranean area. This kind of information is useful for establishing improve-

ment strategies for the prediction aerosols and to achieve the standards set in European Directives for modeling applications.

Kulmala, M., Asmi, A., Lappalainen, H.K., Carslaw, K.S., Pöschl, U., Baltensperger, U. Hov, O., Brenguier, J.-L., Pandis, S.N., Facchini, M.C., Hanson, H.-C., Wiedensohler, A., O'Dowd, C.D., 2009. Introduction: European Integrated Project on Aerosol Cloud Climate and Air Quality interactions (EUCAARI) – integrating aerosol research from nano to global scales. *Atmos. Chem. Phys.*, 9, 2825-2841.

Querol, X., Alastuey, A., Pey, J., Cusack, M., Pérez, N., Mihalopoulos, N., Theodosi, C., Gerasopoulos, E., Kubilay, N., Koçak, M., 2009. Variability in regional background aerosols within the Mediterranean. *Atmos. Chem. Phys.*, 9, 4575-4591.