



## **Observations of the interaction between near-inertial waves and mesoscale eddies**

Antonio Martínez-Marrero (1), Pablo Sangrá (1), Rui Caldeira (2), Borja Aguiar-González (3), and Ángel Rodríguez-Santana (1)

(1) University of Las Palmas de Gran Canaria, IOCAG, Las Palmas de Gran Canaria, Spain (antonio.martinez@ulpgc.es), (2) CIIMAR Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Portugal, (3) NIOZ Royal Netherlands Institute for Sea Research, Den Burg, Netherlands

Trajectories of eight drifters dragged below the surface mixed layer and current meter data from a mooring are used to analyse the interaction between near-inertial waves and mesoscale eddies. Drifters were deployed within eddies generated downstream of Canary and Madeira islands between 1998 and 2007. The mooring was installed in the passage of cyclonic eddies induced by Gran Canaria island during 2006. Rotatory wavelet analysis of Lagrangian velocities shows a clear relationship between the near-inertial waves' intrinsic frequencies and the eddy angular velocities. The results reveal that near-inertial waves reach a minimum frequency of half the planetary vorticity ( $f/2$ ) in the inner core of young anticyclonic eddies rotating with its maximum absolute angular speed of  $f/2$ . The highest amplitudes of the observed inertial motions are also found within anticyclonic eddies evidencing the trapping of inertial waves. Finally, the analysis of the current meter series show frequency fluctuations of the near-inertial currents in the upper 500 meters that are related to the passage of cyclonic eddies. These fluctuations appear to be consistent with the variation of the background vorticity produced by the eddies.