



Remote sensing of dust source characteristics and emission controls on a large playa

Robert Bryant (1), Frank Eckardt (2), Kathryn Vickery (2), Giles Wiggs (3), Martin Hipondoka (4), Jonathan Murray (5), Helen Brindley (5), James King (3), Jo Nield (6), David Thomas (3), and Richard Washington (3)

(1) University of Sheffield, Department of Geography, Sheffield, United Kingdom, (2) Department of Environmental & Geographical Science, University of Cape Town, South Africa, (3) School of Geography and the Environment, University of Oxford, Oxford, United Kingdom, (4) Department of Geography, History and Environmental Studies University of Namibia, Namibia, (5) Space and Atmospheric Physics, Imperial College, London, United Kingdom, (6) Geography and Environment, University of Southampton, Southampton, United Kingdom

Playas are ephemeral, endorheic lake systems that are common arid regions which have been identified as both regionally and globally significant sources of mineral dust. Emissions of dust from large playas can therefore impact significantly on regional climate through a range of land/atmosphere interactions. Nevertheless our understanding of spatial and temporal controls on dust emission from these environments at scales relevant to climate models, are poorly understood. Etosha Pan is a large playa situated in semi-arid northern Namibia. It sits in a basin 1000 m amsl, has a surface area of approximately 5000 km². This playa has been identified as one of the largest sources of mineral dust in the Southern Hemisphere. Here, to uncover sub-basin scale controls on mineral dust emissions, multiple time-series of remote sensing data (2004-2014) for this playa are presented. These depict: (a) mineral aerosol concentration in the vicinity of the playa, and (b) the surface hydrology of the playa basin. Seasonal wind velocities/direction data are also presented (ERA-Interim), and together with aerosol characterisation, they depict clear seasonality in dust emission/transport, significant inter-annual variability in aerosol loadings, and evidence of inter-event variability in plume chemistry. The gross spatial and temporal behaviour of the dust cycle of this ephemeral lake can be directly related to inherent variability in groundwater depth and periods of surface inundation. In most years, surface water interacts with < 30% of the playa surface; often for relatively short periods (< 3 months). However, Etosha Pan can also experience extensive periods of flooding and inundation (>80% basin cover, lasting for up to 12 months). Data from these wet and dry periods, in conjunction with locations of observed dust plumes emanating from the playa surface, allow us to identify direct links between dust emission and surface hydrology; thereby allowing, for the first time, a characterisation of the emission potential of the playa surface. These observations provide a clear context for both field investigation and modelling of this dust source as part of the DO4-Models (Dust Observations for Models) project that aims to understand the variability in dust emission processes at relevant scales for climate modelling.