



## **1.2 million years of aeolian activity in northwestern Western Australian recorded in a deep-sea core in the eastern Indian Ocean**

Felix Temmesfeld (1), Jan-Berend Stuut (1,2), Franck Bassinot (3), David Heslop (4), and Patrick De Deckker (4)  
(1) NIOZ - Royal Netherlands Institute for Sea Research, Marine Geology, Den Burg, Netherlands (jbstuut@nioz.nl — www.nioz.nl/dust), (2) MARUM - Center for Marine Environmental Sciences, University of Bremen, Germany, (3) LSCE (CEA/CNRS/UVSQ), Domaine du CNRS, Gif-sur-Yvette, France, (4) Research School of Earth Sciences, the Australian National University, Canberra, Australia

There is no continuous record of aridity from the Australian continent and we are presenting here, for the first time, a record of aeolian activity for northwestern Western Australia. Our data are based on a 32m long deep-sea core taken offshore North West Cape, the northwestern tip of Western Australia.

Our data rely on 2 adjacent studies: (a) a complete XRF scan of the core which provide elemental ratios which are translated into an aeolian component as well as a fluvial discharge-to-sea component. The first one relates to periods of aridity inland Australia, whereas the second one is interpreted as river discharge during periods of monsoonal activity; (b) a close examination of grain size analyses and observations of wind-blown quartz grains found in close to 600 samples taken from the core.

Our results clearly show a cyclic record of alternating dry and wet periods spanning the last 1.2 million years. Our findings also indicate that monsoonal activity as well as desertification were already in place in northern Australia so long ago, and this has clear implications for the evolution of the arid zone biota, associated fire activities and geomorphological features in northern Australia.