



## **Resilience landscapes for Congo basin rainforests vs. climate and management impacts**

Stephan Alexander Pietsch (1), Sishir Gautam (2), Johannes Elias Bednar (3), Patrick Stanzl (4), Aline Mosnier (1), and Michael Obersteiner (1)

(1) IIASA - International Institute of Applied Systems Analysis, 2361 Laxenburg, Austria (pietsch@iiasa.ac.at), (2) Tree Canada, (3) University of Vienna, (4) VMS Gabon

Past climate change caused severe disturbances of the Central African rainforest belt, with forest fragmentation and re-expansion due to drier and wetter climate conditions. Besides climate, human induced forest degradation affected biodiversity, structure and carbon storage of Congo basin rainforests. Information on climatically stable, mature rainforest, unaffected by human induced disturbances, provides means of assessing the impact of forest degradation and may serve as benchmarks of carbon carrying capacity over regions with similar site and climate conditions. BioGeoChemical (BGC) ecosystem models explicitly consider the impacts of site and climate conditions and may assess benchmark levels over regions devoid of undisturbed conditions.

We will present a BGC-model validation for the Western Congolian Lowland Rainforest (WCLRF) using field data from a recently confirmed forest refuge, show model – data comparisons for disturbed und undisturbed forests under different site and climate conditions as well as for sites with repeated assessment of biodiversity and standing biomass during recovery from intensive exploitation. We will present climatic thresholds for WCLRF stability, and construct resilience landscapes for current day conditions vs. climate and management impacts.