



## **The topography of the Iberian Peninsula from coupled geophysical-petrological inversion of multiple data sets**

Javier Fulla (1,2), Ana Negrodo (2,3), María Charco (2), Imma Palomeras (4), Antonio Villaseñor (5), and Juan Carlos Afonso (6)

(1) Dublin Institute for Advanced Studies, Geophysics section, Dublin, Ireland (jfulla@cp.dias.ie), (2) Institute of Geosciences IGEO (CSIC-UCM). Madrid, Spain, (3) Department of Geophysics and Meteorology, Complutense University of Madrid, Spain, (4) Universidad de Salamanca, Department of Geology Salamanca, Castilla y León, Spain, (5) Institute of Earth Sciences 'Jaume Almera', Barcelona, Spain, (6) Department of Earth and Planetary Sciences, Macquarie University, Sydney, Australia

In this study we have performed a 1D nonlinear Bayesian (probabilistic) inversion of a wide variety of data sets, extensively exploring the parameter space by means of a coupled geophysical-petrological inversion algorithm. The goal is to obtain a robust estimation of the thermal, compositional and density structure of the lithospheric/sublithospheric upper mantle system beneath the Iberian Peninsula, a crucial constraint to understand the complex geodynamic evolution in the study area. The most prominent feature in the modeled lithospheric structure is the progressive northward and northeastward steepening of the lithospheric-asthenospheric boundary (LAB) below the Ebro basin, reaching > 120 km under the central and western Pyrenees. Similarly, absolute maximum values of crustal thickness are obtained in the central Pyrenees, locally exceeding 45 km. Further to the west the Moho discontinuity shallows to about 35 km beneath the Cantabrian Cordillera. A dramatic decrease in both crustal and lithospheric thickness is observed from the central towards the easternmost Pyrenees, reaching depths of about 25 km and 90 km for the Moho and LAB respectively. Average Moho depth values of about 30 km are estimated in the central Iberian Peninsula. A slightly thicker crust is predicted under the Gibraltar arc than under the Betics, consistently with the deeper LAB beneath the former, most likely reflecting the presence of a sinking lithospheric slab. For the rest of the Iberian Peninsula a rather flat topography of LAB and Moho is observed, with moderate lithospheric thinning below the central western and SE Iberian margins. Isostatic topography related to variations in predicted crustal thickness shows local significant discrepancies from observed topography, thus indicating important regional contributions from dynamic and mantle source. The thermal and compositional fields in the lithospheric reveal the imprints of past and ongoing tectonic processes that have their expression in the present-day topography of Iberia.