

## Impacts of light-absorbing impurities on snow and their quantification with bidirectional reflectance measurements

Maria Gritsevich (1), Jouni Peltoniemi (2), Outi Meinander (3), Pavla Dagsson-Waldhauserová (4,5), Nataliya Zubko (2), Teemu Hakala (2), Aki Virkkula (3), Jonas Svensson (3), and Gerrit de Leeuw (3)

(1) Department of Physics, University of Helsinki, Finland (maria.gritsevich@helsinki.fi), (2) Finnish Geospatial Research Institute, Masala, Finland, (3) Finnish Meteorological Institute, Finland, (4) University of Iceland, Háskóli Íslands, Iceland, (5) Agricultural University of Iceland, Landbúnaðarháskólinn, Iceland

In order to quantify the effects of absorbing impurities on snow and define their contribution to the climate change, we have conducted a series of dedicated bidirectional reflectance measurements. Chimney soot, volcanic sand, and glaciogenic silt have been deposited on the snow in the controlled way. The bidirectional reflectance factors of these targets and untouched snow have been measured using the Finnish Geodetic Institute's field goniospectrometer FIGIFIGO, see, e.g., [1, 2] and references therein. It has been found that the contaminants darken the snow, and modify its appearance mostly as expected, with clear directional signal and modest spectral signal. A remarkable feature is the fact that any absorbing contaminant on snow enhances the metamorphosis under strong sunlight [3, 4]. Immediately after deposition, the contaminated snow surface appears darker than the pure snow in all viewing directions, but the heated soot particles start sinking down deeply into the snow in minutes. The nadir measurement remains darkest, but at larger zenith angles the surface of the soot-contaminated snow changes back to almost as white as clean snow. Thus, for on ground observer the darkening by impurities can be completely invisible, overestimating the albedo, but a nadir looking satellite sees the darkest points, now underestimating the albedo.

After more time, also the nadir view brightens, and the remaining impurities may be biased towards more shadowed locations or less absorbing orientations by natural selection. This suggests that at noon the albedo should be lower than in the morning or afternoon. When sunlight stimulates more sinking than melting, albedo should be higher in the afternoon than in the morning, and vice versa when melting is dominating. Thus to estimate the effects caused by black carbon (BC) deposited on snow on climate changes may one need to take into account possible rapid diffusion of the BC inside the snow from its surface. When the snow melt rate gets faster than the diffusion rate (under condition of warm outside temperatures), as it was observed at the end of the experiment reported here, dark material starts accumulating into the surface [5]. The BC deposited on snow at warm temperatures initiates rapid melting process and may cause dramatic changes on the snow surface.

### References

- 1 Peltoniemi J.I., Hakala T., Suomalainen J., Honkavaara E., Markelin L., Gritsevich M., Eskelinen J., Jaanson P., Ikonen E. (2014): Technical notes: A detailed study for the provision of measurement uncertainty and traceability for goniospectrometers. *Journal of Quantitative Spectroscopy & Radiative Transfer* 146, 376-390, <http://dx.doi.org/10.1016/j.jqsrt.2014.04.011>
- 2 Zubko N., Gritsevich M., Zubko E., Hakala T., Peltoniemi J.I. (2016): Optical measurements of chemically heterogeneous particulate surfaces // *Journal of Quantitative Spectroscopy and Radiative Transfer*, 178, 422-431, <http://dx.doi.org/10.1016/j.jqsrt.2015.12.010>
- 3 Peltoniemi J.I., Gritsevich M., Hakala T., Dagsson-Waldhauserová P., Arnalds Ó., Anttila K., Hannula H.-R., Kivekäs N., Lihavainen H., Meinander O., Svensson J., Virkkula A., de Leeuw G. (2015): Soot on snow experiment: bidirectional reflectance factor measurements of contaminated snow // *The Cryosphere*, 9, 2323-2337, <http://dx.doi.org/10.5194/tc-9-2323-2015>
- 4 Svensson J., Virkkula A., Meinander O., Kivekäs N., Hannula H.-R., Järvinen O., Peltoniemi J.I., Gritsevich M., Heikkilä A., Kontu A., Neitola K., Brus D., Dagsson-Waldhauserová P., Anttila K., Vehkamäki M., Hienola A., de Leeuw G. & Lihavainen H. (2016): Soot-doped natural snow and its albedo — results from field experiments. *Boreal Environment Research*, 21, 481-503, <http://www.borenav.net/BER/pdfs/preprints/Svensson1498.pdf>
- 5 Meinander O., Kontu A., Virkkula A., Arola A., Backman L., Dagsson-Waldhauserová P., Järvinen O., Manninen T., Svensson J., de Leeuw G., and Leppäranta M. (2014): Brief communication: Light-absorbing impurities can reduce the density of melting snow, *The Cryosphere*, 8, 991-995, doi:10.5194/tc-8-991-2014.