



## **Intermittent energy dissipation by turbulent reconnection**

Huishan Fu (1), Andris Vaivads (2), Yuri Khotyaintsev (2), Mats André (2), Jinbin Cao (1), Vyacheslav Olshevsky (3), Jonathan Eastwood (4), and Alessandro Retino (5)

(1) School of Space and Environment, Beihang University, Beijing, China, (2) Swedish Institute of Space Physics, Box 537, SE-751 21 Uppsala, Sweden, (3) Center for mathematical Plasma Astrophysics, KU Leuven, Leuven, Belgium, (4) The Blackett Laboratory, Imperial College London, London SW7 2AZ, United Kingdom, (5) Laboratoire de Physique des Plasmas, CNRS/Ecole Polytechnique/UPMC, Palaiseau, France

Magnetic reconnection—the process responsible for many explosive phenomena in both nature and the laboratory—is efficient at dissipating magnetic energy into particle energy. To date, exactly how this dissipation happens remains unclear, owing to the scarcity of multi-point measurements of the ‘diffusion region’ at the sub-ion scale. Here we report such a measurement by Cluster—four spacecraft with separation of  $1/5$  ion scale. We discover numerous current filaments and magnetic nulls inside the diffusion region of magnetic reconnection, with the strongest currents appearing at spiral nulls (O-lines) and the separatrices. Inside each current filament, kinetic-scale turbulence is significantly increased, and the energy dissipation,  $E' \cdot J$ , is 100 times larger than the typical value. At the jet reversal point, where radial nulls (X-lines) are detected, the current, turbulence, and energy dissipations are surprisingly small. All these features clearly demonstrate that energy dissipation in magnetic reconnection occurs at O-lines but not X-lines.