

## **Hennersdorf section (Lake Pannon, Upper Miocene): use of stable isotopes on ostracods for a palaeoenvironmental reconstruction**

Marco CAPORALETTI, Martin GROSS & Werner E. PILLER

Lake Pannon is an ancient lake-system which developed after the vanishing of the Paratethys Sea. Its history starts about 11.6 Ma ago in correspondence to the Sarmatian/Pannonian boundary. During the initial phase the lake was characterized by a brackish and alkaline water body. The following progressive freshening of Lake Pannon's water throughout the Pannonian lead to a high endemism among the benthic fauna (HARZHAUSER et al. 2007).

The purpose of this study is to characterize the palaeoenvironmental changes, which occurred during the early Pannonian in Lake Pannon. In order to pursue this intent oxygen ( $\delta^{18}\text{O}$ ) and carbon ( $\delta^{13}\text{C}$ ) stable isotope analyses on ostracods and bulk sediments were made.

Ostracods and sediments analysed for this study come from the Hennersdorf section; one of the best-known localities of Lake Pannon's marginal area. The Hennersdorf clay pit is located about 10 km south of Vienna (Austria) and is correlated with the lower part of the Pannonian zone E (~10.5 Ma). The outcrop is made up of by about 15 meters of clays differentiated mainly by colour. The lower part is characterized by the presence of several coquina levels, whereas the upper portion is almost barren of molluscs (HARZHAUSER & MANDIC 2004; HARZHAUSER et al. 2008; DANIELOPOL et al. 2011). Ostracod specimens are present in almost the entire section.

A total of 44 samples coming from Hennersdorf section were studied. For each sample up to 5 ostracod species were analysed: *Cyprideis obesa*, *Hemicytheria* sp., *Loxoconcha granifera*, *Loxoconcha* ex gr. *rhomboidea*, *Leptocythere* sp. Only *Cyprideis obesa* is present throughout the section. For the stable isotope analyses the ostracods were carefully cleaned and stored in ethanol; no chemical pre-treatments were used. Ostracod analyses were coupled with sediment analyses (stable isotopes and sieve analysis).

Despite that the use of stable isotope analyses on ostracods is quite widespread, Lake Pannon's results do not show a unequivocally interpretable environmental pattern. The interplay of the different factors (salinity, temperature, vital effects, water influxes, etc.) and the peculiar characteristics of the lake do not permit, probably, to define the palaeoenvironment of the area only through the stable isotopes.

### Acknowledgements

This work is supported by the Austrian Science Fund (P21748-N21 to M.G. and P17738-B03 to Dan L. DANIELOPOL). We further like to express our gratitude to Mathias HARZHAUSER (NHM Wien) and Klaus MINATI (KFU Graz).

### References

- DANIELOPOL, D.L., GROSS, M., HARZHAUSER, M., MINATI, K. & PILLER, W.E. (2011): How and why to achieve greater objectivity in taxonomy, exemplified by a fossil ostracod (*Amplocypris abscissa*) from the Miocene Lake Pannon. – *Joannea Geologie und Paläontologie*, 11: 273-326, Graz.
- HARZHAUSER, M. & MANDIC, O. (2004): The muddy bottom of Lake Pannon – a challenge for dreissenid settlement (Late Miocene; Bivalvia). – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 204: 331-352, Amsterdam.
- HARZHAUSER, M., LATAL, C. & PILLER, W.E. (2007): The stable isotope archive of Lake Pannon as a mirror of Late Miocene climate change. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 249: 335-350, Amsterdam.
- HARZHAUSER, M., KERN, A., SOLIMAN, A., MINATI, K., PILLER, W.E., DANIELOPOL, D.L. & ZUSCHIN, M. (2008): Centennial- to decadal-scale environmental shifts in and around Lake Pannon related to a major Late Miocene lake-level rise. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 270(1-2): 102-115, Amsterdam.

Authors addresses:

Marco Caporaletti & Werner E. Piller  
University of Graz, Institute of Earth Sciences, Heinrichstrasse 26, A-8010 Graz,  
Austria  
marco.caporaletti@uni-graz.at

Martin Gross

Universalmuseum Joanneum, Department for Geology & Palaeontology, Weinzöttl-  
strasse 16, A-8045 Graz, Austria