

## **Ostracod preservation and response to Late Glacial and Early Holocene climate changes in a sub-alpine belt lake of the southern Romanian Carpathians**

Sanda IEPURE, Tadeusz NAMIOŃKO & Enikő K. MAGYARI

The role of ostracods in environmental and climate change reconstructions of Late Pleistocene and Holocene lacustrine environments of the Alpine region is well established (e.g., LÖFFLER 1983; VON GRAFENSTEIN 1999; LAUTERBACH et al. 2011). In contrast, ostracod studies in glacial lakes of the (sub-)alpine belt of the Carpathians in Romania thus far have been completely neglected, although the effects of the last glaciation have been more significant there than elsewhere in the southern Carpathians (REUTHER et al. 2007). The Retezat Mountains in the Romanian Southern Carpathians are home of several peaks of over 2000 m asl that represented the major areas of ice accumulation during the Pleistocene (REUTHER et al. 2007). Due to prevailing Mediterranean circulation of moist air masses this mountain unit is the wettest (1400 mm/yr at 1500–1600 m asl) in the Southern Carpathians (MINDRESCU et al. 2010). Recently, a series of Late Quaternary palaeoecological studies on the glacial lake sediments from the Carpathians has been initiated in order to provide regional palaeoclimatic and palaeoecological records inferred from multiple proxies (plant macrofossils, pollen, siliceous algae, cladocerans, chironomids as well as geochemical and ancient DNA analyses) (MAGYARI et al. 2009, 2011, KORPONAI et al. in press). The present study discusses the preservation of fossil ostracod assemblages and their response to Late Glacial and Early Holocene abrupt climate changes. The investigations were carried out on the lowermost 90 cm section of a 5 m-long core drilled in the sediments of Taul dintre Brazi Lake. This lake is located in the northern part of the Retezat Mountains at 1740 m asl in the sub-alpine belt. Due to its very small surface (about 30 m<sup>2</sup>) and clearly defined catchment area, the lake is regarded as a very sensitively and rapidly responding ecosystem.

The studied core sediment sequence, corresponding to the period between 15.2 and 10.0 cal kyr BP, when an open mixed conifer forest formed around the lake (MAGYARI et al. 2011), yielded a total of 12,273 ostracod valves. However, the ostracod abundances of individual (2 cm-thick) sediment samples differed, varying between 0 (at the base of the sequence and around 14.8 cal kyr BP) and 975 (around 11.8 cal kyr BP), with an average of 342 valves per 10 cm<sup>3</sup> of fresh sediment. Although the overall ostracod abundances were high, the recovered valves were heavily decalcified,

lacking their calcitic part, while still retaining strong chitinous cuticle. Interestingly enough, remains of chitinous exoskeleton of the limbs as well as those of the Zenker organ and even spermatozoa were found well preserved almost throughout the entire studied sequence. The ostracod fossil assemblage of the lake proved to be monospecific, consisting of only *Cypria ophthalmica* (JURINE). The species is nearly cosmopolitan generalist, inhabiting almost any type of inland waters, often doing well in stressed environments, as those highly organically polluted, hypoxic or acidic (MEISCH 2000). It appears that *C. ophthalmica* colonized the lake around 14.3 cal kyr BP, i.e., after the period (15.7–14.5 cal. kyr BP) when low organic content and a shallow and oligotrophic environment prevailed (indicated by a poor assemblage of Cladocera: KORPONAI et al., in press).

*Cypria ophthalmica* showed markedly high fluctuations reflecting abrupt climate changes throughout the Late Pleistocene and the onset of the Holocene associated with the changes in the trophic level of the lake. Overall, the species abundance increased during significant episodes of warming (the first peak was observed in the GI-1c interstadial and the second one at the end of the Younger Dryas YD) and decreased when cooling conditions occurred (during YD) as indicated by the Greenland oxygen isotope record. At around 13.3 cal kyr BP, the abundance of *C. ophthalmica* and cladocerans *Chydorus sphaericus* (O.F. MÜLL.) and *Daphnia longispina* (O.F. MÜLL.) (KORPONAI et al. in press) shows a short rapid increase and reaches a maximum at 13.0 cal kyr BP. The decrease in *C. ophthalmica* abundance was associated with the organic content depletion in the sediment that could be further linked with the re-advance of the M3 glacier in the northern part of the Retezat Mountains and the associated cooling effect. During the YD, the abrupt increase in seasonality caused an evident decrease of *C. ophthalmica*, but the species abundance was maintained as moderate until the onset of the Holocene. However, throughout the entire YD reversal, the lake productivity did not drastically decrease, as evidenced by the sediment stratigraphy and loss of ignition measurements (MAGYARI et al. 2010). From the beginning of the Holocene population of *C. ophthalmica* decreased significantly until 10.1 cal kyr BP when reconstructed temperature was 13.2°C, higher than at the present day in July at this elevation (MAGYARI et al. 2010). This work is the first contribution on Late Glacial–Holocene ostracods of the (sub-)alpine lacustrine habitats in the Romanian Carpathians.

---

### Acknowledgements

This work was possible due to the financial support of the Sectorial Operational Program for Human Resources Development 2007-2013, co-financed by the European Social Fund, under the project number POSDRU 89/1.5/S/60189 with the title „Post-doctoral Programs for Sustainable Development in a Knowledge Based Society” and is also a contribution to the PROLONGE project sponsored by OTKA PD73234 and MERG-CT-2006-041088: Combining Palaeoecology and Palaeogenetics.

## References

- KORPONAI, J., MAGYARI, E.K., BUCZKÓ, K., IEPURE, S., NAMIOTKO, T., CZAKÓ, D., KÖVÉR, C. & BRAUN, M. (in press): Cladocera-response to Lateglacial to early Holocene climate change in a South Carpathian mountain lake. – *Hydrobiologia*, Dordrecht.
- LAUTERBACH, S., BRAUER, A., ANDERSEN, N., DANIELOPOL, D.L., DULSKI, P., HÜLS, M., MILECKA, K., NAMIOTKO, T., OBREMSKA, M. & VON GRAFENSTEIN, U. (2011): Environmental responses to Lateglacial climatic fluctuations recorded in the sediments of pre-Alpine Lake Mondsee (north-eastern Alps). – *Journal of Quaternary Science*, 26 (3): 253-267, Chichester.
- LÖFFLER, H. (1983): Aspects of the history and evolution of alpine lakes in Austria. – *Hydrobiologia*, 100: 143-152, Dordrecht.
- MAGYARI, E.K., BRAUN, M., BUCZKÓ, K., KERN, Z., LÁSZLÓ, P., HUBAY, K. & BÁLINT, M. (2009): Radiocarbon chronology of glacial lake sediments in the Retezat Mts (S Carpathians, Romania): a window to Lateglacial and Holocene climatic and palaeoenvironmental changes. – *Central European Geology*, 52: 225-248, Budapest.
- MAGYARI E.K., TÓTH, M., JAKAB G., BUCZKÓ K., BRAUN, M. & BÁLINT, M. (2010): Vegetation response to rapid climate changes during the lateglacial: a case study from the South Carpathians. – [pages-142.unibe.ch/about/national/hungary/Magyari-Toth.pdf](http://pages-142.unibe.ch/about/national/hungary/Magyari-Toth.pdf).
- MAGYARI, E.K., MAJOR, Á., BÁLINT, M., NÉDLI, J., BRAUN, M., RÁCZ, I. & PARDUCCI, L. (2011): Population dynamics and genetic changes of *Picea abies* in the South Carpathians revealed by pollen and ancient DNA analyses. – *Evolutionary Biology*, 11: 66, New York.
- MEISCH, C. (2000): Freshwater Ostracoda of Western and Central Europe. – In: SCHWOERBEL, J. & ZWICK, P. (Hrsg.): Süßwasserfauna von Mitteleuropa. – 8(3): 1-522, Spektrum Akademischer Verlag, Heidelberg-Berlin.
- MÍNDRESCU M., EVANS I.S. & COX, N.J. (2010): Climatic implications of cirque distribution in the Romanian Carpathians: palaeowind directions during glacial periods. – *Journal of Quaternary Science*, 25(6): 875-888, Chichester.
- REUTHER, A.U., URDEA, P., GEIGER, C., IVY-OCHS, C., NILLER, H.P., KUBIK, P.W. & HEINE, K. (2007): Late Pleistocene glacial chronology of the Pietrele Valley, Retezat Mountains, Southern Carpathians constrained by <sup>10</sup>Be exposure ages and pedological investigations. – *Quaternary International*, 164/165: 151-169, Oxford.
- VON GRAFENSTEIN, U., ERLLENKEUSER, H., BRAUER, A., JOUZEL, J. & JOHNSEN, S.J. (1999): A mid-European decadal isotope-climate record from 15,500 to 5000 years B.P. – *Science*, 284: 1654-1657, Washington DC.

### Authors addresses:

Sanda Iepure

Babes-Bolyai University, Department of Mineralogy, Kogalniceanu 1, 3400, Cluj, and Speleological Institute "Emil Racovitza", Department of Cluj (Biospeleology), Clinicilor 5, 400006 Cluj Napoca, Romania  
siepure@hasdeu.ubbcluj.ro

Tadeusz Namiotko  
University of Gdansk, Department of Genetics, Laboratory of Limnozoology, Kladki 24,  
80-822 Gdansk, Poland  
namiotko@biotech.ug.gda.pl

Enikő K. Magyari  
MTA-MTM Research Group for Paleontology, 1476 Budapest, P.O. Box 222, Hungary  
magyari@bot.nhmus.hu