

took place within the Alpine Wilson cycle. (1) Rifting processes of some 40-50 Ma duration which ended in the Late Triassic. (2) Opening of the Dinaridic Tethys which took place in Late Triassic/Early Jurassic time when a spreading center was set up. This made possible the generation of the oceanic crust during the period of 60-70 Ma. At that time there was probably also subsidence of the northeastern marginal parts of the CP composed of the Triassic formations with unconformably underlying Paleozoic formations, which were thus included in the basinal parts. (3) Subduction processes started in a subsequent Late Jurassic/Early Cretaceous time as indicated by the first emplacement of ophiolites. By the end of these processes, the Paleozoic-Triassic formations overlain by the oceanic crust were probably detached and thrust onto the emplaced ophiolites and their country rocks. The emplacement of ophiolites and stacking of the Paleozoic-Triassic nappes was accompanied by the first strong Alpine metamorphism (120-110 Ma). (4) In the northern Dinarides, in the Late Cretaceous a magmatic arc has been already generated as indicated by the presence of trench sediments and igneous rocks characteristic for such a geotectonic setting. (5) The main deformational (compressional) event (about 40-50 Ma) and medium-pressure metamorphism accompanied by synkinematic granite plutonism took place after the Eocene termination of subduction processes. This deformation produced main NW-SE-trending fold, thrust and imbricate structures with the southwestern vergences. Only north of the presumed subduction zone, opposing northeastern vergences due to obduction were recognized. (6) Post-orogenic evolution started after the Eocene uplift of the Dinarides which gave rise to the separation of the Tethys into the Mediterranean and the Paratethys.

### **Which is the time of rotation? Review of paleomagnetic and K-Ar data from Romania**

Cristian Panaiotu<sup>1</sup>, Zoltan Pecskay<sup>2</sup> and Cristina Panaiotu<sup>1</sup>

<sup>1</sup> Faculty of Physics, University of Bucharest, Bucharest, Romania

<sup>2</sup> Institute of Nuclear research, Hungarian Academy of Sciences, Debrecen, Hungary

Recent K-Ar data from Romania has changed or improved the age of some magmatic rocks sampled for paleomagnetic studies. A first important change concerns the age of basaltic andesites from Mures Valley (Apuseni and Poiana Rusca Mountains).

These rocks, previous considered as Paleogene, have ages between 66 ma and 72 Ma. The second important contribution of the radiometric data concerns the ages of the two groups with different paleomagnetic directions from Miocene magmatic rocks of Apuseni Mountains. Two K-Ar data from the group with a declination around 70° suggest an age between 14.7 - 12.4 Ma. The K-Ar data from the group that show no rotation suggest an age around 11 Ma. The paleomagnetic data support the existence of a domain characterized by a large clockwise rotation in the eastern part of the Carpatho-Pannonian area. The amplitude of the rotation is variable from 70° in the Apuseni Mountains -Banat area to 120° in the Bucegi Mountains. Data from the Apuseni Mountains suggest a very fast rotation during Sarmatian. This rotation was coeval with the counterclockwise rotation of the Gutai Mountains, but took place after the end of the counterclockwise rotation of the North Pannonian Paleogene basin (around 16 Ma). This fast rotation was accommodated in the brittle layer by coeval trusts and strike-slip faults in the East and South Carpathians and extensional grabens and shear zones in the Great Hungarian Plain. These rotations reflect probably the continuously deforming lithosphere beneath the seimogenic upper crusts. K-Ar data from Pannonian-Quaternary volcanic rocks of East Carpathians show the migration of the volcanism along the arc and a short duration of volcanic activity in individual segments. Paleomagnetic rotations are absent in these magmatic rocks, but the migration is in the same sense as the previous clockwise rotation. The above features are all consistent with the slab breakoff model. Cinematic parameters derive from these data will be discussed with respect to the proposed tectonic models for this area.

### **Styles of Miocene thrusting, strike-slip faulting and extension in the eastern Calcareous Alps**

Herwig Peresson and Kurt Decker

Institute of Geology, University Vienna, Austria

The final stage of collisional shortening in the Eastern Alps was characterized by intense brittle deformation of the Calcareous Alps during the Miocene. Thrusting over the European margin occurred until the Early Miocene (Karpatian stage, 17 Ma) as dated by overthrust Molasse sediments. Thrusting was generally directed northwards and exceeded 34 km since the Late Oligocene (Wessely, 1987). The irregular morphology of the overthrust European basement controlled deformation styles in the upper plate. From the