

of space north of these elements they very quickly entered into collision with the collapsing European active margin during Late Carboniferous to Early Permian. Continuing translation of Gondwana westward provided some new space and a new rift appeared north of Tunisia-Sicily (the Sicilian-Lagonegro rift system connected to the Ionian sea-East Mediterranean basin). The western part of the Meliata back-arc rift (Southern Alps-Ivrea) aborted at that time, whilst its eastern part (Hallstatt-Meliata-Dobrogea) oceanised (e.g. Early Triassic MORB pillow lava of N-Dobrogea). During the Late Permian-Early Triassic collision of the detached inner Dinaro-Hellenides Variscan sliver with the Cimmerian elements.

Characteristics of the lithosphere reflected upon electromagnetic parameters along the magneto-telluric profiles in Romania

Maria Stanica

Geological Institute of Romania, Bucharest

Information carried out along the eight MT profiles crossing the Eastern Carpathians and Pannonian Basin - Romanian area are used to reveal the most important aspects related to the lithospheric structure insisting on the suture zone associated to the Carpathian Orogen and Tethyan one. Taking into account the magnetotelluric (MT) results referring to the lithospheric thickness variations, contact alignments and structural relations between the main tectonic units and placement of the enhanced conductivity zones (over 0.1S/m) an explanation of the link between the lithosphere dynamics and crustal deformation is to be attempted. By means of the geoelectrical cross-section elaborated for every profiles, a general spatial image was pursued to be achieved. The MT profiles bring into the light some characteristic aspects emphasizing the structural particularities of these two major tectonic units. A first problem is related to the thickness of the sedimentary cover (1000 - 3000 m in Pannonian Basin) and (13000 - 18000 m in the Eastern Carpathians's Foredeep). The transition zone from the crust to mantle, marked by a decrease of the resistivity (less than 10 ohm.m) and associated especially to Orogenic areas stands for the interesting point to be clarified, as regarding its affiliation on the hand, by means of the electrical properties and of the assessed depth, and the causes which generated it, on the other hand. Obviously, owing to its involvement in revealing certain lithospheric aspects, the top of the asthenosphere (contact between tectonic plates of

various thickness) was pointed out where it was possible.

East Carpathian-Pannonian connection according to geophysical data

Vitali I. Starostenko and O.M. Kharitonov

Institute of Geophysics, Kiev, Ukraine

The geologic boundary between the East Carpathians and Pannonian Basin is eophysical data to be a part of the transcontinental Oder-Caucasian lineament separating the thick from thin lithosphere or hot from cold asthenosphere in Europe. Over this fault crustal thickness varies from 27-30 km in Pannonia to 56 km in East Carpathians. This boundary is very clearly seen in the geothermal field. The main thermal energy source is situated within Pannonia west of the Transcarpathian deep fault. The former caused lithosphere thinning and high-temperature asthenosphere formation under Pannonia at 60-65 km depth. Under the East Carpathians temperature decreases strikingly and the lithosphere thickens. The velocity, density and magnetic properties change. Here under the East Carpathians, the thin crust of the Pannonian basin is alternated by the thick lithospheric armour of the more than 65 km thick Cis-Carpathian through.

A rheological analysis of geophysical data enables us to express the following idea of the structure of this region. Under Pannonian basin the crust is thinned by high temperature to 28 km and thickens towards the Fold Carpathians. The sediments reach in the Transcarpathian through 5 km and rest on a young folded basement lying on plastic granites, which is demonstrated by high temperature (200° C) corresponding with granitoid creep. The plastic granitoids are underlain by 10-12 km thick brittle basites layer whose top is clearly traced by seismic data. The brittle-to-plastic basalts transition is not always a contrasting acoustic boundary and nearly coincides with the 520° C geotherm. The M-discontinuity separates in Pannonia basic from ultrabasic rocks which are in plastic state. The lithosphere thins to 60 km and has been destructed by high (to 1200° C) temperature.

In the East Carpathians bounded by deep faults crust thickens due to its doubling in granitoid and brittle basite layers. The M-discontinuity deepens even more in the Cis-Carpathian foredeep. This is due to sedimentary cover thickening and the appearance of a "crust-mantle mix" layer with 7.5-7.6 km/s longitudinal wave velocity. The large "crust-mantle mix" thickness is due to plunging of a layer of brittle basites ("basaltic plate") driven by