

The use of TM Landsat 5 images satellites (compositions 357, 457 and 375 with histogram equalization stretching enhancements) prove to be a very useful tool for examine the spatial distribution pattern of incipient meandering fluvial rivers.

Among them, two courses, the Pilcomayo and the Bermejo rivers, run about to the "Chaco-Pampeana" plain as typical "meandering" river model. This is a reflection about the flat relief and topography of the region and is related with the weather conditions (a very wet climate).

The sedimentary deposits of these rivers (the Pilcomayo river, the Bermejo river, tributary stream and minor courses) have not a great significance. In fact, these are almost non-existent and mainly they are re-worked sediments of the Upper Tertiary arenaceous sediments. So, they have very little erosional potential.

The deposits of these little rivers are mainly arenaceous lithofacies of fine to very fine sand grains. Grouped planar crossbeds, trough crossbeds stratification is frequent at the beds, in which sometimes, bipolar cross stratification appears. Solitary and grouped channels are common. The downstream accreted mesoform and macroform deposits aren't present and not recognized by now.

It is proposed here that the origin of these low volume courses of meandering design reorganization drainage are quite synchronous with the base sea level change occurred on the 3,000 and/or 6,000 years BP periods according with HAQ et al. eustatic curves (1998). The climatic change, varying from more dry toward more wet conditions at northeastern Argentina in the period mentioned above (TORRA 1998).

These age were accuracy tested upon ^{14}C dating over widely mollusks biofacies related with restricted recent marine transgression whose deposits are present at the entrance of the Río de la Plata estuary and adjoining areas as the coast-line of the Buenos Aires Province (CORTELEZZI & LERMANN 1969).

These brief reflections have the aims to bring forward some ideas about a theme absolutely obscured and never studied in the Chaco-Pampeana region with modern architectural and remote sensing techniques.

CORTELEZZI, C.R. y LERMANN, J.C. (1969): Estudio de las Formaciones Marinas de la Costa Atlántica de la Provincia de Buenos Aires. - VIII Congreso Mundial del INQUA, Paris, 135-164, LEMIT reprint, La Plata.

HAQ, B.U. & v. EYSINGA, F.W.B. (1998): Geological Time Table. - Fifth revised and enlarged edition. (Elsevier Science B.V.) Amsterdam.

TORRA, R. (1998): El Estudio de los Sedimentos del Cuaternario de la Provincia del Chaco y Regiones Aledañas. - III Reunión de Comunicaciones Científicas y Tecnológicas, Universidad Nacional del Nordeste, Actas, 4: 197-200, Resistencia.

Can geochemistry help in stratigraphic problems? The case of the Ituzaingó Formation (Middle Miocene), Northeastern Argentina

TORRA, R.

Güemes 749, 3500, Resistencia, Chaco, Argentina,
roberto_torra@arnet.com.ar

The Ituzaingó Formation sands are a very typical heterolithic succession of sandy-muddy beds that cover all the northeastern of Argentina and the southwest of the Paraguay Republic. The outcrops are of about 117,000 km² (TORRA 1999a, b).

However, recent study of texture in hidrologic drill-holes in the vast area, reveals that these sediments are quite common in the major portions of the 'Chaco-Paraná intracratonic basin'. The extension confirmed until the moment sound of somewhat 450,000 km² (TORRA 1999a).

The sandy lithofacies are characterized for a well sort, very fine grain, very well rounded and gray to white in shades. When reddish, they are ferricretized (epigenetic oxidation of magnetite and ilmenite).

Recent architectural studies over the sandy lithofacies, show a typical assemblage of internal structures that are diagnostics of a tide-dominated depositional environment.

Hummocky cross stratification, herringbone cross stratification, first and second order reactivation surfaces, tidal rhythmic beddings, flaser beddings, bipolar cross stratification, sigmoidal stratification and tidal bundles-beddings are present in all the outcrops (TORRA 2000a).

Typically, an erosion unconformity was proposed between the heterolithic succession and the mudrocks that overlies itself (HERBST 1971).

The trace elements study led to the interpretation that the mud beds that structurally cover in quite conformable arrange the heterolithic succession, has the same geochemical anomalies -or signatures- (TORRA 1999b, 2000b).

The overlying unit was generally interpreted as in erosion discordance (HERBST 1971) in despite of to not accept a simple lithofacies change.

Geochemical trace elements data analysis shows the same anomalies ('signatures') in the muddy lithofacies interleaving with very fine sands. All the mud beds analyzed had the same geochemistry anomalies. As a result, we interpreted them as formed under similar sedimentologic and environmental conditions.

Trace element proves to be a suitable tool in order to help to solve stratigraphic problems always together with architectural analysis at least in the case of the marine Miocene sequence of the Northeastern region of Argentina (TORRA 2000b).

HERBST, R. (1971): Esquema estratigráfico de la Provincia de Corrientes, República Argentina. - Revista Asociación Geológica Argentina, 24 (2): 221-243.

TORRA, R. (1999a): Avances en el conocimiento del Terciario Superior de la Llanura Chaco-Pampeana. - IV Reunión de Comunicaciones Científicas y Tecnológicas. Universidad Nacional del Nordeste, Actas. VI: 93-96, Corrientes.

TORRA, R. (1999b): Ituzaingó Formation. A Key for the Interpretation of the Upper Tertiary Stratigraphy, Mesopotamia-Chaco Paraná Basin, Argentina. - Revista Ciência e Natura. Universidade Federal de Santa Maria. 'in press'.

TORRA, R. (2000a): Reinterpretación Sedimentológica y Edad de las Formaciones Paraná, Ituzaingó, Toropí, Yupó y Puelches (Mioceno Medio), Mesopotamia de Argentina. - Revista Brasileira de Geociencias, Brasília. 'in press'.

TORRA, R. (2000b): Geochemistry of a Tidal Transgressive Heterolithic Succession: The Ituzaingó Formation (Middle Miocene), Argentina. - Chinese Journal of Geochemistry, 19 (1): 1-7.

Geochemical properties and origin of rich-iron manganese concretions in the Ituzaingó Formation (Middle Miocene), Northeastern Argentina

TORRA, R.

Güemes 749, 3500, Resistencia, Chaco, Argentina,
roberto_torra@arnet.com.ar

In the upper beds of the Ituzaingó Formation sands are present outstanding rich-iron manganese concretions previously studied by CORTELEZZI et al. (1997). The Ituzaingó Formation is largely defined by a heterolithic succession composed for interleaved sandy-muddy lithofacies. The upper levels are characterized for a typical thin mudrock bed, 2-4 meters in thickness. In these beds, horizontally disposed, breccial rich-iron manganese concretions appear. They are about 10-40 cm in size (average). Predominant shape is coarser botryoidal or irregular. The main elongation axis is markedly in the sense of the stratification. That is horizontally arranged.

Two selected samples of the Empedrado village and the Arroyo El Sombrero river, were picked up from very well exposed outcrops. The selected samples were analyzed by the technique of