

Where has the Tethyan lithosphere gone after consumption by the Alpine-Carpathian orogeny?

HETÉNYI, G.^{1,2,3}, STUART, G.W.¹, HOUSEMAN, G.A.¹ & the CBP WORKING GROUP⁴

¹School of Earth and Environment, University of Leeds, LS2 9JT, Leeds, UK; ²Laboratoire de Géologie, Ecole Normale Supérieure, CNRS-UMR 8538, Paris, France; ³Department of Geophysics, Eötvös Loránd University, Budapest, Hungary; ⁴E. HEGEDÜS, E. BRÜCKL, S. RADOVANOVIC, A. BRISBOURNE, P. LORINCZI, B. DANDO, G. FALUS, A. KOVÁCS, H. HAUSMANN

In the general scenario of plate tectonics, the outer rigid shell of the Earth is consumed along convergent plate boundaries. This consumption translates into downward movement of the lithosphere into the mantle. This involves almost the full thickness of the lithosphere in case of oceanic subduction, and probably most of the continental mantle lithosphere in case of a continent-continent collision. In the Alpine-Mediterranean area, regional P-wave tomography has shown that subducted lithospheric fragments accumulate in the mantle transition zone beneath most of Europe (e.g. PIROMALLO & MORELLI, JGR, 2003).

Here we present new images of the main mantle discontinuities (nominally at 410 and 660 km) bounding the transition zone beneath the Carpathian-Pannonian region. The ~35 km station spacing of the ~1.5 year Carpathian Basins broadband seismology project deployment along three NW-SE arrays, together with the permanent stations in Hungary and surrounding countries, provide a good 3-D coverage of the transition zone beneath this region, and achieve higher resolution than previous seismological studies there. We process the dataset to enhance P-to-S converted waves (receiver functions), which are then migrated to depth to obtain the 3D image of the discontinuities. The so-called '410' and '660' interfaces, which are in general related to phase changes in olivine, appear clearly. The measured depths show evidence for a transition zone thicker than the global average, which is attributed to the presence of cold material, consistent with the tomographic images. Moreover, below the eastern part of the Pannonian Basin, we determine that the '660' interface is further depressed by at least 15 km. Our interpretation is that this anomalously deep phase change may reveal the initial stages of further downwelling of cold subducted material into the uppermost lower mantle, advecting the phase change surface downward. Similar depression of this phase change is observed beneath some of the currently active oceanic subduction zones.

New insights into the Oligocene to Miocene evolution of the Austrian deep water Molasse Basin: implications for facies distribution and hydrocarbon exploration

HINSCH, R.

Rohöl-Aufsuchungs AG, Schwarzenbergplatz 16, 1015 Vienna, Austria; ralph.hinsch@rohoel.at

The understanding of the detailed geological evolution of the Molasse Basin is crucial for the continued success of exploration in this mature basin. Risks associated with play types can be better assessed considering their evolutionary framework. Results from several research projects increase the understanding of stratigraphic traps by characterising the sedimentological processes that control them and help to find new play types.

Several studies, focussing on the sedimentological and stratigraphic architecture, have been concluded in recent years or are still ongoing. This presentation will give an overview of the objectives, results and implications of these initiatives for evaluation and

analysis of the geological evolution and for exploration of the Molasse Basin.

An initial collaboration with Stanford University integrated sedimentological core analyses with 3D seismic, wireline log data interpretation and outcrop studies in analogue settings. The study yielded a modern sedimentological model for the Upper Puchkirchen Formation which was subsequently applied to exploration. A sequence stratigraphic study examined the sequence framework of the Molasse Basin fill and was able to correlate 5 sequences from the shelf into the deep basin. Studies on seismic and core analyses from the south slope of the Puchkirchen trough show how slope morphology and confinement control sediment distribution in the southern slope deposits. The transition from deep to more shallow marine conditions and the progradation of deltaic sequences into the basin in Eggenburgian/Burdigalian times is described by an intense 3-D seismic interpretation in combination with sedimentological core work. Working on a more local scale, other projects are improving the understanding of the detailed architecture of distinct play elements such as the Upper Puchkirchen Channel or the Basal Hall Formation Channel.

In general, these studies highlight the complex interaction of processes that control sediment distribution in the basin. Morphology, tectonics, sediment input and sea level changes all contribute to the sedimentary architecture of the deep water basin. Continued research needs to detail the interactions in order to enhance trap prediction for continued exploration success.

Reassessing seismic-slip values and potential seismogenic fault areas in the Vienna Basin: implications for seismic potential

HINSCH, R.^{1,2} & DECKER, K.¹

¹Department of Geodynamics and Sedimentology, Univ. of Vienna, Althanstrasse 14, 1090 Vienna, Austria; ²present address: Rohöl-Aufsuchungs AG (RAG), Schwarzenbergplatz 16, 1015 Vienna, Austria, ralph.hinsch@rohoel.at

The Vienna Basin Transfer Fault (VBTF) is a slow moving active fault passing through the most populated and most productive region of Austria and Slovakia. Recent activity is indicated by moderate seismicity ($I_{\max} \sim 8-9$, $M_{\max} \sim 5.7$), quaternary geology (tilted fluvial terraces and fault-controlled Piestocene basins) and geomorphology (linear scarps with hanging valleys). The purpose of this study is to use and discuss different data sets that constrain the seismic potential of the Vienna Basin Transfer Fault (VBTF). Seismic slip values corresponding to the seismic energy release are calculated from an earthquake catalogue spanning the Austrian and Slovakian part of the VBTF in the vicinity of the Vienna Basin (ACORN 2004). A 3-D model of a generalized VBTF in the area of the Vienna Basin is constructed. Segmentations from recent geological interpretations are used to define potential rupture zones on the generalized fault. The seismic slip for the VBTF from Semmering to Dobra Voda is in the range of 0.22 – 0.31 mm/yr, assuming a seismogenic fault thickness of 8 km. Calculating the slip rates for individual fault segments shows, that the rates vary significantly between the segments, from virtually no seismic slip to values as high as 0.77 mm/yr (Dobra Voda segment). Comparing these data to GPS derived velocities (> 1 mm/yr, GRENERCZY et al. 2005) show, that the fault yields a significant seismic slip deficit. Segments of the fault with apparently no seismic slip contrast from segments which might be locked. The modeled fault surface areas within the seismogenic zone (4-14 km depth range) of the VBTF vary from 55 km² to more than 400 km², with segmentation information partly still incomplete. Using these areas with empirical observations (WELLS & COPPERSMITH 1994) confirms, that the areas are sufficiently large to explain

observed earthquakes as well as earthquakes with magnitude >6 as recently postulated by archeo-seismological data (DECKER et al. 2006). Surface areas appear large enough to allow potential earthquakes with magnitudes exceeding 6.5. Given an incomplete historic earthquake catalogue, seismic slip deficits in segments, sufficient large rupture areas and indications of pre-catalogue strong earthquakes we argue, that the maximum earthquake size for the VBTF is significantly larger than the historically observed M_{\max} 5.7, and is more likely in the range M 6-6.5.

- ACORN (2004): Vienna (Central Institute for Meteorology and Geodynamics, Department of Geophysics) - Brno (Institute of Physics of the Earth, University Brno).
 DECKER, K., GANGL, G. & KANDLER, M. (2006): Journal of Seismology **10**/4: 479-495
 GRENERCZY, G., SELLA, G., STEIN, S., KENYERES, A. (2005): Geophysical Research Letters, **32** (16).
 WELLS, D.L. & COPPERSMITH, K.J. (1994): Bulletin of the Seismological Society of America, **84**/4: 974-1002.

Brittle-fault deformation history in the NW Himalaya (Himachal Pradesh, India)

HINTERSBERGER, E.¹, DECKER, K.², THIEDE, R.³ & STRECKER, M.⁴

¹Institut für Geowissenschaften, Universität Potsdam, Karl-Liebknech-Str. 24, 14476 Potsdam, Germany; Geozentrum, Universität Wien, Althanstr. 14, 1090 Wien, Austria; Geologisches Institut, ETH Zürich, Haldenbach-Str. 44, 8092 Zürich, Switzerland; Institut für Geowissenschaften, Universität Potsdam; estherh@geo.uni-potsdam.de, kurt.decker@univie.ac.at; rasmus.thiede@erdw.ethz.ch; strecker@geo.uni-potsdam.de

Since the Indian-Eurasian collision, the Himalayan mountain belt has formed as the southern termination of the Tibetan Plateau. While at present day, thrusting at lower elevations within the Lesser Himalaya is observed, several generations of extensional structures have been detected in the high-elevation regions of the Higher Himalaya, both parallel and perpendicular to the strike of the orogen. In the Sutlej-Spiti River Valleys and the Garhwal in the NW Himalaya (India), focal mechanisms of larger earthquakes and regional GPS measurements reveal ongoing E-W extension in this area. In contrast to other extensional features observed in the Himalaya, this direction is neither parallel nor perpendicular to the NW-SE regional shortening direction.

Here we present new geological data sets such as new structural geological mapping, fault kinematic analysis of hundreds of brittle faults, and satellite imagery analysis covering the area between the Tso Morari Lake in the Tibetan Himalaya in the north and the mountain front in the Garhwal Himalaya in the south.

In the Garhwal Himalaya and the Sutlej-Spiti River Valleys, we collected small brittle normal fault planes on outcrop scale with displacements up to several cm that cover the whole region from Tibetan Himalaya down to the mountain front in a densely spaced network. To analyze fault kinematic data (strike and dip of the fault, slip direction and sense of slip) for these small fault planes, we calculated strain axes for approx. 100 outcrop locations using the program TectonicsFP. This data set as well as field observations such as crosscutting relationships, mineralization at fault planes and correlations with deformation structures in lake sediments allows us to separate different deformation phases. In addition, we have mapped extensional structures in the Spiti River Valley and the Tso Morari Lake, both in the field and on satellite imagery, and compiled them with structures already shown in geological maps.

These data sets allow us to get a detailed overview of the deformation history in this area which is recorded in brittle faults.

In the southern and western part of our study area, brittle faulting records mainly shortening perpendicular to the strike of the orogen. In the northeastern part, there are at least three different directions of extension. The most recent deformation phase is the E-W extension that is also recorded in the seismicity data and GPS measurements. Our data set, however, shows that this extension is affecting a much more larger region towards the south than the seismicity data would suggest.

Mathematische Betrachtungen der Kohlensäuregleichgewichte

HOBIGER, G.

Geologische Bundesanstalt, Neulinggasse 38, 1030 Wien;
 gerhard.hobiger@geologie.ac.at

Das in der Klimadiskussion im Mittelpunkt stehende Kohlendioxid kommt in der Atmosphäre zu 0,03 Vol% vor. Es tritt daher in direkter Wechselwirkung mit der gesamten Hydrosphäre, wodurch sich Kohlensäure und deren Salze in den Gewässern finden. Eine entscheidende Rolle spielen dabei die chemischen Gleichgewichte der Kohlensäure und deren Salze in Wasser mit dem Kohlendioxid der Atmosphäre. Mit dem theoretischen Verständnis dieser chemischen Gleichgewichte kann die Wechselwirkung eines Fließgewässers mit der Atmosphäre modellhaft beschrieben werden. Es werden die theoretischen Grundlagen für die chemischen Gleichgewichte zwischen Kohlendioxid und Wasser mit einer bestimmten Alkalinität allgemein dargestellt. Vom chemischen Gesichtspunkt aus gesehen sind nur wenige einfache chemische Gleichgewichte zu berechnen. Dabei wird gezeigt, welche mathematischen Beziehungen zwischen den einzelnen Parametern existieren. Die Anzahl der möglichen mathematischen Gleichungen werden mit kombinatorischen Methoden berechnet. Da das allgemeine System zwei Freiheitsgrade besitzt, ergeben sich zunächst 16 Systeme mit jeweils drei voneinander unabhängigen Variablen, wobei jeweils zwei bekannt sein müssen um die dritte zu berechnen. Aus diesen 16 Systemen folgen 48 mathematische Beziehungen mit denen das gesamte chemische System beschrieben werden kann. Es ist daher möglich mit nur zwei bekannten Variablen (2 Freiheitsgrade) das gesamte Gleichgewichtssystem zwischen Kohlendioxid und Wasser mit Alkalinität exakt zu berechnen. Exakt bedeutet, dass von allen beteiligten chemischen Gleichgewichten sämtliche Bestimmungsgleichungen ohne Näherungen mitberücksichtigt werden. Nicht berücksichtigt wird der Ionenstärkeeinfluss. Es werden daher ideale Bedingungen angenommen, und die Aktivitätskoeffizienten eins gesetzt. Der Einfluss von nicht idealen Bedingungen kann allerdings jederzeit mit entsprechenden Näherungsformeln (z. B. Debye-Hückel) bei numerischen Berechnungen mitberücksichtigt werden. Wie oben erwähnt genügen im Prinzip nur zwei Variablen um alle anderen zu bestimmen. Dies ist zwar mathematisch richtig, aber in der chemischen Praxis können in sinnvoller Weise nur bestimmte Parameter vorgegeben werden. Dies folgt unmittelbar aus der Chemie der Kohlensäure und aus den beteiligten chemischen Gleichgewichten. Wird im allgemeinen Fall die Alkalinität Null gesetzt, so erhält man direkt das Gleichgewicht von Kohlendioxid mit reinem Wasser. Dieser Spezialfall besitzt nur einen Freiheitsgrad und es existieren 20 unabhängige Gleichungen, die zur Beschreibung dieses Systems verwendet werden können. Dies wird ebenfalls mit kombinatorischen Methoden gezeigt und alle aus den bereits bekannten Gleichungen des allgemeinen Falles hergeleitet. Dieser Spezialfall ist die theoretische Grundlage für eine Verallgemeinerung auf alle Gase, die in Wasser eine 2-basige Säure bilden.