

hydrofracture veins (alternately filled by ore-rich fluids, quartz and calcite), cross-cut the footwall and fault-rock-zone lithologies. While the high-angle normal fault system is characterized by two distinct oriented sets, acting as weakening zones and fluid migration pathways, the subvertical hydrofracture veins predominantly occur in the footwall.

Field based and microstructural studies of the Otzias Bay Detachment reveal that aseismic frictional-viscous creep is interspersed with periods of velocity weakening.

BioSoil – das europäische Waldboden-Monitoring-Projekt

MUTSCH, F.

Bundesforschungs- und Ausbildungszentrum für Wald, Naturgefahren und Landschaft, Institut für Waldökologie und Boden, Seckendorff-Gudent-Weg 8, 1131 Wien; franz.mutsch@baw.gv.at

Im Jahr 2005 hat die EU im Rahmen von Forest Focus eine erste Wiederholungsaufnahme des europäischen Waldboden-Monitorings beauftragt - das Projekt BioSoil. Europaweit werden rund 5000 Flächen beprobt, 139 davon liegen in Österreich. Ende 2009 soll das Projekt abgeschlossen sein.

Vorrangiges umweltpolitisches Ziel dieser europaweiten Erhebung ist, den aktuellen Zustand der europäischen Waldböden zu erfassen und ihre nur langsam ablaufenden Veränderungen zu verfolgen wie beispielsweise:

- die Wirkung bereits erfolgter Maßnahmen (Luftreinhaltung, Waldbodenmelioration, naturnaher Waldbau) zum Schutz und zur Stabilisierung von Waldökosystemen zu überprüfen,
- den Stickstoffstatus von Waldböden und deren Empfindlichkeit gegenüber Stickstoffeinträgen zu untersuchen,
- die Kohlenstoffspeicherung und die Änderung des Kohlenstoffvorrates von Waldböden (Klimawandel, Kyoto-Protokoll) abzuschätzen und
- die Belastung von Böden mit Schadstoffen (z. B. Schwermetallen) zu bewerten.

Außerdem sollen die vorhandenen Arbeitsanleitungen und Vorschriften des ICP Forests für ein europaweit harmonisiertes Vorgehen beim Waldboden-Monitoring erprobt und verbessert werden. Eine Reihe offener Fragen ist dazu zu beantworten:

- Kann eine Wiederholungsaufnahme nach 10-20 Jahren Veränderungen einzelner Bodenparameter erfassen?
- Sind mögliche Veränderungen statistisch signifikant?
- Sind Veränderungen aus Ursachen-Wirkungsbeziehungen erkläbar?
- Können die vorliegenden Methoden im europäischen Maßstab angewandt werden?
- Sind die Ergebnisse spezieller Methoden reproduzierbar?
- Sind die Ergebnisse zwischen den einzelnen Mitgliedsstaaten vergleichbar?
- Sind die Ergebnisse in ein größeres europäisches Bodeninformationssystem integrierbar?

Es werden die methodischen Grundlagen der Geländearbeit und der Laboranalysen erläutert sowie die Möglichkeiten der Qualitätsicherung und der Qualitätskontrollen besprochen. Erste Zwischenergebnisse aus Österreich betreffend die lokale Variabilität chemischer Parameter von Waldböden und deren Veränderungen nach 20 Jahren werden präsentiert. Zusätzlich wird das Problem des Vergleichs neuer Analysedaten mit alten Analysedaten diskutiert, welches durch die Nachanalyse alter Proben teilweise behoben werden kann.

The Permian to Lower Triassic siliciclastic base of central Northern Calcareous Alps revisited: part of a rift zone

NEUBAUER, F., EBNER, C., LACKNER, M. & SCHMIDLECHNER, M.

Dept. Geography and Geology, University of Salzburg,
Hellbrunner Str. 34, A-5020 Salzburg, Austria;
franz.neubauer@sbg.ac.at

Permian to Lower Triassic siliciclastic successions and their petrographic composition of conglomerate, breccias and sandstones of central southern sectors of the Northern Calcareous Alps have been studied in order to reveal their geodynamic setting. These successions exhibit a spatially variable sequence of new-defined formations, which are separated along strike by new detected low-angle normal faults (e.g. the Halm fault). A halfgraben-type siliciclastic infilling includes a ca. 1.5-2 km thick sequence starting with the Filzmoos Conglomerate containing coarse polymict breccia and quartz conglomerate, thick Alpine Verrucano, and the Gerzkopf Fm. representing a thick quartz arenite at the top. These are overlain by the shallow marine Werfen Formation. Only in light-coloured sandstones intercalated within purple siltstones, some trace fossils have been found. The Alpine Verrucano Formation also contains a layer of pure green sericite slates interpreted to represent a completely transposed acidic tuff. The entire sequence is interpreted to represent the transition from desert climate to shallow marine barriere sandstones during prograding extension within a syn-rift geodynamic setting. Overlying siliciclastic sediments are interpreted to monitor changes between terrestrial and marine environments until a full marine environment was established in upper parts of Lower Triassic in a post-rift tectonic setting. The thick graben infill is juxtaposed to relatively thin successions deposited on horst structures. These lack coarse clastic formations and are dominated by fine-grained sandstones.

A higher tectonic unit of the Werfen Imbricate zone comprises similar clastic-dominated but thinner formations as the underlying thick successions. This succession also includes deformed shales of the Haselgebirge Formation underlying the Lower Triassic Werfen Formation.

The new data show that main portions of the siliciclastic base of the Northern Calcareous Alps can be interpreted as a rift succession and that it monitors syndepositional extension. We correlate the units of central southern base Northern Calcareous Alps with recently studied Permian to Lower Triassic units of Lower Austroalpine Quartzphyllite nappe, which represent another syn-rift graben infill.

Cretaceous tectonic processes revealed by Gosau-type collapse basins in the Alpine-Balkan-Carpathian-Dinaric orogenic belt

NEUBAUER, F., GENSER, J. & WIESINGER, M.

Dept. Geography and Geology, University of Salzburg,
Hellbrunner Str. 34, A-5020 Salzburg, Austria;
franz.neubauer@sbg.ac.at, Johann.Genser@sbg.ac.at

The Alpine-Balkan-Carpathian-Dinaric (ABCD) orogenic belt resulted from two stages of closure of two ocean basins (Meliata-Vardar, Penninic) and two, Cretaceous and Eocene-Neogene, stages of plate collision between the European/Moesian plates in the foreland and the Adriatic microplate in the hinterland. Here, we discuss the presence and structural and palaeogeographic relationships of similar Upper Cretaceous collapse basins, which are

commonly known as Gosau basins in the Eastern Alps, in all parts of the ABCD belt. Similar collapse basins occur all along the northern, Alpine-Carpathian-Balkan strand as well as in Dinarides. Common features include: (1) age postdating oceanic, Meliata - Vardar, closure; (2) an age postdating the formation of the Cretaceous-aged nappe stack and peak conditions of the Early Alpine metamorphism; (3) synchronicity with exhumation of metamorphic orogenic wedge and exhumation of metamorphic core complexes as well as of previously subducted continental crust; and (4) deposition of high proportions of siliciclastic material reflecting relief, exhumation and denudation of metamorphic crust. In the eastern part of the Carpathian-Balkan strand, these Gosau basins are associated with a linear belt of calc-alkaline volcanic and shallow-plutonic successions known as Banatite Suite. A lateral diachroneity in onset is observed.

A summary of existing and new own Ar-Ar single-grain age data from these Gosau basins covering the entire ABCD-belt reveals a direct connection with exhuming metamorphic crust. The proportion of Cretaceous-aged white mica is high in the west, and low in the existing reflecting the exhumation of the high-pressure belt in Alps and Apuseni Mountains.

The Meliata/Vardar oceanic basin started to close during the Middle Jurassic. The final closure of the Meliata basin occurred during the Early Cretaceous with the formation of a deep sea trench. The overall process is A-subduction and high-pressure metamorphism of the distal Austroalpine continental margin, and subsequent exhumation of high-pressure rocks and in overall transpressional setting. Geochronologic data monitor continuous footwall accretion of cover and basement/cover nappes between 120 and 50 Ma. Late Cretaceous Gosau basins seal the Meliata suture and nappe stacking structures. Formation of these basins in the Eastern Alps was associated with sinistral wrenching, northward tilting of the orogenic wedge, normal faulting at shallow crustal levels and exhumation of eclogite-bearing crust within Austroalpine units. A model with trench retreat of the subducting Austroalpine/Penninic lithosphere can explain exhumation of high-pressure rocks, accretion of subducting material and subsidence in the upper plate, where Gosau-type basins formed.

Geochemical characteristics of the Miocene Volcanism in SE Styria, Austria

NTAFLOS, T.¹ & KRUMPEL, G.²

¹Dept. of Lithospheric Research, University of Vienna; ²Veitsch-Radex Leoben; theodoros.ntaflos@univie.ac.at, georg.krumpel@rhi-ag.com

The Styrian basin, the westernmost sub-basin of the Pannonian basin system, is located at the eastern margin of the Eastern Alps. The majority of the studied samples were revealed by drilling cores in the southeastern part of the Styrian Basin. The rest of the samples have been collected from quarries.

The rocks are high-K calc-alkaline (andesites and dacites) and alkaline (latites and shoshonites). K/Ar-ages range from 16.3 to 13 Ma with high-K calc-alkaline rocks representing the oldest suite. A striking petrographic difference between both groups is that the calc-alkaline rocks are amphibole bearing and lack of clinopyroxene. Some zoned clinopyroxenes from the alkaline rocks have magmatic corroded cores overgrown by more basic rims suggesting basic magma reflux.

In the spiderdiagrams all the samples show remarkable negative Nb and Ta troughs and high contents of LILE. Additionally the calc-alkaline rocks have negative P anomaly. However, the calc-alkaline rocks are characterized by low contents of LILE and HFSE when compared to the alkaline rocks.

The calc-alkaline rocks from Styria are similar to those from the

Carpathian-Pannonian area that have long been recognized as being typical for subduction related-magmas (SEGHEDI et al. 2004). They can be modelled by an AFC-process controlling the composition of an IAB-magma with F=0.6 and r=0.4. As contaminant is proposed the upper continental crust.

While high-K calc alkaline rocks can be explained by an AFC-process the composition of the alkaline rocks (latites and shoshonites) seems to be a consequence of batch melting processes of a pre-enriched mantle, which can be derived by mixing of primitive lithospheric mantle with material from the continental crust. Model calculations reveal that the fraction of continental material in the mixture was 4% and the batch melting process took place in the stability field of spinel-lherzolite at 5%.

From the geodynamical point of view, the calc-alkaline lavas can be seen as a consequence of the subduction of the Rhenodanubian-flysche or the subduction of the West-Carpathian-Flysche. This subduction was related to the N-S convergence of Africa and Europe in the Early Miocene. In contrast the alkaline magmatism was the result of decompressional melting of a metasomatic pre-enriched mantle. The process of decompressional melting may be related to the extension that characterizes the western margin of the Pannonian Basin during the Middle Miocene (SEGHEDI et al. 2004).

SEGHEDI, I., DOWNES, H., SZAKACS, A., MASON, P.R.D., THIRWALL, M.F., ROSU, E., PECSKAY, Z., MARTON, E. & PANAIOTU C. (2004): Neogene-Quaternary magmatism and geodynamics in the Carpathian-Pannonian region: a synthesis. - *Lithos*, **72**: 117-146.

Multiple Particle Interactions

O'CONNOR, A.¹ & MULCHRONE, K.²

¹Department of Geodynamics and Sedimentology, Universität Wien, Austria, Althanstrasse 14, 1090 Wien; ²Department of Applied Mathematics, University College Cork, Ireland; a.oconnor@ucc.ie, k.mulchrone@ucc.ie

A mathematical model is developed to account for the rotational disturbances of rigid ellipsoids caused by interactions between neighbours. The equations of motion of rigid particles in isolation are adapted via the Kuramoto model which implements synchronisation among the set of interacting particles. The orientation of the ellipsoids is determined by three angles (the Euler angles) which in turn determine the angular velocity at any point in time. Rather than adjusting the perturbed velocity field, a noise term is appended to the equations that determine the angular velocity. This term depends on the neighbouring particles' degree of coupling, proximity and relative position. Simulations reveal a preferred orientation at an angle to the shear direction that increases with the degree of coupling.

Slump structures and paleoslope: Case study in pelagic limestones of the Ammergau Formation, Ampelsbach gorge, Northern Calcareous Alps, Tyrol, Austria

ORTNER, H. & KILIAN, S.

Institut für Geologie und Paläontologie, Universität Innsbruck, Innrain 52, 6020 Innsbruck; sinah.kilian@studentuibk.ac.at, hugo.ortner@uibk.ac.at

One method to estimate paleoslope orientation is to study and interpret slump folds. Existing methods for analysis of slump structures require collection of orientation data on a scale similar