

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 20/1	Graz 2014
PANGEO AUSTRIA 2014		Graz, 14. September 2014 – 19. September 2014	

No late Permian tsunamites in Guryul Ravine (Kashmir, India) but frequent storms and seismic activity

KRYSZYN, L.¹, HORACEK, M.², BRANDNER, R.³, PARCHA, S.K.⁴

¹ University of Vienna, Department for Paleontology, Althanstraße 14, 1090 Vienna, Austria

² BLT Wieselburg, Lehr- und Forschungszentrum Francisco-Josephinum, Rottenhauserstr. 1, 3250 Wieselburg, Austria and University of Vienna, Department for Lithospheric Research, Althanstraße 14, 1090 Vienna, Austria

³ University of Innsbruck, Department for Geology, Innrain 52, 6020 Innsbruck, Austria

⁴ Wadia Institute of Himalayan Geology, Gen. Mahadeo Singh Road 33, 248001 Dehra Dun, India

Recent claims for tsunami-related event beds induced by the Siberian Trapp basalts in this section (Brookfield et al., 2013) have to be questioned. Identical storm generated carbonate beds occur not only during a short interval close to the Permian-Triassic (P-T) boundary but through a major part of the late Permian (Changhsingian) succession there – as low as 26 m below the so-called tsunami beds. Moreover, during our recent study in a closely neighbouring place called Mandakpal (less than 10 km to the southeast), no signs of tsunamites have been detected in time-correlative finegrained sediments. Based on sedimentary and trace fossil evidence we interpret the late Permian of Guryul as relatively shallow, neritic and delta-influenced. The so-called tsunamites are shelly-enriched discontinuous carbonate lenses fed downslope through local channels. Judging from the distinct facies change from sandy towards muddy background sedimentation at the base of the “tsunamites” a sudden deepening may be explained by local and still rift-related tectonics along the NIM (North-Indian Gondwana Margin) which led to episodic seismic induced sediment redeposition in channel influenced places like Guryul. Synsedimentary tectonic activity with tilting and eventual Horst and Graben structure building along the large NIM is indicated by margin inversion during the P-T boundary interval leading to sedimentary breaks and 20 times thinner, condensed limestone deposits far offshore from Guryul in Spiti (Krystyn et al., 2007) and Tibet (Orchard et al., 1996). Thus, local seismic activity seems to be a far more logic explanation of the Guryul “tsunamites” than the eruption of the Siberian Traps more than 6000 km away.

This is a contribution to IGCP572.