



Oxidation of SO₂ by stabilized Criegee intermediate (sCI) radicals as a crucial source for atmospheric sulfuric acid concentrations

M. Boy¹, D. Mogensen^{1,2}, S. Smolander¹, L. Zhou¹, T. Nieminen¹, P. Paasonen^{1,7}, C. Plass-Dülmer³, M. Sipilä¹, T. Petäjä¹, L. Mauldin^{1,4,5}, H. Berresheim⁶, and M. Kulmala¹

¹Department of Physics, P.O. Box 48, University of Helsinki, 00014 Helsinki, Finland

²Helsinki University Centre for Environment, P.O. Box 27, University of Helsinki, 00014 Helsinki, Finland

³Hohenpeissenberg Meteorological Observatory, German Weather Service, Hohenpeissenberg, Germany

⁴Department of Atmospheric and Oceanic Sciences, University of Colorado at Boulder, P.O. Box 311, Boulder, Colorado 80309-0311, USA

⁵Institute for Arctic and Alpine Research, University of Colorado at Boulder, P.O. Box 450, Boulder, Colorado 80309-0450, USA

⁶Center for Climate and Air Pollution Studies, School of Physics, National University of Ireland Galway, Galway, Ireland

⁷International Institute for Applied Systems Analysis (IIASA), Schlossplatz 1, 2361 Laxenburg, Austria

Correspondence to: M. Boy (michael.boy@helsinki.fi)

Received: 24 September 2012 – Published in Atmos. Chem. Phys. Discuss.: 22 October 2012

Revised: 3 March 2013 – Accepted: 17 March 2013 – Published: 12 April 2013

Abstract. The effect of increased reaction rates of stabilized Criegee intermediates (sCIs) with SO₂ to produce sulfuric acid is investigated using data from two different locations, SMEAR II, Hyytiälä, Finland, and Hohenpeissenberg, Germany. Results from MALTE, a zero-dimensional model, show that using previous values for the rate coefficients of sCI + SO₂, the model underestimates gas phase H₂SO₄ by up to a factor of two when compared to measurements. Using the rate coefficients recently calculated by Mauldin et al. (2012) increases sulfuric acid by 30–40%. Increasing the rate coefficient for formaldehyde oxide (CH₂OO) with SO₂ according to the values recommended by Welz et al. (2012) increases the H₂SO₄ yield by 3–6%. Taken together, these increases lead to the conclusion that, depending on their concentrations, the reaction of stabilized Criegee intermediates with SO₂ could contribute as much as 33–46% to atmospheric sulfuric acid gas phase concentrations at ground level. Using the SMEAR II data, results from SOSA, a one-dimensional model, show that the contribution from sCI reactions to sulfuric acid production is most important in the canopy, where the concentrations of organic compounds are the highest, but can have significant effects on sulfuric acid concentrations up to 100 m. The recent findings that the reaction of sCI + SO₂ is much faster than previously thought

together with these results show that the inclusion of this new oxidation mechanism could be crucial in regional as well as global models.

1 Introduction

New particle formation in the troposphere is important for the global concentration of cloud condensation nuclei (CCN) (Spracklen et al., 2008; Makkonen et al., 2009; Pierce and Adams, 2009; Wang and Penner, 2009; Merikanto et al., 2010). Recently, Makkonen et al. (2012) presented the effect of new particle formation on anthropogenic climate forcing in present-day and future (year 2100) conditions and concluded that with total aerosol forcing diminishing in response to air pollution controls taking effect (especially reductions in sulfur dioxide (SO₂)), warming from increased greenhouse gas concentrations can potentially increase at a very rapid rate. According to several studies (e.g. Kulmala and Pirjola, 2000; Paasonen et al., 2010; Sipilä et al., 2010; Zhao et al., 2010; Lauros et al., 2011; Kirkby et al., 2011), sulfuric acid (H₂SO₄) is one of the initial or required molecules in the nucleation mechanism. In order to quantify future atmospheric sulfuric acid concentrations, a complete

understanding of the sink and source terms is crucial. Although the precursors for sulfuric acid (at least some of them), as well as the main sink term (condensation on atmospheric aerosols) have been measured in several places, the closure between measured and calculated sulfuric acid concentrations has rarely been investigated (Eisele et al., 1993; Weber et al., 1997; Boy et al., 2005).

Criegee intermediate (CI) radicals can play a crucial role in tropospheric oxidation, as suggested more than a decade ago by Calvert et al. (2000). The CI formation mechanism starts from the ozonolysis of alkenes, with an addition of ozone to the double bond forming a primary ozonide with high excess energy. The excess energy causes the primary ozonide to decompose instantaneously to the Criegee intermediate, which will still possess excess energy. In order to release its excess energy, the Criegee intermediate either decomposes into different products or collisionally stabilizes (we refer to the latter as a stabilized Criegee intermediate (sCI)). The stabilized CI can then react with various atmospheric compounds, particularly H₂O, NO_x, SO₂, CO and many others.

Recently, Welz et al. (2012) pointed out that ozonolysis of unsaturated hydrocarbons (e.g. terpenes) is a major removal process in the troposphere for organic compounds and proceeds via Criegee intermediate radicals. In their work, the authors reported a more than three orders of magnitude higher reaction rate constant for formaldehyde oxide (CH₂OO) with sulfur dioxide than was reported in the literature until today ($k = 3.9 \times 10^{-11} \pm 0.7 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ at 298 K and 4 torr). The potential for different CIs to oxidize compounds like SO₂ is much stronger than previously expected and is thus crucial to be included in all chemical models.

New atmospheric observations supported by laboratory experiments and theoretical considerations point to the existence of compounds (most probably stabilized CIs) which have significant capacities to oxidize atmospheric trace gases like sulfur dioxide (Mauldin III et al., 2012). These authors claimed rate constants for Criegee intermediate radicals originating from the ozonolysis of α -pinene and limonene to be $6 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$ and $8 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$, respectively. These new rate constants are about one order of magnitude higher than assumed as for example in the Master Chemical Mechanism published by the University of Leeds, Great Britain (<http://mcm.leeds.ac.uk/MCM/>).

In this work we investigated the effect of increased reaction rate constants of SO₂ with Criegee intermediate radicals on the atmospheric concentrations of sulfuric acid for two different stations: SMEAR II, Hyytiälä, Finland, and Hohenpeissenberg, Germany. Our results show that depending on the concentrations of the organic compounds, their contribution via the reaction of stabilized Criegee intermediates to atmospheric gas phase sulfuric acid concentrations is crucial at ground level. We further studied the role of this new oxidation mechanism in the lower troposphere (up to 2 km) with the one-dimensional chemical transport model SOSA for the

Finnish station SMEAR II. In the last two sections we will give a short discussion about the uncertainties in the calculations of different parameters and present a sensitivity study on the effect of the reaction rates between Criegee intermediates and water on the overall budget of sulfuric acid.

2 Measurements

For our model investigations, we used data from two different stations. These data included: measured concentrations of volatile organic compounds (VOC), hydroxyl radical (OH), sulfuric acid, and other inorganic gases in addition to particle size distributions and basic meteorological data. The next three subsections will present information about the stations and the used data sets.

2.1 Hyytiälä

Measurements in Hyytiälä were performed at the University of Helsinki's SMEAR II station (Station for Measuring Ecosystem–Atmosphere Relations; Hari and Kulmala (2005); 61° 51' N, 24° 17' E). The surroundings of the station are a rather homogeneous Scots pine (*Pinus sylvestris*) forest, which was sown in 1962. The station is equipped with permanent instrumentation for measuring basic meteorological parameters, aerosol concentrations, photosynthesis and soil properties. In this work, we utilize measurements of temperature, wind speed, spectral irradiance, global and diffuse solar radiation intensity, SO₂, CO, NO, NO_x, O₃, VOC, OH and H₂SO₄ concentrations. Furthermore, particle size number concentrations from DMPS (differential mobility particle sizer) and APS (aerodynamic particle sizer) are included. A detailed description of the station and instrumentation (name, branch and detection limit) can be found under Kulmala et al. (2001a) and at <http://www.atm.helsinki.fi/SMEAR/index.php>.

For Hyytiälä we first selected 7 days (29 July to 4 August) during the HUMPPA-COPEC-2010 (Hyytiälä United Measurement of Photochemistry and Particles – Comprehensive Organic Particle and Environmental Chemistry) campaign which took place in July and August 2010 (Williams et al., 2011). For the results in Sect. 4.3, we used measurements from the whole year of 2010.

2.2 Hohenpeissenberg

The measurements at Hohenpeissenberg in rural southern Germany were conducted at the Meteorological Observatory Hohenpeissenberg (47° 48' N, 11° 0' E; e.g. Birmili et al., 2003, Rohrer and Berresheim, 2006). The observatory is operated by the German Weather Service (DWD) and is located atop the Hohenpeissenberg, at an altitude of 985 m a.s.l. and about 300 m above the surrounding terrain, which consists mainly of meadows and forests. At night, the site is generally above the nocturnal boundary layer and local emissions from

surrounding vegetation and anthropogenic sources do not reach the site. Data utilized from this station were global and diffuse radiation intensity, SO₂, CO, NO, NO_x, O₃, VOC, OH and H₂SO₄ concentrations. In addition, we used particle size number concentrations from a DMPS system.

For Hohenpeissenberg, a long-term data set from the year 2000 recorded during the HAFEX campaign (Birmili et al., 2003) covering the period January to August was available. However, the organic compounds, which are crucial for our studies were only measured at certain times per day. Based on the measurements for these parameters in combination with the availability of all other requested data, we selected 7 days where at least seven measurement points of VOC concentrations were performed during the day. These selection criteria limited our studies to the following days: 9, 10, 27 and 28 April and 18 to 20 June 2000.

2.3 OH, H₂SO₄ and VOC concentration measurements at the two stations

The sulfuric acid and OH concentrations were measured with a chemical ionization mass spectrometer (CIMS; Berresheim et al., 2000; Petäjä et al., 2009). The CIMS instruments applied in Hyytiälä (Petäjä et al., 2009) and in Hohenpeissenberg (Berresheim et al., 2000 see also <http://www.dwd.de/luftchemie>) are of slightly different design but calibrated with a similar procedure. The uncertainty of the CIMS instruments at the 2σ level has been estimated to be 39 % for sulfuric acid and 54 % for OH concentration (Mauldin III et al., 1999; Berresheim et al., 2000).

The measurements of monoterpene concentrations at the two sites were performed using different instruments. At Hyytiälä the total monoterpene concentration was obtained with proton transfer reaction mass spectrometry (PTR-MS; DeGouw and Warneke, 2007), whereas at Hohenpeissenberg the total monoterpene concentration is calculated as a sum of the single monoterpene concentrations measured using gas chromatography mass spectrometry (GC-MS; Plass-Duelmer and Berresheim, 2007). The monoterpenes detected with the applied GC-MS were α-thujene, tricyclene, α-pinene, camphene, sabinene, myrcene, β-pinene, α-phellandrene, Δ³-carene, α-trepenene, limonene and β-phellandrene.

3 Methods

In this work we used two different models, MALTE (Model to predict new Aerosol formation in the Lower Troposphere; Boy et al., 2006, 2008; Lauros et al., 2011) and SOSA (model to Simulate the concentrations of Organic vapours and Sulfuric Acid; Boy et al., 2011; Mogensen et al., 2011). Although MALTE – similar to SOSA – is a one-dimensional chemistry transport model, we used the latter to investigate the effect of the studied oxidation schemes on temporal trends in concen-

tration profiles for the SMEAR II station in Hyytiälä. This exploration can only be done with SOSA because the model is parallelized and runs on a computer cluster, so that the chemistry in different atmospheric layers can be computed in parallel. In the next two sections we will discuss the two models briefly, and in Sect. 3.3 we will define the selected chemistry scenarios in particular.

3.1 MALTE

The one-dimensional model MALTE is described in detail by Boy et al. (2006, 2008) and Lauros et al. (2011). The model reproduces the diurnal variation of boundary layer meteorology, chemistry, emissions and particle formation. Here, we used the zero-dimensional version of the model with different chemistry scenarios but without any aerosol dynamics.

The chemistry is calculated using the Kinetic PreProcessor (KPP) (Damian et al., 2002) and selected organic reactions from the Master Chemical Mechanism v3.2 (MCM-Leeds; Jenkin et al., 1997; Saunders et al., 2003) via the website <http://mcm.leeds.ac.uk/MCM/>. We included photochemical and inorganic reactions from the Master Chemical Mechanism v3.2 and from Atkinson et al. (2004), together with spectral irradiance measurements from the SMEAR II station (Boy and Kulmala, 2002). Measured inorganic gas concentrations (NO_x, SO₂ and CO), OH and VOC concentrations from SMEAR II and Hohenpeissenberg are used as input. The condensation sink for vapours was calculated from aerosol number size distributions measured with the twin DMPS and APS for Hyytiälä and by DMPS for Hohenpeissenberg using the method presented by Kulmala et al. (2001b).

3.2 SOSA

The one-dimensional chemistry transport model SOSA is described in detail by Boy et al. (2011) and Mogensen et al. (2011), thus we will only give a brief overview of the model here. SOSA consists of three modules dealing with meteorology, emissions and chemistry. The measured input data are obtained from SMEAR II and includes inorganic gas concentrations (NO_x, SO₂ and CO) together with condensation sinks for sulfuric acid and nitric acid, based on DMPS and APS data (Boy et al., 2003).

The meteorological module is described by a 1-D version of the coupled plant–atmosphere boundary layer model SCADIS (Sogachev et al., 2002, 2005; Sogachev and Panferov, 2006; Sogachev, 2009; Boy et al., 2011). SCADIS employs a 1.5 turbulent kinetic energy–specific dissipation closure scheme (including the Reynolds averaged Navier–Stokes equations for flow) using a number of parameterizations. The model is capable of describing the physical processes forming the meteorological regime within and above the forest canopy under different environmental conditions. The model describes the atmospheric boundary layer

evolution and the mixing of the chemical species within a model domain of a flexible amount of layers. A resolution of 100 layers is used in this study. The separation between the model layers increases logarithmically from the bottom to the top of the column (3000 m). Meteorological data from the European Centre for Medium-range Weather Forecasts (ECMWF) were used for nudging the model variables towards the observations and upper boundary meteorological conditions at 3000 m.

The emissions of biogenic organic vapours from the canopy are calculated with an implemented modification of the MEGAN model (Model of Emissions of Gases and Aerosols from Nature), version 2.04 (Guenther et al., 2006). MEGAN uses estimates of plant species composition, representative species-specific emission factors, and information about leaf temperature and solar radiation on sun and shade leaves at different canopy levels to simulate landscape average emissions. We have assumed a Scots pine forest environment and used 16 different canopy characteristics, such as leaf data together with scattering and reflection coefficients, to describe the conifer forest. Included also are standard emission potentials (SEP) for isoprene, α -pinene, β -pinene, Δ^3 -carene, limonene, sabinene, ocimene, farnesene, β -caryophyllene, 2-methyl-3-buten-2-ol, cineole, linalool and unspecified monoterpenes and sesquiterpenes (Hakola et al., 2006). The emission scheme has been verified by comparing VOC gas concentrations in the papers by Boy et al. (2011) and Mogensen et al. (2011).

The chemistry module is similar to the one used in MALTE. However, the concentrations of the hydroxyl radical and the organic vapours are predicted by the model SOSA, whereas measurements are used as input in MALTE.

3.3 Selected chemistry scenarios




To investigate the quantitative contribution of the stabilized Criegee intermediates to the oxidation of sulfur dioxide, we ran MALTE with 4 different chemistry scenarios which will be explained in detail below. All important rates related to the Criegee intermediates' chemistry (e.g. the decomposition rates of the CIs or the reaction rates of the CIs with other molecules like H₂O) are used as stated in the MCM-Leeds if not explicitly mentioned below. However, we will provide a discussion about the uncertainties of these values in Sects. 5 and 6.

Scenario A: Tropospheric relevant inorganic reactions selected from the Master Chemical Mechanism and Atkinson et al. (2004) were used.

Scenario B: The same inorganic chemistry as under scenario A plus additional organic reactions for isoprene and monoterpenes from the MCM. As the MCM currently only includes full chemistry pathways for the terpenes α -pinene, β -pinene and limonene, we adopted rate constants from Atkinson et al. (1994) for Δ^3 -carene, sabinene, myrcene and camphene with OH, O₃ and NO₃. The further chemical steps

for the first order reaction products of Δ^3 -carene, sabinene, myrcene and camphene with OH, O₃ and NO₃ have been approximated with the chemical paths for α -pinene, β -pinene and limonene depending on their mixing ratios and location of the double bond(s). We have approximated the chemical path for Δ^3 -carene, which is a bicyclic monoterpene with an endocyclic double bond, with α -pinene, which is also a bicyclic monoterpene with an endocyclic double bond. The chemical paths for sabinene and camphene, which are both bicyclic monoterpenes with an exocyclic double bond, have been approximated with the chemical path for β -pinene, which is likewise a bicyclic monoterpene with an exocyclic double bond. Myrcene, an acyclic monoterpene with three double bonds, was also added to the β -pinene path. By limiting our chemistry for the monoterpenes to these 7 compounds, we are able to cover generally more than 95 % of the observed monoterpene concentrations in Hyytiälä and Hohenpeissenberg (Plass-Dülmer and Berresheim, 2007; Bäck et al., 2012).

Scenario C: In this scenario we used the same chemical reactions as described in scenario B, but we increased the reaction rates of the sCIs with SO₂ based on the new values recently suggested by Mauldin III et al. (2012). Currently, the MCM assumes a reaction constant of $7 \times 10^{-14} \text{ cm}^3 \text{ s}^{-1}$ for all sCIs with SO₂, but with different compound-specific stabilization and loss rates for the different Criegee intermediates. Equations (1)–(3) present the crucial reactions for the oxidation of sulfur dioxide based on the MCM scheme, and Table 1 gives the original and modified reaction rate constants.




Combining the formation and the stabilization rates of the Criegee intermediates (multiplication of the yields of the two-step reactions) results in nearly equal values for α -pinene, β -pinene and limonene with 0.2, 0.102 and 0.135, respectively. However, the rates of the sCIs (not presented in Table 1) with water vapour (sometimes called water channel) from the MCM-Leeds show two-times higher loss for APINBOO and LIMOO compared to NOPINOO. This would lead to remarkably higher sulfur dioxide oxidation if high concentrations of β -pinene are present (which is not the case for the two selected stations).


Scenario D: In this scenario we investigate the effect of the stabilized Criegee intermediate formaldehyde oxide (CH₂OO), which is formed by the ozonolysis of isoprene and β -pinene (see Eq. 4). We increased the reaction rate as published by Welz et al. (2012) and also included a higher reaction rate for this sCI with NO₂ ($7 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$) as stated by Welz et al. (2012). Further, we used for the reaction of formaldehyde oxide with water the upper limit of the reaction rate from Welz et al. (2012) of $4 \times 10^{-15} \text{ cm}^3 \text{ s}^{-1}$. The

Table 1. Reaction rates for the most crucial chemical equations presented under Eqs. (1)–(4).

	k_1 (cm ³ s ⁻¹)	k_2 (s ⁻¹)	k_3 (MCM) (cm ³ s ⁻¹)	k_3 (new) (cm ³ s ⁻¹)
α -Pinene	$6.3 \times 10^{-16} \cdot \exp(-580/T) \cdot 0.4$	$1 \times 10^6 \cdot 0.5$	7×10^{-14}	6×10^{-13}
β -Pinene	$1.5 \times 10^{-17} \cdot 0.6$	$1 \times 10^6 \cdot 0.17$	7×10^{-14}	6×10^{-13}
Limonene	$2.95 \times 10^{-15} \cdot \exp(-783/T) \cdot 0.27$	$1 \times 10^6 \cdot 0.5$	7×10^{-14}	8×10^{-13}
Isoprene	$1.03 \times 10^{-14} \cdot \exp(-1995/T) \cdot 0.3$	$1 \times 10^6 \cdot 0.22$	7×10^{-14}	3.9×10^{-11}

**Fig. 1.** Measured temperature and global and diffuse radiation for the selected time periods at both stations.

combined formation and stabilization rate originating from isoprene is only 0.066. However, it should be considered that isoprene is the most abundant VOC in the atmosphere, and due to its high concentrations isoprene could still have a remarkable effect in the oxidation of SO₂.



4 Results and discussion

The results of our investigation of the oxidation capacity of sCIs to form sulfuric acid will be presented in three subsections. In the first we discuss the ambient measured conditions of several relevant parameters including the concentrations of OH and H₂SO₄. In the following subsection our zero-dimensional model results for the two stations will be presented, and in the last we show the average seasonal and yearly effect of the new oxidation mechanism in the lower part of the troposphere over the SMEAR II station in Hyytiälä.

4.1 Ambient conditions at the two stations

Figure 1 shows the temperature and radiation (global and diffuse) for the selected days at Hyytiälä and Hohenpeissenberg. During the days at the SMEAR II, exceptionally warm air masses with both the lowest and highest temperatures of 12.8 and 32.7 °C were present. Most of the time clouds formed during the day, which is visible in the small difference between the values of these two measured radiation parameters (Williams et al., 2011). At Hohenpeissenberg three different meteorological periods can be categorized: at the beginning of April, cold air masses with temperatures always below 10.4 °C but high solar irradiance with values up to 800 W m⁻² were observed; this was followed by a more cloudy but warmer period at the end of April; and a clear sky period with temperatures between 10 and 26 °C in June.

The condensation sink (CS) and the concentrations of sulfur dioxide for both stations are presented in Fig. 2. These two parameters are crucial in the production of H₂SO₄: CS is the most important sink term for sulfuric acid and proportional to the surface area of the existing atmospheric aerosols, while SO₂ is one of its precursors. The most obvious finding from the figure is that the values of both parameters are similar at both stations. Daily variations are not pronounced,




Fig. 2. Measured condensational sink and sulfur dioxide concentrations for the selected periods at both stations.

however the presence of more or less polluted air masses in Hyytiälä and Hohenpeissenberg can also be recognised as well. At the SMEAR II station, we observed relatively high concentrations of CS and SO₂ on the first day followed by much cleaner air over the next two days. During the last four days, both parameters are similar to values in the first three days with the exception of the last evening when the CS value decreased towards its lowest value of 0.0008 s⁻¹. Compared to Hyytiälä, Hohenpeissenberg has smaller variations both in and between the three different periods. It seems that during all selected days at this station no air mass changes with dramatic increases or decreases of pollution occurred.

A relatively strong difference in the daily pattern at the two stations can be found when observing the concentrations of organic compounds, hydroxyl radicals and sulfuric acid, which are presented in Fig. 3. The sum of the monoterpenes and isoprene measured at the SMEAR II station shows only a very small distinctive diurnal trend. This observation is in contrast with earlier studies at the same location where night-time values were a factor of 2–3 higher than during the day and the absolute concentrations were clearly higher compared to our selected days (e.g. Mogensen et al., 2011). One reason could be the nearly constant high temperatures during the selected period causing the closing of the stomata of the plants to decrease evaporation and loss of water. Also, the positive temperature dependence of the mixing layer height probably affects this pattern: even though the monoterpene emissions during the warmer day are higher, the concentration is more effectively diluted through the higher mixing layer. A completely different picture appears for Hohenpeissenberg with a clearly pronounced daily pattern for all three parameters. However, the ratio between day and night concentrations of the organic compounds, and vice versa, is as is normally observed at Hyytiälä. Daytime concentrations are

larger by up to a factor of 10, but the absolute concentrations are similar to those measured at the SMEAR II station. Measured VOCs are emitted at lower altitudes in the surroundings and then transported to the site by updrafts during convective meteorological conditions (high solar irradiance). At night, the observatory at Hohenpeissenberg is above the mixing layer, and thus the monoterpenes emitted in the surroundings do not reach the station before the increase of the mixing layer height in the morning. The reader should also remember that the volatile organic compounds in Hohenpeissenberg were not measured continuously with a PTR-MS as was the case for SMEAR II, but by online GC-MS (see Sect. 2.3). For this reason only time–space point measurements are available for the station in Germany, with linear interpolation being used to retrieve the daily variation.

The same discrepancy as for the organics can also be observed in the daily patterns between the two stations for the OH and H₂SO₄ concentrations. At both stations the profiles of these two parameters show a similar behaviour, but the increase during daytime is more pronounced at Hohenpeissenberg. However, at daytime the absolute values of the parameters are in the same range as would be expected at least for sulfuric acid by comparing the sink and source terms. We should also remember that at this time of the year the amount of daylight in Hyytiälä is around 18 h and that photochemistry is only inactive for a very short period during night. In Hyytiälä there exist strong differences between the hydroxyl radical and sulfuric acid concentrations on the first day, which could point to an extra source term of H₂SO₄ related to the observed much more highly polluted air mass during this time (see Fig. 2). Most probably the reaction products of organic compounds contribute to the oxidation of SO₂, which will be investigated in detail in Sect. 4.2.




Fig. 3. Measured concentrations of different organic vapours, hydroxyl radicals and sulfuric acid for the selected periods at both stations.

4.2 Process study near the ground at the two stations

In this subsection we used the zero-dimensional version of the model MALTE to investigate the oxidation capacity of the stabilized CI described in Sect. 3.3 to form sulfuric acid at the two selected stations, Hyytiälä and Hohenpeissenberg. During the simulations we kept the concentrations of the OH radicals and the VOCs constrained by the measurements. All other organic compounds and H₂SO₄ were calculated by the model. This setup was used to minimize the errors resulting from the model and to receive the best estimates for our study.

Figure 4 shows a scatter plot for the concentrations of the four Criegee intermediates using scenario B for the chemistry against the missing sulfuric acid concentrations calculated by subtracting the measured sulfuric acid concentrations from those predicted by MALTE running the chemistry scenario B. For the Hyytiälä plot we see a very similar behaviour for formaldehyde oxide and the sum of the three CIs resulting from the ozonolysis of the monoterpenes. This reflects the nearly identical behaviour of the precursors during the chosen period presented in Fig. 3. At Hohenpeissenberg the variability of the precursors is more pronounced, with larger increases of the monoterpene concentrations during the day. However, at both stations the sCIs from the monoterpene ozonolysis are larger by a factor 3–5 compared to CH₂OO. The important observation from this figure is the trend between the two plotted variables. With increasing concentrations of the stabilized Criegee intermediates, we see a larger number of sulfuric acid molecules which cannot be predicted using the model chemistry from the MCM with unchanged reaction rates for the oxidation of sulfur dioxide with the sCIs. This linear dependency can be observed at both sta-




Fig. 4. Scatter plot of organic vapours against the missing sulfuric acid concentrations for the selected periods at both stations.

tions and could indicate the possible participation of the sCIs in SO₂ oxidation.

The next step in our study was to run the model with the different chemistry scenarios explained in Sect. 3.3 to predict the contributions of each different sulfur dioxide oxidation mechanism. Table 2 shows the mean measured and modelled sulfuric acid concentrations together with the correlation coefficient and the coefficient of variation of the root-mean-square error (CV(RMSE)) between the measurements and simulations for each scenario and station. In Table 3 we present the percentage contribution of the different reactions schemes described in Sect. 3.3 in relation to the measured concentrations. If we compare the measured and modelled sulfuric acid concentrations at each station, we realize that for both locations the model underestimates

Table 2. Statistical analysis between measured and modelled sulfuric acid concentrations for Hyttiälä and Hohenpeissenberg (RMSE – root mean square error).

	Hyttiälä				Hohenpeissenberg			
	Mean meas.	Mean model	Corr. coef.	CV(RMSE) (%)	Mean meas.	Mean model	Corr. coef.	CV(RMSE) (%)
Scenario A	1.2 E6	3.3 E5	0.98	137.4	1.3 E6	7.4 E5	0.91	47.1
Scenario B	1.2 E6	4.3 E5	0.96	119.9	1.3 E6	8.3 E5	0.92	41.8
Scenario C	1.2 E6	7.6 E5	0.95	77.4	1.3 E6	1.2 E6	0.89	32.2
Scenario D	1.2 E6	8.2 E5	0.93	72.3	1.3 E6	1.2 E6	0.89	32.2

Table 3. Contribution of the different chemical paths to form sulfuric acid in the atmosphere.

	Hyttiälä	Hohenpeissenberg
Inorganic chemistry (scenario A)	28.9 %	49.5 %
Organic chemistry from MCM (scenario B)	9.1 %	10.2 %
sCIs from Monoterpenes with increased reaction rate (scenario C)	31.7 %	38.1 %
sCIs from CH ₂ O with increased reaction rate (scenario D)	5.8 %	2.9 %
Underestimation by scenario D	24.4 %	−0.6 %

the measurements by approximately a factor of 2 using the chemistry from the Master Chemical Mechanism as explained under scenario B. If we consider the new reaction rates for the sCIs resulting from the ozonolysis of the monoterpenes (scenario C), we increase calculated H₂SO₄ concentrations between 30–40 %. By also including the new reaction rate from formaldehyde oxide as in scenario D, we increase the simulated sulfuric acid concentrations for both stations by 3–6 %. Adding all the different factors together, we find the model underestimates measured concentrations of H₂SO₄ in Hyttiälä by 24 % and shows a perfect agreement for Hohenpeissenberg.

All scenarios show a very high correlation between the modelled and measured values, with $R \geq 0.88$. This correlation shows that there is a similar trend in the two data sets, but it gives no information about the difference between both data sets. For this reason we calculated the RMSE (sometimes also called root-mean-square deviation – RMSD) to measure the difference between our predicted and measured sulfuric acid concentrations. At the last step we normalized these values with the mean measured concentrations, and we present the values in percentage in Table 2. It is obvious that scenarios C and D have the lowest CV(RMSE) values com-

pared to all other runs and that the root-mean-square error is reasonably smaller at Hohenpeissenberg.

Figures 5 and 6 show the measured and modelled (scenarios B and D) sulfuric acid concentrations for Hyttiälä and Hohenpeissenberg, respectively. Included in the figures are uncertainty simulations for scenario B for the measured OH-concentration with $\pm 30\%$. For Hyttiälä, it is obvious that the inclusions of the new reaction rates for the sCIs nearly always dramatically improve the simulated sulfuric acid concentrations. Only at the end of our selected period does the model overestimate the measurements continuously for several hours. At Hohenpeissenberg the situation differs for the three different periods. During the first two days, the difference between the measured sulfuric acid concentrations and the one modelled without using the improved reaction rates for the sCIs is very small. However, the concentrations of the monoterpenes and isoprene are reasonably smaller, thus the contributions of the new oxidation path for sulfur dioxide is very small. If we now consider the last three days in this plot, we see a much greater difference between measured and modelled concentrations for scenario B. This difference becomes smaller when we use the new reaction rates of the sCIs on all days. The contribution of these mechanisms seems to be crucial for Hohenpeissenberg, especially during the night-time when the isoprene and monoterpene concentrations reach a certain value. Scatter plots for both stations of the measured and modelled (scenario D) H₂SO₄ concentrations, presented in Fig. 7, show that for both stations most of the values are in the 50 % interval, which could be explained by taking the uncertainty of all the measurements into account. During time periods with sulfuric acid concentrations below 3×10^5 molecules cm^{−3}, the model still seems to underestimate compared to the measured concentrations. This divergence could be related to the fact that we only changed the rates for four reactions of stabilized Criegee intermediates for our model investigations and very likely many more exist in the real atmosphere, which are not currently included. Another possible explanation which we will not discuss further in this manuscript but should be mentioned is the low concentrations of OH radicals during night-times and possibly an increase in the uncertainties of the instrument at these low concentrations.




Fig. 5. Measured and modelled sulfuric acid concentrations for Hyttiälä; the definitions of the different model scenarios are presented in Sect. 3.3.




Fig. 6. Measured and modelled sulfuric acid concentrations for Hohenpeissenberg; the definitions of the different model scenarios are presented in Sect. 3.3.

Figures 8 and 9 give the contributions, as a percentage, of the different chemical pathways for the entire periods at the two stations. The “H₂SO₄ by SO₂ and Cls” data were calculated by first running the model with scenario D and then subtracting the values predicted by a run with scenario B. In this case only the difference between the new and old reaction rate constants for the sCIs are considered as a source term for the concentrations of sulfuric acid. The difference in the contributions at the two stations is remarkable. While in Hyttiälä both mechanisms seem to be active at all times with nearly similar contributions, the situation appears completely different at Hohenpeissenberg. Here we recognise a clear dominance of the OH path during daytime and the converse at nights when the sCIs contribute between 50 and 80 % depending on the precursor concentrations. As mentioned in Sect. 2, the meteorological situation of the summer 2010 in Hyttiälä was very exceptional during the HUMPPA-COPEC campaign and would not be considered the normal situation for this station. In the next subsection we will show that there




Fig. 7. Scatter plot of measured against modelled sulfuric acid concentrations for the selected periods at both stations based on simulation with scenario D (for the definition of the scenarios, see Sect. 3.3); black thick lines indicate the 1 : 1 range and black thin lines the 50 % uncertainty interval.




Fig. 8. Cumulative percentage contribution of the different SO₂ oxidation mechanisms for Hyttiälä.

is also in Hyttiälä a clear trend of OH-generated H₂SO₄ during daytime and sCI-generated H₂SO₄ at night. The selected period shown for Hyttiälä was abnormal and gave a good opportunity to test the new chemical paths under extreme conditions and investigate if the results are still valid.

4.3 Vertical seasonal impact at Hyttiälä

In this last section we will investigate the seasonal impact of the oxidation of SO₂ by sCIs on the sulfuric acid concentrations in the lower part of the troposphere at Hyttiälä during 2010 using the chemistry transport model SOSA. In contrast with the model MALTE in the previous subsection where measured values were used, OH and VOCs will now be calculated online and not taken from measurements. This change is necessary because no long-term (over a season or year) vertical measurements for these compounds exist, but are necessary to do this type of study.




Fig. 9. Cumulative percentage contribution of the different SO₂ oxidation mechanisms for Hohenpeissenberg.

After the run we calculated for every time step and for every day in 2010 the H₂SO₄ production ratio. We define the H₂SO₄ production ratio as the ratio between H₂SO₄ produced only by OH oxidation of SO₂ divided by H₂SO₄ produced both by OH oxidation of SO₂ and also by reactions of stabilized Criegee intermediates with SO₂. We can write this ratio, *R*, as

$$R = \frac{[\text{H}_2\text{SO}_4]_{\text{OH}}}{[\text{H}_2\text{SO}_4]_{\text{OH}} + [\text{H}_2\text{SO}_4]_{\text{sCI}}} \quad (5)$$

This ratio is calculated for every height layer and averaged seasonally over the year 2010 with spring season being March, April and May; summer being June, July and August; autumn being September, October and November; and winter being December, January and February.

Figure 10 presents the sulfuric acid production ratio in the lowest 500 m for all seasons at the SMEAR II station at Hyytiälä and Table 4 gives the average values for different height levels from the ground to the height mentioned in the first column. In spring, summer and autumn, the situations are quite similar, with a contribution from the stabilized Criegee intermediates to the total sulfuric acid production decreasing from 15–20 % in the lowest 20 m to 5–7 % when considering the first 500 m. However, nearly double the contribution of the sCIs is observed during winter due to the low solar irradiance at this high latitude station during the months of December to February. One remarkable result from the plots in Fig. 10 is the expected daily distribution with relatively low sulfuric acid production ratios during night-time. As already mentioned above, this observation reflects exactly the monoterpene concentrations measured at this station in earlier studies (Mogensen et al., 2011) and stands in contrast with the terpene profiles recorded during the HUMPPA-COPEC campaign.

The last figure of our study (Fig. 11) shows the sulfuric acid production ratio for two different height intervals (500 m and 50 m) averaged over the whole year of 2010 at the SMEAR II station. In the first plot we see that up to a

Table 4. Seasonal mean values of the sulfuric reaction ratio averaged from the ground up to different heights calculated according to Eq. (5).

Height	Season			
	Spring	Summer	Autumn	Winter
22 m	0.85	0.86	0.80	0.61
56 m	0.86	0.87	0.81	0.63
108 m	0.88	0.88	0.83	0.70
225 m	0.92	0.91	0.88	0.80
505 m	0.94	0.93	0.93	0.89
2001 m	0.97	0.96	0.95	0.95

height of 100 m the average daily contribution of the sCIs can be around 19 %, with higher contributions during night-time. Above this height the OH production term dominates, and the ozonolysis of the monoterpene contributes only with a very small fraction to the total sulfuric acid production. The second plot also shows one shortcoming of the model SOSA, or rather the emission module used, which is that it currently does not include any parameterization of VOC emissions from the soil or floor of the canopy. For this reason the H₂SO₄ production ratio increased quite strongly in the lowest 3 m compared to the upper part of the canopy. Overall, our results show that the effect of sCIs can have a significant impact in the lowest part of the troposphere (first 100 m) when calculating the sulfuric acid concentrations. Considering that most of the H₂SO₄ measurements are performed at ground level, the inclusion of this new oxidation mechanism is crucial. However, when we consider the vertical dependence over the SMEAR II station (up to 2 km), the model shows a decreasing importance with only some percent of the produced sulfuric acid concentrations (5 % up to 2 km) originating from the reaction of sulfur dioxide with stable Criegee intermediates.

5 Uncertainties

The aerosol condensational sink (CS) determines how rapidly molecules will condense onto pre-existing aerosols (Kulmala et al., 2001b). In these calculations we use measured particle number concentrations from the smallest sizes at around 3 nm up to several μm. However, the uncertainty of the predicted CS values due to potentially different hygroscopic growth behaviour depending on the chemical composition of the particles is difficult to estimate and could have a significant effect for the simulated sulfuric acid concentrations.

The Criegee chemistry has been investigated both experimentally as well as theoretically, but is still somewhat unexplored and leaves us with a lot to wish for concerning the chemical mechanisms, pathways, products, stabilizations and yields, thermal lifetimes, pressure dependency and




Fig. 10. Seasonal average values over the first 500 m above the surface at SMEAR II for model simulations with the chemistry model from scenario B divided by scenario D (for the definition of the scenarios see Sect. 3.3). The colour bar gives the ratio as described in Eq. (5).




Fig. 11. Yearly average values for 50 and 500 m above the surface at SMEAR II for model simulations with the chemistry model from scenario B divided by scenario D (for the definition of the scenarios see Sect. 3.3). The colour bar gives the ratio as described in Eq. (5).

reaction rate coefficients. Donahue et al. (2011) gives a great general overview of the dependences on conditions in atmospheric Criegee chemistry. Below we will briefly touch those uncertainties that influence the results gained in the presented manuscript.

Firstly, it should be emphasised that CIs have not been directly observed in the gas phase until very recently, by Welz et al. (2012). Instead their presence, lifetime and reaction rate coefficients have been estimated based on indirect measurements, by investigating changes in the products of ozonolysis of specific alkenes upon addition of different reagents or scavengers (e.g. Alam et al., 2011; Presto and Donahue, 2004).

Chemical paths: The CI chemistry of simple alkenes is still not fully understood, but the situation gets still worse for more complex systems. For example, α -pinene and limonene (endocyclic terpenes) produce two excited CIs, both with

a carbonyl and a carbonyl oxide functional group. This structure gives the possibility for intra-molecular reactions, which is not well investigated. Furthermore, limonene also has an exocyclic double bond where we would also expect an ozone addition (Leungsakul et al., 2005; Jiang et al., 2010); however, this is not included in the MCM.

Stabilization: In order to react with other molecules, the CI must be collisionally stabilized. MCM assumes that one stable CI is formed for each alkene (including dienes). Browsing through the literature, one finds that there also seems to be some disagreement concerning the yields of the sCI produced from various alkenes, e.g. for α -pinene yields of 0.34 (Zhang and Zhang, 2005) and 0.15 (Drozd and Donahue, 2011) have been reported, while for β -pinene the yields are given as 0.37 (Nguyen et al., 2009) and 0.22 (Zhang and Zhang, 2005).

Thermal lifetime: After stabilization of the CI, the CI must have a long enough thermal lifetime in order to react with other molecules. Again, direct measurements of the lifetimes of the stabilized intermediates have not been possible so far, and reported values span orders of magnitudes (e.g. Welz et al., 2012; Olzmann et al., 1997).

Reaction rate coefficients: The stabilized CI is assumed to react with SO₂, CO, NO, NO₂ and H₂O. Direct measurements, indirect determinations via measurements and computational calculations of specific CIs reaction rate coefficients with these various compounds span over several orders of magnitudes and must be thought to be extremely uncertain (e.g. Welz et al., 2012; Mauldin III et al., 2012; Johnson and Marston, 2008; Hatakeyama and Akimoto, 1994; Hatakeyama et al., 1986; Kurtén et al., 2011). Measurements of the reaction rate coefficients are often done under low pressure due to practical issues. When applying reaction rate coefficients measured under low pressure conditions, one has to bear in mind that reaction rate coefficients

Table 5. Sensitivity studies on the different reaction rates between the stabilized Criegee intermediates and water on the overall sCI contribution to form sulfuric acid in the atmosphere. Explanations of the different scenarios are provided in Sects. 3 and 6.

	Hyytiälä	Hohenpeissenberg
Scenario D	75.6 %	100.6 %
Scenario D1	39.8 %	63.4 %
Scenario D2	35.3 %	54.7 %
Scenario D3	100.2 %	134.6 %
Scenario D4	227.6 %	231.1 %

under tropospheric conditions can be pressure and temperature dependent, reactions are commonly affected by the presence of water, and if the reactions have a substantial stabilization component, the rate constants could be substantially larger at atmospheric pressure (Welz et al., 2012). In our study we have only changed the reaction rate coefficients for reaction between CI and SO₂ for the Criegee intermediates resulting from the ozonolysis of monoterpenes. In the case of formaldehyde oxide, we also implemented the recommended reaction rates for NO₂ and H₂O by Welz et al. (2012). However, the large uncertainty on the CI reaction rate coefficients is not only thought to be for reactions with SO₂, but must also be expected for reaction with other trace gases.

6 Sensitivity study on the water channel

One of the highest uncertainties concerning the lifetimes of the stabilized Criegee intermediates relates to the value of the rate constant for the sCI with water. Hatakeyama and Aimoto (1994) reported that the rate constants for this reaction vary widely (2×10^{-19} to 1×10^{-15} cm³ s⁻¹). In our study, which is based on the work from Welz et al. (2012) and Mauldin et al. (2012), a very important difference has to be taken into account. The rates from Welz are based on direct measurements whereas Mauldin used an indirect method. In the second case no values for the reaction rate for sCI plus H₂O were given by the authors and the “overall” reaction rate included already a loss of the sCI by the reaction with water. This may be the reason for the significantly lower reaction rate between the Criegees and sulfur dioxide published by Mauldin compared to Welz. To investigate the effect of the water reaction rates with the stabilized Criegee intermediates, we performed sensitivity studies by changing these rates over two orders of magnitude. The rate constants in the MCM-Leeds for the sCI–H₂O reaction, which were used until now for the Criegees originating from the ozonolysis of the monoterpenes, are 1.4×10^{-17} cm³ s⁻¹ (α -pinene and limonene) and 6×10^{-18} cm³ s⁻¹ (β -pinene). In the case of formaldehyde oxide, we used the rate as mentioned before from Welz et al. (2012) of 4×10^{-15} cm³ s⁻¹. To show how crucial the impact of these rates are on the simulated

sulfuric acid concentrations, we increased the sCI–H₂O reaction rate for the monoterpene-originated sCIs by one and two orders of magnitude (D1 and D2, respectively) and then decreased the rate for formaldehyde oxide plus water by one and two orders of magnitude (D3 and D4, respectively). The results, presented in Table 5, show the contribution of the oxidation of sulfur dioxide by the sCI for the single cases when all the other rates kept constant as described in scenario D. The results point out that the knowledge of this reaction rate is crucial. An increase of the reaction rate for the sCIs from monoterpenes by 10 will decrease the overall formation rate of sulfuric acid down to 39.8 % and 63.4 % for Hyytiälä and Hohenpeissenberg, respectively, when compared to the measurements. On the other side an increase in the reaction rate for formaldehyde oxide by 10 would improve the situation for Hyytiälä but lead to an overestimation for Hohenpeissenberg by more than 34 %. One way to investigate the reaction rates for the sCI with water and with sulfur dioxide could be a very detailed long-term sensitivity study by taking the changes of the relative humidity as one factor to research when the best agreement between the simulations and measurements are achieved. This could improve our knowledge in the reaction rates for the Criegees with water and sulfur dioxide but will still leave a lot of uncertainties on the formation of the stabilized Criegee intermediates.

7 Conclusions

This study dramatically changes the current understanding of atmospheric sulfuric acid production: we have shown that oxidation of SO₂ by stabilized Criegee intermediate radicals can be a crucial source for atmospheric sulfuric acid production in VOC rich environments. Depending on the concentrations of several investigated organic compounds (reaction products from ozone oxidation of isoprene and monoterpenes), their contribution via the reaction of stabilized Criegee intermediates to atmospheric sulfuric acid gas phase concentrations could be as high as 46 % at ground level. Taking into account that most of the H₂SO₄ measurements are performed at ground level, the inclusion of this new oxidation mechanism could be crucial in regional as well as global models.

Our model investigations of the boundary layer in Hyytiälä showed that the contribution from the sCIs to sulfuric acid production is, as expected, most important in the canopy, where the concentration of organic compounds is highest. However, our overall results show that the effect of sCIs up to 100 m is very important to consider when calculating the sulfuric acid concentration.

We assume that other stabilized Criegee intermediate radicals produced from ozone oxidation of other unsaturated organic compounds exist and contribute to the production of sulfuric acid, and we urge for more investigation into this field. As a last remark we want to point to the high

uncertainties in calculating the concentrations of the stabilized Criegee intermediates and rate constants for the reactions of the sCI and other molecules. Until a complete knowledge of these parameters has been achieved, a final statement about the quantitative effects Criegee radicals have on the atmospheric chemistry is not possible.

Acknowledgements. We thank the HUMPPA-COPEC-2010 team for providing and measuring gas concentrations used in this paper. Furthermore, financial support from Cryosphere-Atmosphere Interactions in a Changing Arctic Climate (CRAICC), the Helsinki University Centre for Environment (HENVI), the Academy of Finland (251427, 139656), PEGASOS project funded by the European Commission (FP7-ENV-2010-265148), and the Otto A. Malm foundation and computational resources from CSC – IT Center for Science Ltd. are all gratefully acknowledged. Harald Berresheim would like to thank Deutscher Wetterdienst and Science Foundation Ireland (grant 09/RFP/GEO2176) for support.

Edited by: D. Heard

References

- Alam, M. S., Camredon, M., Rickard, A. R., Carr, T., Wyche, K. P., Hornsby, K. E., Monks, P. S., and Bloss, W. J.: Total radical yields from tropospheric ethene ozonolysis, *Phys. Chem. Chem. Phys.*, 13, 11002–11015, 2011.
- Atkinson, R.: Gas-phase tropospheric chemistry of organic compounds, *J. Phys. Chem. Ref. Data, Monograph*, 2, 1–216, 1994.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- Bäck, J., Aalto, J., Henriksson, M., Hakola, H., He, Q., and Boy, M.: Chemodiversity of a Scots pine stand and implications for terpene air concentrations, *Biogeosciences*, 9, 689–702, doi:10.5194/bg-9-689-2012, 2012.
- Berresheim, H., Elste, T., Plass-Dülmer, C., Eisele, F. L., and Tanner, D. J.: Chemical ionization mass spectrometer for long-term measurements of atmospheric OH and H₂SO₄, *Int. J. Mass Spectrom.*, 202, 91–109, 2000.
- Birmili, W., Berresheim, H., Plass-Dülmer, C., Elste, T., Gilge, S., Wiedensohler, A., and Uhrner, U.: The Hohenpeissenberg aerosol formation experiment (HAFEX): a long-term study including size-resolved aerosol, H₂SO₄, OH, and monoterpene measurements, *Atmos. Chem. Phys.*, 3, 361–376, doi:10.5194/acp-3-361-2003, 2003.
- Boy, M. and Kulmala, M.: The part of the solar spectrum with the highest influence on the formation of SOA in the continental boundary layer, *Atmos. Chem. Phys.*, 2, 375–386, doi:10.5194/acp-2-375-2002, 2002.
- Boy, M., Rannik, U., Lehtinen, K., Tarvainen, V., Hakola, H., and Kulmala, M.: Nucleation events in the continental boundary layer: long-term statistical analysis of aerosol relevant characteristics, *J. Geophys. Res.*, 108, 4667–4679, 2003.
- Boy, M., Kulmala, M., Ruuskanen, T. M., Pihlatie, M., Reissell, A., Aalto, P. P., Keronen, P., Dal Maso, M., Hellen, H., Hakola, H., Jansson, R., Hanke, M., and Arnold, F.: Sulphuric acid closure and contribution to nucleation mode particle growth, *Atmos. Chem. Phys.*, 5, 863–878, doi:10.5194/acp-5-863-2005, 2005.
- Boy, M., Hellmuth, O., Korhonen, H., Nilsson, E. D., ReVelle, D., Turnipseed, A., Arnold, F., and Kulmala, M.: MALTE – model to predict new aerosol formation in the lower troposphere, *Atmos. Chem. Phys.*, 6, 4499–4517, doi:10.5194/acp-6-4499-2006, 2006.
- Boy, M., Kazil, J., Lovejoy, E. R., Korhonen, H., Guenther, A., and Kulmala, M.: Relevance of ion-induced nucleation of sulphuric acid and water in the lower troposphere over the boreal forest at northern latitudes, *Atmos. Res.*, 90, 151–158, doi:10.1016/j.atmosres.2008.01.002, 2008.
- Boy, M., Sogachev, A., Lauros, J., Zhou, L., Guenther, A., and Smolander, S.: SOSA – a new model to simulate the concentrations of organic vapours and sulphuric acid inside the ABL – Part I: Model description and initial evaluation, *Atmos. Chem. Phys.*, 11, 43–51, doi:10.5194/acp-11-43-2011, 2011.
- Calvert, J., Atkinson, R., Kerr, J. A., Madronich, S., Moortgat, G. K., Wallington, T. J., and Yarwood, G.: *The Mechanisms of Atmospheric Oxidation of the Alkenes*, Oxford University Press, Oxford, UK, 2000.
- Damian, V., Sandu, A., Damian, M., Potra, F., and Carmichael, G. R.: The kinetic preprocessor kpp – a software environment for solving chemical kinetics, *Comput. Chem. Eng.*, 26, 1567–1579, 2002.
- De Gouw, J. and Warneke, C.: Measurements of volatile organic compounds in the Earth's atmosphere using proton-transfer-reaction mass spectrometry, *Mass Spectrom. Rev.*, 26, 223–257, 2007.
- Donahue, N. M., Drozd, G. T., Epstein, S. A., Presto, A. A., and Kroll, J. H.: Adventures in ozoneland: down the rabbit-hole, *Phys. Chem. Chem. Phys.*, 13, 10848–10857, 2011.
- Drozd, G. T. and Donahue, N. M.: Pressure dependence of stabilized Criegee intermediate formation from a sequence of alkenes, *J. Phys. Chem. A*, 115, 4381–4387, 2011.
- Eisele, F. L. and Tanner, D. J.: Measurement of the gas phase concentration of H₂SO₄ and methane sulfonic acid and estimates of H₂SO₄ production and loss in the atmosphere, *J. Geophys. Res.*, 98, 9001–9010, 1993.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- Hakola, H., Tarvainen, V., Bäck, J., Ranta, H., Bonn, B., Rinne, J., and Kulmala, M.: Seasonal variation of mono- and sesquiterpene emission rates of Scots pine, *Biogeosciences*, 3, 93–101, doi:10.5194/bg-3-93-2006, 2006.
- Hari, P. and Kulmala, M.: Station for Measuring Ecosystem–Atmosphere Relations (SMEAR II), *Boreal Environ. Res.*, 10, 315–322, 2005.
- Hatakeyama, S. and Akimoto, H.: Reactions of Criegee Intermediates in the gas phase, *Res. Chem. Intermed.*, 20, 503–524, 1994.
- Hatakeyama, S., Kobayashi, H., Lin, Z. Y., Tagaki, H., and Akimoto, H.: Mechanism for the reaction of peroxyethylene with sulfur dioxide, *J. Phys. Chem.*, 90, 4131–4135, 1986.

- Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, 1997.
- Jiang, L., Xu, Y.-S., and Ding, A.-Z.: Reaction of stabilized criegee intermediates from ozonolysis of limonene with sulfur dioxide: ab Initio and DFT study, *J. Phys. Chem. A*, 114, 12452–12461, 2010.
- Johnson, D. and Marston, G.: The gas-phase ozonolysis of unsaturated volatile organic compounds in the troposphere, *Chem. Soc. Rev.*, 37, 699–716, 2008.
- Kirkby, J. and CLOUD-team: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–423, 2011.
- Kulmala, M. and Pirjola, L.: Stable sulphate clusters as a source of new atmospheric particles, *Nature*, 404, 66–69, 2000.
- Kulmala, M., Haemeri, K. K., Aalto, P., Maekelae, J., Pirjola, L., Nilsson, E. D., Buzorius, G., Rannik, U., Dal Maso, M., Seidl, W., Hoffmann, T., Jansson, R., Hansson, H.-C., O'Dowd, C., and Viisanen, Y.: Overview of the international project on biogenic aerosol formation in the boreal forest (BIOFOR), *Tellus B*, 53, 324–343, 2001a.
- Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen, P., Hämeri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus B*, 53, 479–490, 2001b.
- Kurtén, T., Lane, J. R., Jørgensen, S., and Kjaergaard, H. G.: A computational study of the oxidation of SO₂ to SO₃ by gas-phase organic oxidants, *J. Phys. Chem. A*, 115, 8669–8681, 2011.
- Lauros, J., Sogachev, A., Smolander, S., Vuollekoski, H., Sihto, S.-L., Mammarella, I., Laakso, L., Rannik, Ü., and Boy, M.: Particle concentration and flux dynamics in the atmospheric boundary layer as the indicator of formation mechanism, *Atmos. Chem. Phys.*, 11, 5591–5601, doi:10.5194/acp-11-5591-2011, 2011.
- Leungsakul, S., Jaoui, M., and Kamens, R. M.: Kinetic mechanism for predicting secondary organic aerosol formation from the reaction of d-limonene with ozone, *Environ. Sci. Technol.*, 39, 9583–9594, 2005.
- Makkonen, R., Asmi, A., Korhonen, H., Kokkola, H., Järvenoja, S., Räisänen, P., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., Järvinen, H., Lohmann, U., Bennartz, R., Feichter, J., and Kulmala, M.: Sensitivity of aerosol concentrations and cloud properties to nucleation and secondary organic distribution in ECHAM5-HAM global circulation model, *Atmos. Chem. Phys.*, 9, 1747–1766, doi:10.5194/acp-9-1747-2009, 2009.
- Makkonen, R., Asmi, A., Kerminen, V.-M., Boy, M., Arneth, A., Hari, P., and Kulmala, M.: Air pollution control and decreasing new particle formation lead to strong climate warming, *Atmos. Chem. Phys.*, 12, 1515–1524, doi:10.5194/acp-12-1515-2012, 2012.
- Mauldin III, R. L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T., Stratmann, F., Kerminen, V.-M., and Kulmala, M.: A new atmospherically relevant oxidant, *Nature*, 488, 193–196, doi:10.1038/nature11278, 2012.
- Merikanto, J., Spracklen, D. V., Pringle, K. J., and Carslaw, K. S.: Effects of boundary layer particle formation on cloud droplet number and changes in cloud albedo from 1850 to 2000, *Atmos. Chem. Phys.*, 10, 695–705, doi:10.5194/acp-10-695-2010, 2010.
- Mogensen, D., Smolander, S., Sogachev, A., Zhou, L., Sinha, V., Guenther, A., Williams, J., Nieminen, T., Kajos, M. K., Rinne, J., Kulmala, M., and Boy, M.: Modelling atmospheric OH-reactivity in a boreal forest ecosystem, *Atmos. Chem. Phys.*, 11, 9709–9719, doi:10.5194/acp-11-9709-2011, 2011.
- Nguyen, T. L., Peeters, J., and Vereecken, L.: Theoretical study of the gas-phase ozonolysis of beta-pinene (C₁₀H₁₆), *Phys. Chem. Chem. Phys.*, 11, 5643–5656, 2009.
- Olzmann, M., Kraka, E., Cremer, D., Gutbrod, R., and Anderson, S.: Energetics, kinetics, and product distributions of the reactions of ozone with ethene and 2,3-Dimethyl-2-butene, *J. Phys. Chem. A*, 101, 9421–9429, 1997.
- Paasonen, P., Nieminen, T., Asmi, E., Manninen, H. E., Petäjä, T., Plass-Dülmer, C., Flentje, H., Birmili, W., Wiedensohler, A., Hörrak, U., Metzger, A., Hamed, A., Laaksonen, A., Facchini, M. C., Kerminen, V.-M., and Kulmala, M.: On the roles of sulphuric acid and low-volatility organic vapours in the initial steps of atmospheric new particle formation, *Atmos. Chem. Phys.*, 10, 11223–11242, doi:10.5194/acp-10-11223-2010, 2010.
- Petäjä, T., Mauldin, III, R. L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, *Atmos. Chem. Phys.*, 9, 7435–7448, doi:10.5194/acp-9-7435-2009, 2009.
- Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation and primary emission rates, *Atmos. Chem. Phys.*, 9, 1339–1356, doi:10.5194/acp-9-1339-2009, 2009.
- Plass-Dülmer, C. and Berresheim, H.: Volatile Organic Compound Measurements at Hohenpeissenberg as part of GAW, in “The German Contribution to the WMO/GAW Program: Upon the 225th anniversary of GAW Hohenpeissenberg Observatory”, edited by: Fricke, W., GAW Report 167, (WMO TD No. 1336), Deutscher Wetterdienst, 2007.
- Presto, A. A. and Donahue, N. M.: Ozonolysis fragment quenching by nitrate formation: the pressure dependence of prompt OH radical formation, *J. Phys. Chem. A*, 108, 9096–9104, 2004.
- Rohrer, F. and Berresheim, H.: Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet radiation, *Nature*, 442, 184–187, 2006.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, doi:10.5194/acp-3-161-2003, 2003.
- Sipilä, M., Berndt, T., Petäjä, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin III, R. L., Hyvärinen, H., Lihavainen, H., and Kulmala, M.: The role of sulphuric acid in atmospheric nucleation, *Science*, 327, 1243–1246, 2010.
- Sogachev, A.: A note on two-equation closure modelling of canopy flow, *Bound.-Lay. Meteorol.*, 130, 423–435, 2009.
- Sogachev, A. and Panferov, O.: Modification of two-equation models to account for plant drag, *Bound.-Lay. Meteorol.*, 121, 229–266, 2006.
- Sogachev, A., Menzhulin, G., Heimann, M., and Lloyd, J.: A simple three dimensional canopy planetary boundary layer simulation model for scalar concentrations and fluxes, *Tellus B*, 54, 784–819, 2002.
- Sogachev, A., Panferov, O., Gravenhorst, G., and Vesala, T.: Numerical analysis of flux footprints for different landscapes, *Theor. Appl. Climatol.*, 80, 169–185, 2005.

- Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V. M., Sihto, S. L., Riipinen, I., Merikanto, J., Mann, G. W., Chipperfield, M. P., Wiedensohler, A., Birmili, W., and Lihavainen, H.: The contribution of particle formation to global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 35, L06808, doi:10.1029/2007GL033038, 2008.
- Wang, M. and Penner, J. E.: Aerosol indirect forcing in a global model with particle nucleation, *Atmos. Chem. Phys.*, 9, 239–260, doi:10.5194/acp-9-239-2009, 2009.
- Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measurements of new particle formation and ultrafine particle growth rates at a clean continental site, *J. Geophys. Res.*, 102, 4375–4385, 1997.
- Welz, O., Savee, J. D., Osborn, D. L., Vasu, S. S., Percival, C. J., Shallcross, D. E., and Taatjes, C. A.: Direct kinetic measurements of Criegee Intermediate (CH₂OO) formed by reaction of CH₂I with O₂, *Science*, 335, 204–207, 2012.
- Williams, J., Crowley, J., Fischer, H., Harder, H., Martinez, M., Petäjä, T., Rinne, J., Bäck, J., Boy, M., Dal Maso, M., Hakala, J., Kajos, M., Keronen, P., Rantala, P., Aalto, J., Aaltonen, H., Paatero, J., Vesala, T., Hakola, H., Levula, J., Pohja, T., Herrmann, F., Auld, J., Mesarchaki, E., Song, W., Yassaa, N., Nölscher, A., Johnson, A. M., Custer, T., Sinha, V., Thieser, J., Pouvesle, N., Taraborrelli, D., Tang, M. J., Bozem, H., Hosaynali-Beygi, Z., Axinte, R., Oswald, R., Novelli, A., Kubistin, D., Hens, K., Javed, U., Trawny, K., Breitenberger, C., Hidalgo, P. J., Ebben, C. J., Geiger, F. M., Corrigan, A. L., Russell, L. M., Ouwersloot, H. G., Vilà-Guerau de Arellano, J., Ganzeveld, L., Vogel, A., Beck, M., Bayerle, A., Kampf, C. J., Bertelmann, M., Köllner, F., Hoffmann, T., Valverde, J., González, D., Riekkola, M.-L., Kulmala, M., and Lelieveld, J.: The summertime Boreal forest field measurement intensive (HUMPPA-COPEC-2010): an overview of meteorological and chemical influences, *Atmos. Chem. Phys.*, 11, 10599–10618, doi:10.5194/acp-11-10599-2011, 2011.
- Zhang, D. and Zhang, R.: Ozonolysis of α -pinene and β -pinene: kinetics and mechanism, *J. Chem. Phys.*, 122, 114308, doi:10.1063/1.1862616, 2005.
- Zhao, J., Eisele, F. L., Titcombe, M., Kuang, C., and McMurry, P. H.: Chemical ionization mass spectrometric measurements of atmospheric neutral clusters using the cluster-CIMS, *J. Geophys. Res.*, 115, D08205, doi:10.1029/2009JD012606, 2010.