



The fractionation of nitrogen and oxygen isotopes in macroalgae during the assimilation of nitrate

P. K. Swart¹, S. Evans^{1,*}, T. Capo², and M. A. Altabet³

¹Division of Marine Geology and Geophysics, Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149, USA

²Division of Marine Biology and Fisheries, Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149, USA

³School for Marine Science and Technology, University of Massachusetts Dartmouth, New Bedford, MA 02744, USA

* now at: Department of Geosciences, Boise State University, 1910 University Drive, Boise, Idaho 83725-1535, USA

Correspondence to: P. K. Swart (pswart@rsmas.miami.edu)

Received: 27 March 2014 – Published in Biogeosciences Discuss.: 16 May 2014

Revised: 9 September 2014 – Accepted: 30 September 2014 – Published: 13 November 2014

Abstract. In order to determine and understand the stable isotope fractionation of ^{18}O and ^{15}N manifested during assimilation of NO_3^- in marine macro-benthic algae, two species (*Ulva* sp. and *Agardhiella* sp.) have been grown in a wide range of NO_3^- concentrations (2–500 μM). Two types of experiments were performed. The first was one in which the concentration of the NO_3^- was allowed to drift downward as it was assimilated by the algae, between 24 hour replacements of media. These experiments proceeded for periods of between 7 and 10 days. A second set of experiments maintained the NO_3^- concentration at a low steady-state value by means of a syringe pump. The effective fractionation during the assimilation of the NO_3^- was determined by measuring the $\delta^{15}\text{N}$ of both the (i) new algal growth and (ii) residual NO_3^- in the free-drift experiments after 0, 12, 24 and 48 h. Modelling these data show that the fractionation during assimilation is dependent upon the concentration of NO_3^- and is effectively 0 at concentrations of less than $\sim 2 \mu\text{M}$. The change in the fractionation with respect to concentration is the greatest at lower concentrations (2–10 μM). The fractionation stabilizes between 4 and 6 ‰ at concentrations of between 50 and 500 μM . Although the $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ values of NO_3^- in the residual solution were correlated, the slope of relationship also varied with respect to NO_3^- concentration, with slopes of greater than unity at low concentration. These results suggest shifts in the dominant fractionation mechanism of ^{15}N and ^{18}O between concentrations of 1 and 10 μM NO_3^- . At higher NO_3^- concentrations (> 10–50 μM), frac-

tionation during assimilation will lead to $\delta^{15}\text{N}$ values in algal biomass lower than the ambient NO_3^- and ^{15}N enrichments in the residual NO_3^- .

1 Introduction

Nitrogen availability is an important factor in controlling algal growth in marine environments, representing a limiting nutrient throughout much of the global ocean (Dugdale and Wilkerson, 1986). In many studies, information on nitrogen sources and its cycling has been obtained by examining the ratio of the stable isotopes of nitrogen (^{14}N and ^{15}N) as well as oxygen (^{18}O and ^{16}O) in the case of NO_3^- . Isotope ratios are expressed using the conventional “delta” notation ($\delta^{15}\text{N}$ or $\delta^{18}\text{O}$) in parts per thousand (‰) deviation from the atmospheric N_2 standard or, in the case of oxygen, from Vienna standard mean ocean water (VSMOW). During cycling of NO_3^- , isotope fractionation takes place, as quantified by the associated fractionation factor (α). For algal NO_3^- uptake, α can be calculated using Eq. (1).

The term epsilon (ϵ) is also commonly used and is related to α by Eq. (2).

$$\alpha = \frac{\frac{^{15}\text{N}}{^{14}\text{N}}_{\text{NO}_3^-}}{\frac{^{15}\text{N}}{^{14}\text{N}}_{\text{algae}}} \quad (1)$$

$$\epsilon = (\alpha - 1) \times 1000 \quad (2)$$

The term ϵ can refer to fractionation of either ^{15}N ($^{15}\epsilon$) or ^{18}O ($^{18}\epsilon$) relative to the more abundant isotope of the element. In some of these processes, such as the fixation of atmospheric nitrogen, no significant isotopic fractionation takes place ($^{15}\epsilon \sim 0.0\%$) (Hoering and Ford, 1960), and consequently the $\delta^{15}\text{N}$ of N_2 fixing organisms is similar to that of atmospheric N_2 (0% by convention). In other processes, such as the denitrification of NO_3^- , $^{15}\epsilon$ values are higher than 20% (Barford et al., 1999; Delwiche and Steyn, 1970; Granger et al., 2006; Miyake and Wada, 1971), leading to large increases in the $\delta^{15}\text{N}$ of the residual reservoir of NO_3^- . While the $\delta^{15}\text{N}$ of microalgae has been studied in order to understand its use as a paleoceanographic proxy (Altabet, 1989; Altabet et al., 1991; Haug et al., 1998; Sigman et al., 2003), variations in the $\delta^{15}\text{N}$ of macroalgae have also been widely used as possible indicators of anthropogenic influences (Carballeira et al., 2013; Costanzo et al., 2001; Heaton, 1986). Generally speaking, nitrogen derived from sewage is isotopically enriched in ^{15}N , and it has been argued that even modest enrichments of ^{15}N in macroalgae might reflect enhanced input from such sources (Lapointe et al., 2004). Other studies have shown that such enrichments could occur through normal processes including fractionation during assimilation (Lamb et al., 2012; Stokes et al., 2011) and that there are not always simple relationships between the input of anthropogenic wastes and $\delta^{15}\text{N}$ values (Viana and Bode, 2013).

Studies of isotope fractionation during the assimilation of dissolved inorganic nitrogen by marine microalgae have reported a wide range of values. In one study, reported $^{15}\epsilon$ values ranged from 0.7 to 23% for the assimilation of NO_3^- by *Phaeodactylum tricorutum* (Wada and Hattori, 1978), a marine diatom. Another study reported $^{15}\epsilon$ values between 2.2 and 6.2% for 12 different marine phytoplankton cultures kept at a NO_3^- concentration of 100 μM (Needoba et al., 2003). Other research also reports wide ranges in $^{15}\epsilon$ values for both NO_3^- and NH_4^+ for a variety of different microalgae (Horrihan et al., 1990; Lajtha and Michener, 1994; Montoya et al., 1990; Wada and Hattori, 1978). At least part of these large ranges in $^{15}\epsilon$ values probably resulted from variations in experimental conditions and are perhaps artefacts resulting from differences in aeration, light and nutrient drawdown. In addition, changing nutrient concentration might be an important controlling parameter, and several studies have shown that microalgae show varying fractionation as a function of concentration (Hoch et al., 1992; Pennock et al., 1996; Waser et al., 1998) that is likely due to changes in physiology and perhaps uptake mechanism.

In contrast to microalgae, there have been relatively few studies of ^{15}N fractionation in macroalgae. Some of these studies have relied on spiking the natural environment with high nitrate and ammonium concentrations (Teichberg et al., 2007), while others have used transplant experiments (Deutsch and Voss, 2006). Neither of these investigations reported $^{15}\epsilon$ values for fractionation during the assimilation of NO_3^- . The study of Cohen and Fong (2005) grew the green algae *Enteromorpha intestinalis* under varying concentrations of NO_3^- and NH_4^+ , and, although they did not report values for ^{15}N fractionation, they concluded that the $\delta^{15}\text{N}$ of the algae was not dependent upon concentrations of dissolved inorganic nitrogen. These experiments used a combination of increases in NO_3^- and NH_4^+ , with the lower NO_3^- concentration experiments containing high amounts of NH_4^+ and vice versa. Under such experimental conditions it would have been difficult to isolate any potential concentration dependence upon fractionation manifested during assimilation. Given the possibility of a concentration dependence of ^{15}N fractionation for NO_3^- in microalgae, we revisit here whether such a dependency is found in macroalgae. We have used two different approaches over a range of different concentrations. In the first series of experiments, two species of macroalgae, *Ulva* sp. and *Agardhiella* sp., were grown over a range in nominal NO_3^- concentrations of 10, 50, 100 and 500 μM . As the algae within each culture consumed the NO_3^- in the solution, the solutions were replaced every 24 h. These were the so-called free-drift experiments. In the second set of experiments, NO_3^- levels were maintained at a low level (< 2 μM) by continual addition from a syringe pump. Hence these experiments cover the range of NO_3^- concentrations used in most previous experiments (> 100 μM) as well as those seen under natural conditions.

2 Methods

Samples of the green algae *Ulva* sp. and the rhodophyte algae *Agardhiella* sp. were collected from cultures held at the *Aplysia* Mariculture Laboratory's algal aquaculture facility (University of Miami). These species were maintained in a system of seven, 9000 L fiberglass tanks supplied with filtered seawater at a rate of $\sim 22 \text{ L min}^{-1}$. Radiant energy and temperature are monitored constantly, and algal growth rates are optimized by adjusting nutrient levels weekly. These stocks are kept continually as a food source for other organisms in the facility. In preparation for these experiments the algal thalli were rinsed with filtered seawater and gently scrubbed to remove surface epiphytes. Prior to experimentation, the macroalgae were maintained within 2 L flasks at 26 °C and approximately 100 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ for a 14-day acclimation period. During the acclimation period, filtered and autoclaved seawater was changed every 2 days, enriched to 500 $\mu\text{M N}$ (250 $\mu\text{M NaNO}_3$ and 250 $\mu\text{M NH}_4\text{Cl}$) and 44 $\mu\text{M KH}_2\text{PO}_4$, with $f/2$ medium supplements of B

Table 1. Changes in the $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ of NO_3^- during the *Ulva* experiments.

Time	<i>Ulva</i> sp.				<i>Agardiella</i> sp.			
	NO_3^-	Final	Final	f	NO_3^-	Final	Final	f
	μM	$\delta^{15}\text{N}$ ‰	$\delta^{18}\text{O}$ ‰		μM	$\delta^{15}\text{N}$ ‰	$\delta^{18}\text{O}$ ‰	
Tissue	14	5.1	–	0.26	14	1.8	–	0.21
0	14	3.3	21.8	1.00	14	3.5	17.3	1.00
12	9	4.5	22.4	0.60	3	6.3	23.1	0.22
24	4	4.8	24.9	0.26	3	6.4	22.6	0.21
48	< 1	Lost	Lost	0.03	–	–	–	–
Tissue	60	6.0	–	0.51	55	1.2	–	0.24
0	60	3.4	23.0	1.00	55	2.9	21.9	1.00
12	46	4.1	24.2	0.76	–	–	–	–
24	31	4.9	25.5	0.51	13	6.3	25.6	0.24
48	1	9.5	30.1	0.02	–	–	–	–
Tissue	103	2.9	–	0.78	104	0.7	–	0.68
0	103	3.2	23.6	1.00	104	3.1	23.3	1.00
12	90	3.5	24.1	0.87	–	–	–	–
24	90	4.1	24.4	0.78	71	5.1	25.1	0.68
48	36	6.3	27.6	0.35	–	–	–	–
Tissue	485	1.2	–	0.99	514	–3.0	–	0.96
0	485	3.2	23.2	1.00	514	2.7	23.1	1.00
12	467	3.2	23.5	0.96	–	–	–	–
24	481	2.7	23.1	0.99	495	3.1	23.3	0.96
48	435	3.4	23.7	0.90	439	4.1	25.0	0.85

For the tissue each analysis represents the mean of two replicates; for the NO_3^- each analysis represents the mean of four analyses.

Table 2. N isotopic composition of syringe experiment algae.

	NO_3^- (μM)	$\delta^{15}\text{N}$ ‰	σ	n
Initial		3.3	0.3	16
<i>Ulva</i>	3	1.3	0.3	7
<i>Agardiella</i>	7	3.4	0.0	2
<i>Agardiella</i>	10	2.9	0.0	2

vitamins (vitamin B₁₂, biotin and thiamine) and trace metals (Fe, Cu, Mo, Zn, Co and Mn) (Guillard, 1975). The cultures were continually aerated throughout the incubations.

2.1 Experimental protocol

2.1.1 Free-drift experiments

In these experiments the effect of varied nutrient availability on the nitrogen isotopic composition of new algal growth with respect to varied NO_3^- concentration was investigated. Nominal concentrations of 10, 50, 100 and 500 μM N (NaNO_3) were supplied in a medium of autoclaved, filtered (0.2 μm cartridge filter) seawater enriched with the

same KH_2PO_4 , B vitamin and trace metal supplements outlined for the acclimation medium (note that the actual targeted and measured concentrations were slightly different and the values used are reported in Tables 1 and 2). Sub-samples of *Ulva* and *Agardiella* (0.25–0.5 g wet weight; 2.5–3.0 cm) were taken from acclimation flasks, any visible epiphytes were again removed and the algae samples were placed in 2 L flasks filled with incubation medium. The media were replaced every 24 h, at which time each algal sample was rinsed to prevent epiphyte accumulation. The experiments proceeded for a period of 7–9 days. Water samples were collected after each 24 h period and analysed for the concentrations of NO_3^- and NH_4^+ . At the conclusion of the incubations, final accumulated biomass was weighed and, as the new algal growth produced was clearly visible, material which had grown only under the experimental conditions was trimmed off (Fig. 1). This material was dried (40 °C, 48 h) and then ground with mortar and pestle for subsequent N isotopic analyses and C / N determination. In order to examine the effect of assimilation on the $\delta^{15}\text{N}$ of residual NO_3^- , special experiments were performed in which the same water was kept in the algal cultures for periods of up to 48 h. After




Figure 1. Pictures showing samples of *Agardhiella* sp. grown in different concentrations of NO_3^- . From left to right, pictures show the initial individual and specimens grown in solutions containing nominally ambient, 10, 50, 100 and 500 μM NO_3^- . All experiments in which NO_3^- was added showed approximately similar growth rates, but reduced uptake of N at lower N concentrations. At the end of the experiment the ends of the algae were trimmed and analysed for their $\delta^{15}\text{N}$, $\delta^{13}\text{C}$ and C/N ratio. The new growth could be distinguished by comparison with the size of the original fragment (see Fig. 1a) and the change in colour.

12, 24 and 48 h, water samples were taken and the $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ of the NO_3^- measured.

2.1.2 Constant NO_3^- concentration experiments

At low concentrations of NO_3^- ($< 10 \mu\text{M}$), the algae rapidly assimilated NO_3^- and concentrations decreased to values of less than $3 \mu\text{M}$ within a few hours. In order to maintain a consistent low concentration and provide sufficient NO_3^- for the algal growth, NO_3^- was continuously added by means of a syringe pump. The rate of addition was initially determined by using the uptake rates calculated from the free-drift experiments and then adjusted slightly after the analysis of the NO_3^- concentration in the experiment. In these experiments concentrations started at $\sim 10 \mu\text{M}$ and stabilized at $3 \mu\text{M}$ throughout the growth period.




Figure 2. The relationship between $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ during the free-drift experiments for the two species of algae studied. Error bars represent mean analytical error for the various analyses and have been removed for clarity from the *Agardhiella* sp. data.

2.2 Analytical protocol

2.2.1 Stable isotopes

Algal biomass

The organic carbon and nitrogen content as well as the stable nitrogen ($\delta^{15}\text{N}$) and carbon ($\delta^{13}\text{C}$) isotopic composition of the algae was determined using a CN analyser (ANCA, Europa Scientific) interfaced with a continuous-flow isotope-ratio mass spectrometer (CF-IRMS) (20–20, Europa Scientific). Prior to analysis the algae samples were dried and 3–6 mg were placed in tin capsules. Data obtained from the mass spectrometer provide the C/N ratio of the samples in addition to the isotopic content of the organic matter. Samples of the nutrient salts added were analysed in a similar manner to determine the initial $\delta^{15}\text{N}$ of the medium. The $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ of the initial NO_3^- were also analysed as dissolved inorganic nitrogen (see below). Internal laboratory standards, calibrated to VPDB and atmospheric N_2 , were analysed every 10 samples, and data were corrected relative to the mean of the two nearest standards. External precision is approximately $\pm 0.2 \text{‰}$ for $\delta^{15}\text{N}$ and $\pm 0.1 \text{‰}$ for $\delta^{13}\text{C}$. The C/N ratio was calculated by comparing the integrated area of the major beams (mass 28 for N and mass 44 for C) to standards with known C/N ratios. The external precision for this method is $< 0.1 \text{‰}$.

Dissolved inorganic nitrogen

The $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ composition of the samples were determined using a GV IsoPrime with an external automated

purge-and-trap system at the University of Massachusetts, Dartmouth, SMAST campus. The NO_3^- was converted to N_2O using Cd reduction to NO_2^- followed by azide treatment (McIlvin and Altabet, 2005). Data are reported relative to atmospheric N_2 and VSMOW for nitrogen and oxygen, respectively. Each run of NO_3^- samples consisted of one operational blank (low-nutrient seawater treated with azide), three NO_2^- standards, a cadmium blank (low-nutrient seawater treated with cadmium) and three NO_3^- standards (USGS 34, 35 and an internal Altabet lab standard), followed by the prepared samples. Three randomly selected samples were also prepared in triplicate to check for method and machine reproducibility. The run ended with three more NO_2^- standards, three NO_3^- standards, a Cd blank and an operational blank. Analytical precision measured from multiple determinations on standards was approximately $\pm 0.2\text{‰}$ for $\delta^{15}\text{N}$ and $\pm 0.7\text{‰}$ for $\delta^{18}\text{O}$ (NO_3^- only).

Isotopic data produced from each run were scrutinized for standard precision throughout individual runs. Samples were corrected for the small amount ($\sim 15\%$) of oxygen exchange that occurs between the sample and water during the conversion to nitrous oxide, fractionation due to oxygen removal and the 1 : 1 addition of azide N to NO_2^- -N in the formation of N_2O (see McIlvin and Altabet, 2005, for an in-depth discussion of $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ corrections).

2 Nutrient concentrations

Concentrations of NO_3^- , NO_2^- and NH_4^+ in the growth solutions were analysed prior to, during and after each experiment. Nitrate and nitrite concentrations were determined by diazotization before and after reduction with cadmium (Grasshoff, 1976). Ammonium concentrations were determined with the indophenol-blue method. Note that the measured concentrations of the NO_3^- were slightly different than initial target concentrations.

3 Results

3.1 Nitrogen isotopes in algal material

3.1.1 Free-drift experiments

Results from the free-drift nutrient experiments from *Ulva* and *Agardhiella* are presented in Table 1. In each of the treatments the $\delta^{15}\text{N}$ of the new algal growth during each experiment and the residual NO_3^- concentrations left in each treatment after the 24 h incubations were determined. Although concentrations of NO_2^- and NH_4^+ were measured, none was detected. The $\delta^{15}\text{N}$ of the newly grown *Agardhiella* material decreased from 1.8‰ (14 μM) to 1.6‰ in the 50 μM treatment, to 0.7‰ in the 103 μM treatment and finally to -3.0‰ in the 485 μM experiment. Similar results were found in the experiments using *Ulva* although the $\delta^{15}\text{N}$ values were all

higher (Table 1). For example, in the lowest two NO_3^- treatments, the $\delta^{15}\text{N}$ of the *Ulva* was actually more positive than that of the NO_3^- in the growth medium (Table 1).

The C/N ratios and the $\delta^{13}\text{C}$ values of the algae are included in the supplementary information.

3.1.2 Syringe experiments

The results from all the syringe experiments are listed in Table 2. The $\delta^{15}\text{N}$ value of *Ulva* and *Agardhiella* exhibited small decreases.

3.2 Isotopic analysis of dissolved inorganic nitrogen

3.2.1 Free-drift experiments

The data from the free-drift experiments are presented in Table 1 with the trend in the $\delta^{15}\text{N}$ of the NO_3^- mirroring that of the solid algae. The mean $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values of the initial NO_3^- were $+3.3 \pm 0.3\text{‰}$ and $+23 \pm 0.3$, respectively ($n = 12$), and as the NO_3^- was consumed the residual NO_3^- became isotopically enriched in ^{15}N and ^{18}O . The $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values in both *Agardhiella* sp. ($r^2 = 0.60$, $n = 13$) and *Ulva* sp. ($r^2 = 0.79$, $n = 25$) experiments were positively correlated to each other, exhibiting a slope close to unity for both algae species (1.1 for *Ulva* sp. and 1.17 for *Agardhiella* sp.). In the lower-concentration experiments the slope increased to approximately 2.

4 Discussion

In order to calculate the fractionation during assimilation, the change in the $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ of the NO_3^- and the algal tissue was modelled using a Rayleigh distillation model. In the case of N, the $^{15}\text{N}/^{14}\text{N}$ of the new algal growth (RA) at time (t) is given by Eq. (3), while the $^{15}\text{N}/^{14}\text{N}$ of the residual NO_3^- (R) at t is given by Eq. (4).

$$\text{RA}t = \text{Ri} \frac{1 - f^{1/\alpha}}{1 - f} \quad (3)$$

$$\text{R}t = \text{Ri} f^{(\frac{1}{\alpha}-1)} \quad (4)$$

In these equations, f represents the fraction of the initial NO_3^- remaining; Ri the $^{15}\text{N}/^{14}\text{N}$ ratio of the initial NO_3^- ; Rt and RA the $^{15}\text{N}/^{14}\text{N}$ ratio of the NO_3^- and new algal growth, respectively, after a specific time during which f has been determined; and α the fractionation factor. The fractionation ($^{15}\epsilon$) can also be calculated using the approach of Mariotti et al. (1981) which utilizes a plot of the isotopic composition of the NO_3^- with respect to $\ln f$ or $\ln(\text{NO}_3^-(t)/\text{NO}_3^-(i))$ as in Eq. (5).

$$\delta t = \delta i - \epsilon \ln f \quad (5)$$

In Eq. (5), ϵ ($^{15}\epsilon$) is the slope of the relationship between $\delta^{15}\text{N}t$ and $\ln f$. The term δt is equal to the $\delta^{15}\text{N}$ of the NO_3^-

Table 3. Calculated fractionation (ϵ) for experiments using linear model.

#	Species	NO ₃ ⁻	NO ₃ ⁻	Nitrogen	Oxygen
		μM	ϵ (solid) ‰	ϵ (DIN) ‰	ϵ (DIN) ‰
1	<i>Ulva</i>	2.6	2.1		
2	<i>Ulva</i>	14	-3.2	0.8	1.5
2	<i>Ulva</i>	60	-0.2	1.5	3.6
2	<i>Ulva</i>	103	0.3	2.9	3.8
2	<i>Ulva</i>	485	2.0	3.5	5.6
1	<i>Agardhiella</i>	7	0	nm	nm
1	<i>Agardhiella</i>	10	0.4	nm	nm
2	<i>Agardhiella</i>	14	3.2	1.9	nm
2	<i>Agardhiella</i>	55	3.4	2.4	2.6
2	<i>Agardhiella</i>	104	3.0	5.1	4.8
2	<i>Agardhiella</i>	514	6.3	8.3	12.9

1 represents syringe experiment; 2 represents free drift; nm stands for not measured.

at time (t) when the concentration is equal to NO₃⁻ (t), and δi is equal to $\delta^{15}\text{N}$ of the NO₃⁻ at the initial time when the concentration is equal to NO₃⁻ (i). In the free-drift experiments where the $\delta^{15}\text{N}$ of the solution was sampled multiple times, the $\delta^{15}\text{N}$ values were measured at various concentrations as the NO₃⁻ was assimilated by the algae and hence values of $\ln f$ calculated. A similar approach was used to calculate $^{18}\epsilon$ using the $\delta^{18}\text{O}$ data.

An alternative method for calculating the fractionation factor used the measurement of the $\delta^{15}\text{N}$ of new algal tissue as a function of the expression in Eq. (6).

As f tends to 0 (all the NO₃⁻ was consumed), the $\delta^{15}\text{N}$ of the algae (δA) tended to approach the $\delta^{15}\text{N}$ of the initial NO₃⁻. Hence, utilizing Eq. (6), the slope of the relationship was equivalent to $(\alpha - 1) \times 1000$ or ϵ .

$$\delta A t = \delta i + \epsilon \frac{f \ln f}{1 - f} \quad (6)$$

In each of the experiments the slope of the line was determined by plotting the initial $\delta^{15}\text{N}$ of the NO₃⁻ at a f value of 0 and the measured $\delta^{15}\text{N}$ of the algae at the appropriate f value corresponding to the decrease in the concentration of NO₃⁻ at the end of 24 h.

While it is possible to arrive at an estimate of fractionation using either the solid sample or the NO₃⁻ data, the method of measuring the $\delta^{15}\text{N}$ of the residual NO₃⁻ may provide a more accurate method for a number of reasons. First, in the case of the measurement of the $\delta^{15}\text{N}$ of the tissue in the free-drift experiments, the f factor is calculated by averaging the amount of NO₃⁻ utilized during a 24 h period. This assumes that the algae grows equally throughout the 24 h period, rather than perhaps faster when the NO₃⁻ concentration is high and lower as the concentration is reduced. In addition, as the concentration of NO₃⁻ is reduced to low concentrations, the fractiona-

tion of ^{15}N will also change (see later discussion). It might be possible to model these changes, but the interpretation would be dependent upon a number of assumptions which could not be validated with the present data set. In this regard the $\delta^{15}\text{N}$ of the tissue grown in the syringe experiments might be more reliable in providing an estimate of fractionation as a constant amount of NO₃⁻ is supplied throughout the growth period, and therefore Rayleigh type modelling is unnecessary. In addition, both the syringe and the free-drift tissue measurements might suffer from the inability to precisely separate new and old algal tissue growth and the possibility of translocation of N-bearing compounds in the algal tissue. In contrast, the $\delta^{15}\text{N}$ of the residual NO₃⁻ provides a direct measurement of the fractionation during assimilation. While the results obtained between the two methods are similar, in cases where there are differences we feel that the data obtained from the $\delta^{15}\text{N}$ of the NO₃⁻ provide the best estimate of fractionation.

4.1 Modelling

As the NO₃⁻ removed from the medium was balanced by algal assimilation, isotopic fractionation produced corresponding changes in both the $\delta^{15}\text{N}$ of the residual NO₃⁻ and the $\delta^{15}\text{N}$ of new algal growth. These data are reported in Table 1, and the fractionation factors estimated for ^{15}N (and ^{18}O when applicable) using Eqs. (5) and (6) are reported in Table 3.

4.1.1 *Ulva*

The $^{15}\epsilon$ values calculated from the $\delta^{15}\text{N}$ of the algal growth and the NO₃⁻ show a decrease towards 0 with decreasing concentration of NO₃⁻ (Table 3, Fig. 3). At the higher concentrations, the estimate of $^{15}\epsilon$ obtained from the algal growth (~ 3 ‰) and that obtained from residual NO₃⁻ $\delta^{15}\text{N}$ are statistically the same, while at the lower initial NO₃⁻ concentrations values of $^{15}\epsilon$ obtained from the algal $\delta^{15}\text{N}$ are significantly lower (Fig. 4). If the observation that fractionation varies as a function of the concentration of NO₃⁻ is correct, then Eq. (3) can only yield a mean estimate of $^{15}\epsilon$ as during the experiment the concentration of NO₃⁻ changes considerably as it is assimilated. In fact the data from the NO₃⁻ free-drift experiment (Table 1) is best fitted by a quadratic equation confirming a change in fractionation with changing concentration (Fig. 5). Using a Chi-squared test, the improvement in the fit between the linear and non-linear model can be shown to be statistically significant at the 99 % level in both the 60 and 103 μM experiments. The first differential of the quadratic equation therefore provides an estimate of ϵ at any value of f . Using the data from the experiments which were initiated at concentrations of 14, 60 and 103 μM NO₃⁻ and calculating the mean $^{15}\epsilon$ value derived from each experiment with respect to concentration rather than f , a robust estimate of ϵ with respect to changing NO₃⁻ can be obtained




Figure 3. Calculation of $^{15}\epsilon$ using linear regression through $\delta^{15}\text{N}$ values of NO_3^- with respect to $\ln f$ for experiments in which *Ulva* sp. was incubated for 48 h. Samples of water were taken at 12, 24 and 48 h and measured for the concentration of remaining NO_3^- and its $\delta^{15}\text{N}$ (and $\delta^{18}\text{O}$). The numbers refer to the different initial concentrations of NO_3^- used in each experiment (See Table 1).

(Fig. 6). Data from the 500 μM experiment were not used in this estimate as a result of the small change in concentration of NO_3^- (and as a consequence a small change in f which occurred during the experiment at high concentration). Although the estimates of $^{15}\epsilon$ obtained from the non-linear equation predict a value of less than 0 at concentrations lower than $\sim 1 \mu\text{M}$, none of the experiments attained these low concentrations and therefore this observation will need to be confirmed. In addition the one syringe experiment performed with *Ulva* at a constant concentration of $\sim 3 \mu\text{M}$ yielded a $^{15}\epsilon$ value of 1 ‰, higher than the values estimated from the NO_3^- drawdown experiments. Hence such data were inconsistent with a $^{15}\epsilon$ value below 0. In the free-drift experiments however, the $\delta^{15}\text{N}$ of the measured algae was greater than that in the initial NO_3^- (5.1 and 4.0 ‰ in the 14 μM and 60 μM treatments, respectively, compared to the initial NO_3^- and algal values of 3.3 and 3.1 ‰, respectively) (Table 1), giving estimates of $^{15}\epsilon$ less than 0 ($\epsilon = -3.2$ ‰). While the data of the solids appear inconsistent with the data measured on the $\delta^{15}\text{N}$ of the NO_3^- , based on previous discussion our feeling is that assimilation factors calculated from the analysis of the $\delta^{15}\text{N}$ of the NO_3^- are correct and that the $\delta^{15}\text{N}$ of the solid material might therefore be some kind of artefact as discussed earlier. Regardless of whether $^{15}\epsilon$ is less than 0, both approaches show a decrease in $^{15}\epsilon$ with decreasing concentrations, and a rate of change of appears to be greatest at the lowest concentration, i.e. between 1 and 10 μM (Fig. 6).




Figure 4. Estimate of fractionation factor (α) and ϵ during the assimilation of NO_3^- by *Ulva* sp. based on the $\delta^{15}\text{N}$ analysis of the algal material (solid) and the DIN (data from Fig. 3).

With higher concentrations (between 10 and 50 to 500 μM) the fractionation appears to reach a constant positive value ($\epsilon = 3$ to 4 ‰).

4.1.2 *Agardhiella*

Based on both the algal and $\text{NO}_3^- \delta^{15}\text{N}$ data, this species also exhibited a strong dependence between fractionation and NO_3^- concentration. Values of $^{15}\epsilon$ were close to 0, or slightly negative, at low concentrations ($< 10 \mu\text{M}$) and increased between 100 and 500 μM , reaching a value of ~ 8 ‰ at 500 μM (Fig. 7). As a result of the fact that at most only three samples were taken for measurement of the $\delta^{15}\text{N}$ (and $\delta^{18}\text{O}$) of the NO_3^- during the free-drift experiments, it was not considered valuable to fit anything more than a straight line to the data, and therefore a more refined equation relating the change in ϵ to the concentration of NO_3^- was not calculated. As in the case of *Ulva*, there was a suggestion that $^{15}\epsilon$ values might fall below 0 at low concentrations, although the $\delta^{15}\text{N}$ of the solid material did not increase at low NO_3^- concentrations as seen in *Ulva* sp.

4.2 Concentration dependence of the fractionation factor

In microalgae and bacteria the uptake and fractionation of NO_3^- has been proposed to be a three-step process (Granger et al., 2004; Hoch et al., 1992; Karsh et al., 2012, 2014; Mariotti et al., 1982; Shearer et al., 1991): first a transport step across the cellular membrane (ϵ_{in}), then a nitrate reductase step (ϵ_{NR}) and finally a flux out of the cell (ϵ_{out}). The overall fractionation manifested by the organism, expressed as ϵ_{org} , is related to the influx, efflux and nitrate reductase




Figure 5. The changing fractionation as a consequence of decreasing NO_3^- concentration, as shown in Fig. 3, necessitates the use of a non-linear curve fitting to the data. The use of a quadratic equation shows an improved fit to the data and allows the slope of the relationship to be calculated at specific concentrations using the first differential of the equation. Data from the 485 μM experiment have been omitted as a result of the small change in the f value.

fractionation by Eq. (8), in which γ is the proportion of efflux relative to influx (Karsh et al., 2014).

$$\varepsilon_{\text{org}} = \varepsilon_{\text{in}} + \gamma (\varepsilon_{\text{NR}} + \varepsilon_{\text{out}}) \quad (7)$$

The estimated fractionation associated with these processes in a marine diatom (*Thalassiosira weissflogii*) are $^{15}\varepsilon_{\text{in}} = 2 \text{‰}$, $^{15}\varepsilon_{\text{out}} = 1.2 \text{‰}$ and $^{15}\varepsilon_{\text{NR}} = 26.6 \text{‰}$ (Karsh et al., 2012, 2014). As the majority of the fractionation is associated with the NR step, the degree to which this is expressed in the external medium and also in the organism is controlled by the amount of efflux relative to influx (γ). Accepting the possibility that there may be differences between microalgae and the organisms used in this study, we have nevertheless used this model as a basis with which to explain the observations of a concentration dependence on $^{15}\varepsilon_{\text{org}}$ made in this paper. In this regard it is helpful to examine the work of Needoba et al (2004), who measured the $\delta^{15}\text{N}$ of the internal and external NO_3^- pools. They determined that the maximum difference in $\delta^{15}\text{N}$ occurred in situations in which ε_{org} was at a minimum, thus indicating that the efflux from the cell was small. Conversely when fractionation was high, the difference between the $\delta^{15}\text{N}$ of the external and internal pools




Figure 6. Average fractionation factors (α) and ε values calculated using the mean values estimated from the first differential of the quadratic fits shown in Figure 5 for *Ulva* sp. Error bars represent $\pm\sigma$ of the estimate calculated using equations shown in Fig. 5.

was at a minimum and efflux maximal. As in both cases, the greatest potential for isotope fractionation is at the NR step (Karsh et al., 2012; Ledgard et al., 1985), the principal explanation for dependence on external concentration must relate to the ratio of NO_3^- uptake to efflux from the cell. At lower external concentrations, NO_3^- is limiting and the $\delta^{15}\text{N}$ of the internal pool is highly elevated. However, most of the NO_3^- is consumed and efflux is minimal, and, although the same amount of fractionation at the NR step takes place, this isotopic signal is not communicated to the external environment. At high concentrations the reverse is true: NO_3^- is not limiting, and the fractionation experienced at the NR step is translated to the external environment. Based on our findings we propose that macroalgae may behave similarly in many respects to microalgae. However, the only study we are aware of dealing with macroalgae concluded that the concentration of NO_3^- did not influence the fractionation of ^{15}N (Cohen and Fong, 2005) and would therefore appear to be in conflict with the results of this study. However, in the Cohen and Fong (2005) research the only experiments in which the concentration of NH_4^+ was not altered, in addition to NO_3^- , were carried out at relatively high concentrations of NO_3^- ($> 50 \mu\text{M}$). This is above the level at which the fractionation appeared to be constant in our study.




Figure 7. Estimate of fractionation (ϵ) exerted during the incorporation of NO_3^- into *Agardhiella* sp. based on the $\delta^{15}\text{N}$ analysis of the algal material and the DIN. Error bars represent $\pm\sigma$ of replicate measurements. The solid and the DIN data are based on the free-drift results.

4.3 Oxygen isotopic composition of NO_3^-

The measurement of the $\delta^{18}\text{O}$ of nitrate is a relatively new technique which has helped explain both the source of NO_3^- and the mechanism of fractionation of N and O isotopes during assimilation (Granger et al., 2004, 2010; Leichter et al., 2007; Wankel et al., 2006, 2009). The data presented here suggest that in a manner similar to ^{15}N the fractionation of ^{18}O is dependent upon the concentration of NO_3^- in the external environment (Tables 1 and 2; Fig. 8). While generally the fractionation of $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ are related in a 1 : 1 ratio (Granger et al., 2004), in this study the slope of the data seems to have a value of greater than the ideal 1 : 1 relationship (Fig. 2). It was argued by Granger et al. (2004) that this 1 : 1 relationship was consistent with fractionation of N and O during NR, whereas fractionation during uptake would give a 2 : 1 relationship. In more recent work it was shown that there are different degrees of fractionation for N compared to O during uptake and efflux, which would cause the relationship between $^{18}\epsilon$ and $^{15}\epsilon$ to rise significantly above unity, when fractionation is low (Karsh et al., 2014). Such data are in agreement with our study in that the $^{18}\epsilon : ^{15}\epsilon$ ratio is closest to unity in the highest-concentration ($\sim 500 \mu\text{M}$) experiments and increases with lower initial concentrations of NO_3^- , reaching a value of ~ 2 at $10 \mu\text{M}$. This 2 : 1 relationship corresponds to the lowest amount of fractionation observed ($\epsilon \sim 0\text{‰}$). Using the rationale suggested by Granger et al. (2004), this pattern is consistent with a change in fractionation from a process predominantly controlled by NR, to one in which fractionation is controlled by the relative difference between the fractionation of O and N during up-




Figure 8. Relationship between the change in concentration of NO_3^- and the $\delta^{18}\text{O}$ of the NO_3^- in the *Ulva* sp. free-drift experiments. Error bars represent $\pm\sigma$ of the analytical precision on the $\delta^{18}\text{O}$ measurements.

take (1.4) and efflux (2.3) (Karsh et al., 2014). If the results of these experiments are correct, then the relationship between $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ also should not be linear, but rather a quadratic, similar to that observed between the $\delta^{15}\text{N}$ and concentration discussed earlier. However, as a result of the larger error on the $\delta^{18}\text{O}$ compared to $\delta^{15}\text{N}$ (0.7 vs. 0.2‰), this pattern was not evident in the data collected in these experiments.

4.4 Biogeochemical implications

The observation of a concentration dependence upon ^{15}N fractionation during denitrification has been previously made for microbes (Kritee et al., 2012). Both the results of that study and the data presented here suggest that there is a relationship between fractionation and concentration during assimilation, which has implications for the application of nitrogen isotopes for detection of N sources. It is clear that under typical N-limiting conditions both micro- and macroalgae have the same isotopic composition as the ambient nitrate. However, when NO_3^- concentrations are elevated, algae fractionate the external NO_3^- pool, forming biomass which is relatively isotopically more negative than the ambient NO_3^- . The residual NO_3^- effluxed from the cell consequently becomes isotopically more positive regardless of the $\delta^{15}\text{N}$ of the original NO_3^- . Consider a hypothetical coastal estuary in which there is a significant input of NO_3^- from artificial fertilizers (with a $\delta^{15}\text{N} \sim 0\text{‰}$) applied to adjacent agricultural areas. As a result of the high NO_3^- concentrations, the fractionation during assimilation by algae would be

greater than 0, initially producing algal material with $\delta^{15}\text{N}$ values more negative than the original NO_3^- . As the NO_3^- is consumed, the $\delta^{15}\text{N}$ of the residual NO_3^- would become more positive. Eventually isotopically positive algal material would be formed from waters which originally had a $\delta^{15}\text{N}$ close to 0‰. It might be incorrectly assumed in such instances that the algal material was affected by nitrogen derived from an isotopically positive source, when in fact the positive values were produced as a result of fractionation during assimilation.

5 Conclusions

1. There is a concentration dependence upon the fractionation of ^{15}N and ^{18}O exerted during macroalgal assimilation of NO_3^- . This dependence varies according to species but approaches 0 at low concentrations in both of the algal species studied here. This concentration dependence essentially means that in most open marine environments, which have NO_3^- concentrations of less than $2\ \mu\text{M}$, there is minimal fractionation during assimilation. In environments with higher concentrations of NO_3^- , the fractionation is greater than 0, leading to enrichment in the ^{15}N of the residual NO_3^- regardless of the $\delta^{15}\text{N}$ of the original source of that NO_3^- . The observation of the concentration dependence of $^{15}\epsilon$ helps to explain the wide range of values reported in the literature where experiments were carried over a wide range of NO_3^- concentrations.
2. The change in the $^{15}\epsilon$ shows the largest rate of variation at low NO_3^- concentrations, and there is a suggestion $^{15}\epsilon$ may fall below 0. This might imply that organic material formed under very low NO_3^- concentrations could manifest an inverse isotopic effect.
3. The $^{18}\epsilon$ also shows a dependence on concentration and is related to $^{15}\epsilon$ in a 1 : 1 manner at higher concentrations ($> 100\ \mu\text{M}$) of NO_3^- . At lower concentrations the slope of $^{18}\epsilon : ^{15}\epsilon$ approaches values of 2 : 1.
4. The change of the fractionation of both ^{15}N and ^{18}O with respect to the concentration of NO_3^- supports a model in which there is a change in the origin of the fractionation from one in which the control is exerted by the NR step to one in which the control is exerted by difference between fractionation exerted in uptake and efflux.

The Supplement related to this article is available online at doi:10.5194/bg-11-6147-2014-supplement.

Acknowledgements. We would like to thank the staff of the *Aplysia* facility and the Stable Isotope Laboratory at the University of Miami. Funding for this project was provided by EPA grants to P. K. Swart. We would like to thank Quinn Devlin for assistance in the laboratory and helpful discussions. Additional funding for this project was provided by the Stable Isotope Laboratory at the University of Miami.

Edited by: B. A. Bergamaschi

References

- Altabet, M. A.: A time-series study of the vertical structure of nitrogen and particle dynamics in the Sargasso Sea, *Limnol. Ocean.*, 34, 1185–1201, 1989.
- Altabet, M. A., Deuser, W. G., Honjo, S., and Stienen, C.: Seasonal and depth-related changes in the source of sinking particles in the North-Atlantic, *Nature*, 354, 136–139, 1991.
- Barford, C. C., Montoya, J. P., Altabet, M. A., and Mitchell, R.: Steady-state nitrogen isotope effects of N_2 and N_2O production in *Paracoccus denitrificans*, *Appl. Environ. Microbiol.*, 65, 989–994, 1999.
- Carballeira, C., Viana, I. G., and Carballeira, A.: $\delta^{15}\text{N}$ values of macroalgae as an indicator of the potential presence of waste disposal from land-based marine fish farms, *J. Appl. Phycol.*, 25, 97–107, 2013.
- Cohen, R. A. and Fong, P.: Experimental evidence supports the use of $\delta^{15}\text{N}$ of the opportunistic green macroalga *Enteromorpha intestinalis* (Chlorophyta) to determine nitrogen sources to estuaries, *J. Phy.*, 41, 287–293, 2005.
- Costanzo, S. D., O' Donohue, M. J., Dennison, W. C., Loneragan, N. R., and Thomas, M.: A new approach for detecting and mapping sewage impacts, *Mar. Pollut. Bull.*, 42, 149–156, 2001.
- Delwiche, C. C. and Steyn, P. L.: Nitrogen isotope fractionation in soils and microbial reactions, *Env. Sci. Tech.*, 4, 929–935, 1970.
- Deutsch, B. and Voss, M.: Anthropogenic nitrogen input traced by means of $\delta^{15}\text{N}$ values in macroalgae: Results from in-situ incubation experiments, *Sci. Total Environ.*, 366, 799–808, 2006.
- Dugdale, R. C. and Wilkerson, F. P.: The use of ^{15}N to measure nitrogen uptake in eutrophic oceans; experimental considerations, *Limnol. Ocean.*, 31, 673–689, 1986.
- Granger, J., D. M. Sigman, J. A. Needoba, and Harrison, P. J.: Coupled nitrogen and oxygen isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton, *Limnol. Oceanogr.*, 49, 1763–1773, 2004.
- Granger, J., Sigman, D. M., Prokopenko, M. G., Lehmann, M. F., and Tortell, P. D.: A method for nitrite removal in nitrate N and O isotope analyses, *Limnol. Oceanogr.-Meth.*, 4, 205–212, 2006.
- Granger, J., Sigman, D. M., Rohde, M. M., Maldonado, M. T., and Tortell, P. D.: N and O isotope effects during nitrate assimilation by unicellular prokaryotic and eukaryotic plankton cultures, *Geochim. Cosmochim. Acta*, 74, 1030–1040, 2010.
- Grasshoff, K.: *Methods of Seawater Analysis*, Verlag Chemie, Weinheim, 1976.
- Guillard, R. R. L.: Culture of phytoplankton for feeding marine invertebrates, in: *Culture of Marine Invertebrate Animals*, edited by: Smith, W. L. and Chanley, M. H., Plenum Press, New York, USA, 29–60, 1975.

- Haug, G. H., Pedersen, T. F., Sigman, D. M., Calvert, S. E., Nielsen, B., and Peterson, L. C.: Glacial/interglacial variations in production and nitrogen fixation in the Cariaco Basin during the last 580 kyrs, *Paleocean*, 13, 427–432, 1998.
- Heaton, T. H.: Isotopic studies of nitrogen pollution in the hydrosphere and atmosphere: a review, *Chem. Geol.*, 59, 87–102, 1986.
- Hoch, M. P., Fogel, M. L., and Kirchman, D. L.: Isotope fractionation associated with ammonium uptake by a marine bacterium, *Limnol. Ocean.*, 37, 1447–1459, 1992.
- Hoering, T. C. and Ford, H.: The isotope effect in the fixation of nitrogen by *Azotobacter*, *Am. J. Chem. Soc.*, 82, 376–378, 1960.
- Horrigan, S. G., Montoya, J. P., Nevins, J. L., and McCarthy, J. J.: Natural isotopic composition of dissolved inorganic nitrogen in the Chesapeake Bay, *Est. Coast. Shelf Sci.*, 30, 393–410, 1990.
- Karsh, K. L., Granger, J., Kritee, K., and Sigman, D. M.: Eukaryotic assimilatory nitrate reductase fractionates N and O isotopes with a ratio near unity, *Env. Sci. Tech.*, 46, 5727–5735, 2012.
- Karsh, K. L., Trull, T. W., Sigman, D. M., Thompson, P. A., and Granger, J.: The contributions of nitrate uptake and efflux to isotope fractionation during algal nitrate assimilation, *Geochim. Cosmochim. Acta*, 132, 391–412, doi:10.1016/j.gca.2013.09.030, 2014.
- Kritee, K., Sigman, D. M., Granger, J., Ward, B. B., Jayakumar, A., and Deutsch, C.: Reduced isotope fractionation by denitrification under conditions relevant to the ocean, *Geochim. Cosmochim. Acta*, 92, 243–259, 2012.
- Lajtha, K. and Michener, R. H.: *Stable Isotopes in Ecology and Environmental Science*, Blackwell, London, 1994.
- Lamb, K., Swart, P. K., and Altabet, M. A.: Nitrogen isotopic systematics in the Florida reef tract, *Bull. Mar. Sci.*, 88, 119–146, 2012.
- Lapointe, B. E., Barile, P. J., and Matzie, W. R.: Anthropogenic nutrient enrichment of seagrass and coral reef communities in the Lower Florida Keys: discrimination of local versus regional nitrogen sources, *J. Exp. Mar. Biol. Ecol.*, 308, 23–58, 2004.
- Ledgard, S. F., Woo, K. C., and Bergersen, F. J.: Isotopic fractionation during reduction of nitrate and nitrite by extracts of spinach leaves, *Austral. J. Plant Phys.*, 12, 631–640, 1985.
- Leichter, J. J., Paytan, A., Wankel, S., and Hanson, K.: Nitrogen and oxygen isotopic signatures of subsurface nitrate seaward of the Florida Keys reef tract, *Limnol. Ocean.*, 52, 1258–1267, 2007.
- Mariotti, A., Germon, J. C., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A., and Tardieux, P.: Experimental-determination of nitrogen kinetic isotope fractionation: Some principles-Illustration for the denitrification and nitrification processes, *Plant Soil*, 62, 413–430, 1981.
- Mariotti, A., Mariotti, F., Champigny, M. L., Amarger, N., and Moyse, A.: Nitrogen isotope fractionation associated with nitrate reductase activity and uptake of NO_3^- by Pearl Millet, *Plant Physiol.*, 69, 880–884, 1982.
- McIlvin, M. R. and Altabet, M. A.: Chemical conversion of nitrate and nitrite to nitrous oxide for nitrogen and oxygen isotopic analysis in freshwater and seawater, *Anal. Chem.*, 77, 5589–5595, 2005.
- Miyake, Y. and Wada, E.: The isotope effect on the nitrogen in biochemical, oxidation-reduction reactions, *Records of Oceanographic Works in Japan*, 11, 1–6, 1971.
- Montoya, J. P., Horrigan, S. G., and McCarthy, J. J.: Natural abundance of ^{15}N in particulate nitrogen and zooplankton in the Chesapeake Bay, *Mar. Ecol.-Prog. Ser.*, 65, 35–61, 1990.
- Needoba, J. A., Waser, N. A., Harrison, P. J., and Calvert, S. E.: Nitrogen isotope fractionation in 12 species of marine phytoplankton during growth on nitrate, *Mar. Ecol.-Prog. Ser.*, 255, 81–91, 2003.
- Needoba, J. A., Sigman, D. M., and Harrison, P. J.: The mechanism of isotope fractionation during algal nitrate assimilation as illuminated by the $^{15}\text{N}/^{14}\text{N}$ of intracellular nitrate, *J. Phy.*, 40, 517–522, 2004.
- Pennock, J. R., Velinsky, D. J., Ludlam, J. M., Sharp, J. H., and Fogel, M. L.: Isotopic fractionation of ammonium and nitrate during uptake by *Skeletonema costatum*: Implications for $\delta^{15}\text{N}$ dynamics under bloom conditions, *Limnol. Ocean.*, 41, 451–459, 1996.
- Shearer, G., Schneider, J. D., and Kohl, D. H.: Separating efflux and influx components of net nitrate uptake by *Synechococcus*-R2 under steady state conditions, *J. Gen. Microbiol.*, 137, 1179–1184, 1991.
- Sigman, D. M., Lehman, S. J., and Oppo, D. W.: Evaluating mechanisms of nutrient depletion and C-13 enrichment in the intermediate-depth Atlantic during the last ice age, *Paleocean.*, 18, 1072, doi:10.1029/2002PA000818, 2003.
- Stokes, M. D., Leichter, J. J., Wing, S., and Frew, R.: Temperature variability and algal isotopic heterogeneity on a Floridian coral reef, *Mar. Ecol.*, 32, 364–379, 2011.
- Teichberg, M., Heffner, L. R., Fox, S., and Valiela, I.: Nitrate reductase and glutamine synthetase activity, internal N pools, and growth of *Ulva lactuca*: responses to long and short-term N supply, *Mar. Biol.*, 151, 1249–1259, 2007.
- Viana, I. G. and Bode, A.: Stable nitrogen isotopes in coastal macroalgae: Geographic and anthropogenic variability, *Sci. Total Environ.*, 443, 887–895, 2013.
- Wada, E. and Hattori, A.: Nitrogen isotope effects in the assimilation of inorganic nitrogenous compounds by marine diatoms, *Geomicro. J.*, 1, 85–101, 1978.
- Wankel, S. D., Kendall, C., Francis, C. A., and Paytan, A.: Nitrogen sources and cycling in the San Francisco Bay Estuary: a nitrate dual isotopic composition approach, *Limnol. Ocean.*, 51, 1654–1664, 2006.
- Wankel, S. D., Kendall, C., and Paytan, A.: Using nitrate dual isotopic composition ($\delta^{15}\text{N}$ and $\delta^{18}\text{O}$) as a tool for exploring sources and cycling of nitrate in an estuarine system: Elkhorn Slough, California, *J. Geophys. Res. Biogeosci.*, 114, G01011, doi:10.1029/2008JG000729, 2009.
- Waser, N. A., Yin, K. D., Yu, Z. M., Tada, K., Harrison, P. J., Turpin, D. H., and Calvert, S. E.: Nitrogen isotope fractionation during nitrate, ammonium and urea uptake by marine diatoms and coccolithophores under various conditions of N availability, *Mar. Ecol.-Prog. Ser.*, 169, 29–41, 1998.