



# Radiocarbon isotopic evidence for assimilation of atmospheric CO<sub>2</sub> by the seagrass *Zostera marina*

K. Watanabe and T. Kuwae

Coastal and Estuarine Environment Research Group, Port and Airport Research Institute, 3-1-1 Nagase, Yokosuka 239-0826, Japan

Correspondence to: K. Watanabe (watanabe-ke@ipc.pari.go.jp)

Received: 2 April 2015 – Published in Biogeosciences Discuss.: 21 May 2015

Revised: 16 October 2015 – Accepted: 20 October 2015 – Published: 30 October 2015

**Abstract.** Submerged aquatic vegetation takes up water-column dissolved inorganic carbon (DIC) as a carbon source across its thin cuticle layer. It is expected that marine macrophytes also use atmospheric CO<sub>2</sub> when exposed to air during low tide, although assimilation of atmospheric CO<sub>2</sub> has never been quantitatively evaluated. Using the radiocarbon isotopic signatures ( $\Delta^{14}\text{C}$ ) of the seagrass *Zostera marina*, DIC and particulate organic carbon (POC), we show quantitatively that *Z. marina* takes up and assimilates atmospheric modern CO<sub>2</sub> in a shallow coastal ecosystem. The  $\Delta^{14}\text{C}$  values of the seagrass (−40 to −10‰) were significantly higher than those of aquatic DIC (−46 to −18‰), indicating that the seagrass uses a <sup>14</sup>C-rich carbon source (atmospheric CO<sub>2</sub>, +17‰). A carbon-source mixing model indicated that the seagrass assimilated 0–40% (mean, 17%) of its inorganic carbon as atmospheric CO<sub>2</sub>. CO<sub>2</sub> exchange between the air and the seagrass might be enhanced by the presence of a very thin film of water over the air-exposed leaves during low tide. Our radiocarbon isotope analysis, showing assimilation of atmospheric modern CO<sub>2</sub> as an inorganic carbon source, improves our understanding of the role of seagrass meadows in coastal carbon dynamics.

## 1 Introduction

Submerged aquatic vegetation assimilates dissolved inorganic carbon (DIC) from the water column as a carbon source. Seagrasses take up DIC across their thin cuticle layer (Hemminga and Duarte, 2000), as their leaves lack stomata despite being angiosperms (Larkum and Den Hartog, 1989). An alternative carbon source, atmospheric CO<sub>2</sub> (C<sub>air</sub>), can-

not directly reach seagrasses when they are completely submerged; however, seagrasses can take up C<sub>air</sub> when their leaves are exposed to air during low tide (Clavier et al., 2011; Jiang et al., 2014; Leuschner and Rees, 1993). Seagrasses rely largely on aqueous CO<sub>2</sub> (CO<sub>2</sub>(aq)) as a carbon source for photosynthesis in nature (Beer and Koch, 1996). Some seagrass species, however, can use bicarbonate ions (HCO<sub>3</sub><sup>−</sup>) as a major carbon source (Beer and Rehnberg, 1997; Beer et al., 2002), although there is considerable interspecific variation in HCO<sub>3</sub><sup>−</sup> utilization (Campbell and Fourqurean, 2013). As CO<sub>2</sub>(aq) is in limited supply under normal seawater conditions (pH ≈ 8), comprising only 1% (roughly 10–15 μmol L<sup>−1</sup>) of the DIC pool, photosynthesis in seagrasses under high light conditions is frequently limited by carbon availability (Campbell and Fourqurean, 2013; Invers et al., 2001; Zimmerman et al., 1995). Under normal seawater pH conditions, the bicarbonate ion (HCO<sub>3</sub><sup>−</sup>) is the most abundant inorganic carbon species, accounting for nearly 90% of the DIC pool (Plummer and Busenberg, 1982; Zeebe and Wolf-Gladrow, 2001). Some seagrass species indirectly use HCO<sub>3</sub><sup>−</sup> under low-CO<sub>2</sub>(aq) conditions (Beer et al., 2002; Campbell and Fourqurean, 2013), using one or both of the following suggested mechanisms: (1) extracellular dehydration of HCO<sub>3</sub><sup>−</sup> into CO<sub>2</sub>(aq) via membrane-bound enzymes (Beer and Rehnberg, 1997) and/or (2) electrogenic proton (H<sup>+</sup>) extrusion into an boundary layer on the leaf surface, facilitating HCO<sub>3</sub><sup>−</sup>/H<sup>+</sup> cotransport (Hellblom et al., 2001).

Diffusion of CO<sub>2</sub> in water is much slower than that in air. During low tide, air-exposed aquatic macrophytes have a thin film of water between the air and their leaves, which promotes the uptake of C<sub>air</sub>, in contrast to high tide, when there is a thick water layer inhibiting the uptake of C<sub>air</sub> (Ji


and Tanaka, 2002). Previous studies have shown the possibility of C<sub>air</sub> uptake by seagrasses by using evidence from stable carbon isotope ratios ( $\delta^{13}\text{C}$ ) in seagrasses and the two carbon sources (DIC and C<sub>air</sub>; Clavier et al., 2011; Cooper and McRoy, 1988; Raven et al., 2002). However, the  $^{13}\text{C}$  method has considerable uncertainty because in addition to the source of carbon, the  $\delta^{13}\text{C}$  values of seagrasses are also determined by other factors such as the chemical species of DIC (CO<sub>2</sub>(aq) or HCO<sub>3</sub><sup>-</sup>) and photosynthetic carbon demand. The chemical species in the carbonate system (CO<sub>2</sub>(aq), HCO<sub>3</sub><sup>-</sup>, and carbonate ion; CO<sub>3</sub><sup>2-</sup>) have distinct  $\delta^{13}\text{C}$  values, and isotopic fractionations change depending on pH and temperature (Zeebe and Wolf-Gladrow, 2001; Zhang et al., 1995). Because the  $\delta^{13}\text{C}$  of HCO<sub>3</sub><sup>-</sup> (0‰) is isotopically distinct from that of both CO<sub>2</sub>(aq) (-9‰) and C<sub>air</sub> (-8‰) under normal seawater conditions (pH ≈ 8),  $\delta^{13}\text{C}$  values in seagrasses become higher with increasing of HCO<sub>3</sub><sup>-</sup> use (Campbell and Fourqurean, 2009; Hemminga and Mateo, 1996; Raven et al., 2002). However, quantification of the contribution of C<sub>air</sub> is impossible because of the  $\delta^{13}\text{C}$  value overlap between CO<sub>2</sub>(aq) and C<sub>air</sub> although low  $\delta^{13}\text{C}$  in seagrasses could be explained by the assimilation of either  $^{13}\text{C}$ -depleted CO<sub>2</sub>(aq) or C<sub>air</sub>. Also, changes in the photosynthetic carbon demand driven by irradiance fluctuations affect the isotopic fractionation factor (Hemminga and Mateo, 1996; Raven et al., 2002).

The natural abundance of radiocarbon ( $^{14}\text{C}$ ) has recently been used to assess food web structures (Ishikawa et al., 2014) and the origin and components of organic-matter pools (Goñi et al., 2013), as carbon sources have specific  $^{14}\text{C}$  concentrations ( $\Delta^{14}\text{C}$ ). The  $\Delta^{14}\text{C}$  of inorganic carbon also has specific values depending on the source, such as DIC or C<sub>air</sub>. The  $\Delta^{14}\text{C}$  of DIC generally differs from that of atmospheric CO<sub>2</sub> because of the longer residence time of carbon in aquatic ecosystems than in the atmosphere (Ishikawa et al., 2014; Stuiver and Braziunas, 1993). Moreover, the calculation of  $\Delta^{14}\text{C}$  by internal correction using  $\delta^{13}\text{C}$  values eliminates any effects from isotopic fractionation (Stuiver and Polach, 1977), overcoming one of the major uncertainties in the conventional  $\delta^{13}\text{C}$  approach. This study is the first to show quantitative evidence that the seagrass *Zostera marina* assimilates modern C<sub>air</sub>, based on the  $\Delta^{14}\text{C}$  values of the seagrass and two carbon sources.

## 2 Material and methods

### 2.1 Field surveys

Field surveys were conducted in 2014 during the growing season of *Z. marina* (May, July, September and November) in Furen Lagoon, Japan (Fig. 1; 43°19'46.5" N, 145°15'27.8" E). The lagoon is covered by ice from December to April. Furen Lagoon is brackish (salinity, ~30), and the northern part of the lagoon receives freshwater from the



**Figure 1.** Location of Furen Lagoon and sampling stations. The area offshore of Furen Lagoon is affected by both the Oyashio and the Soya warm current. The northern part of the lagoon receives freshwater from the Furen, Yausubetsu, and Pon-Yausubetsu rivers. Closed circles show lagoon stations. Water samples for DIC were collected at stations F1–F7. Seagrass samples were collected at stations F3, F4, F8 and F9 (marked with \*). POC samples were collected at stations R1–R3. The green-shaded areas indicate seagrass meadows.

Furen, Yausubetsu, and Pon-Yausubetsu rivers. The lagoon is covered by large seagrass meadows (67 % of the total area) dominated by *Z. marina*. The offshore of the lagoon (Sea of Okhotsk) is influenced by the dynamics of both the Oyashio and the Soya warm current. Surface water samples (depth, 0.1 m) for DIC (concentration and isotopic signatures) and total alkalinity (TA) in the water column were collected from a research vessel along the salinity gradient at seven stations in the lagoon (Fig. 1; stations F1–F7). At each station, one water sample was collected for measuring DIC and TA, and the salinity of the surface water was recorded with a conductivity–temperature sensor (COMPACT-CT; JFE Advantech, Nishinomiya, Japan). The samples for isotopic analysis of DIC were collected into 500 mL hermetically sealed glass bottles (Duran bottle; SCHOTT AG, Mainz, Germany), which were poisoned by adding saturated mercuric chloride solution (400  $\mu\text{L}$  per bottle) to prevent changes in DIC due to biological activity. The samples for measuring DIC concentration and TA were collected into 250 mL Duran bottles (SCHOTT AG), which were poisoned with saturated mercuric chloride solution (200  $\mu\text{L}$  per bottle). Seagrass (*Z. marina*) leaves were collected at four stations covered by *Z. marina* meadows (Fig. 1; stations F3, F4, F8 and F9) along the salinity gradient. The stations were located in sub-tidal zones (mean water depth, 0.83–1.12 m). The above-

ground wet-weight biomass of the seagrass, estimated from randomly thrown quadrats (0.0625 m<sup>2</sup>), ranged from 500 to 6800 g m<sup>-2</sup>. Three or four independent samples of seagrass leaves were collected at each station. Both the biofilm and epiphytes covering the leaves were gently removed by hands with powder-free gloves and washed off using ultrapure water (Milli-Q water; Millipore, Billerica, MA, USA). To estimate the  $\Delta^{14}\text{C}$  of C<sub>air</sub>, leaves of a terrestrial plant (giant reed, *Phragmites australis*) were collected near the lagoon. Plant samples were freeze-dried and subsamples were homogenized. To remove carbonate, the plant samples were acidified with 1 N HCl and dried again.

Water samples for the isotopic analysis of terrestrial particulate organic carbon (POC) were collected at three riverine stations (Fig. 1; stations R1–R3). Samples for POC were obtained by filtration (approximately 1 L) onto pre-combusted (450 °C for 2 h) glass-fiber filters (GF/F, Whatman, Maidstone, Kent, UK).

## 2.2 Carbon isotope analysis

We determined the stable carbon isotope ratios ( $\delta^{13}\text{C}$ ) and radiocarbon concentrations ( $\Delta^{14}\text{C}$ ) of seagrass leaves, terrestrial plant leaves, DIC samples and POC samples. Prior to  $\Delta^{14}\text{C}$  and  $\delta^{13}\text{C}$  measurements, samples were subjected to graphite purification as follows. DIC samples for  $\Delta^{14}\text{C}$  and  $\delta^{13}\text{C}$  analysis were acidified (pH < 2) with H<sub>3</sub>PO<sub>4</sub> and sparged using ultra-high purity mixed N<sub>2</sub>/H<sub>2</sub> gas. The powdered plant leaves and POC samples for  $\Delta^{14}\text{C}$  and  $\delta^{13}\text{C}$  analysis were combusted in an elemental analyzer (either a Euro EA3000, EuroVector, Milan, Italy; or a Flash 2000, Thermo Fisher Scientific, Inc., Waltham, Massachusetts, USA). For each process, the CO<sub>2</sub> evolved was collected cryogenically and purified in a vacuum line. The purified CO<sub>2</sub> was then reduced to graphite using hydrogen and an iron catalyst at 650 °C for 10 h. The <sup>13</sup>C and <sup>14</sup>C concentrations were measured using an accelerator mass spectrometer (AMS). The AMS results are reported as  $\Delta^{14}\text{C}$  (‰) values (Stuiver and Polach, 1977) as follows:

$$\Delta^{14}\text{C}(\text{‰}) = \delta^{14}\text{C} - 2(\delta^{13}\text{C} + 25)(1 + \delta^{14}\text{C}/1000). \quad (1)$$

The  $\Delta^{14}\text{C}$  values have been corrected by the radioactive decay of an international standard (oxalic acid) since AD 1950 (Stuiver and Polach, 1977). The  $\delta^{13}\text{C}$  values are reported relative to Vienna Pee Dee Belemnite.  $\delta^{13}\text{C}$  data were corrected using an internal standard. The analytical precision of the AMS was within 0.7 ‰ for  $\delta^{13}\text{C}$  and 3 ‰ for  $\Delta^{14}\text{C}$ .

## 2.3 Carbonate system analysis

DIC concentration and TA were determined on a batch-sample analyzer (ATT-05; Kimoto Electric, Osaka, Japan). The precision of the analyses was 4 μmol L<sup>-1</sup> for DIC and 3 μmol L<sup>-1</sup> for TA. The concentrations of CO<sub>2</sub>(aq), HCO<sub>3</sub><sup>-</sup>, and CO<sub>3</sub><sup>2-</sup> were estimated using chemical equilibrium rela-

tionships and the TA and DIC concentrations of the water samples (Zeebe and Wolf-Gladrow, 2001). The  $\delta^{13}\text{C}$  values of CO<sub>2</sub>(aq) ( $\delta^{13}\text{C}_{\text{CO}_2(\text{aq})}$ ) and HCO<sub>3</sub><sup>-</sup> ( $\delta^{13}\text{C}_{\text{HCO}_3^-}$ ) were calculated as follows (Zeebe and Wolf-Gladrow, 2001; Zhang et al., 1995):

$$\delta^{13}\text{C}_{\text{HCO}_3^-} = \delta^{13}\text{C}_{\text{DIC}} - (\varepsilon_{\text{db}} \cdot [\text{CO}_2(\text{aq})] + \varepsilon_{\text{cb}} \cdot [\text{CO}_3^{2-}]) / [\text{DIC}], \quad (2)$$

$$\delta^{13}\text{C}_{\text{CO}_2(\text{aq})} = \delta^{13}\text{C}_{\text{HCO}_3^-} + \varepsilon_{\text{db}}, \quad (3)$$

$$\varepsilon_{\text{db}} = \varepsilon(\text{CO}_2(\text{aq}) - \text{HCO}_3^-) = -9866/T + 24.12(\text{‰}), \quad (4)$$

$$\varepsilon_{\text{cb}} = \varepsilon(\text{CO}_3^{2-} - \text{HCO}_3^-) = -867/T + 2.52(\text{‰}), \quad (5)$$

where [CO<sub>2</sub>(aq)], [CO<sub>3</sub><sup>2-</sup>], and [DIC] are the concentrations of CO<sub>2</sub>(aq), CO<sub>3</sub><sup>2-</sup> and DIC, respectively; *T* is water temperature (K); and  $\varepsilon_{\text{db}}$  and  $\varepsilon_{\text{cb}}$  are factors for the isotopic fractionation between CO<sub>2</sub>(aq) and HCO<sub>3</sub><sup>-</sup>, and between CO<sub>3</sub><sup>2-</sup> and HCO<sub>3</sub><sup>-</sup>, respectively.

## 2.4 Data analysis

Because DIC taken up by seagrasses is a mixture of DIC from two sources (terrestrial and oceanic) each having distinct  $\Delta^{14}\text{C}$  values, it is reasonable to use salinity as a proxy for the extent of mixing of these two sources as well as for the salinity gradient-based comparison between  $\Delta^{14}\text{C}$  of DIC and seagrass. This comparison was therefore possible even though DIC and *Z. marina* samples were not necessarily collected from the same stations (Fig. 1). Analyses of covariance (ANCOVA) were used to examine the difference in  $\Delta^{14}\text{C}$  value between seagrass leaves and DIC. These differences provide evidence that the seagrasses assimilate C<sub>air</sub>. We selected salinity, categorical data (seagrass leaves or DIC) and the interaction term as the explanatory variables.


The relative contribution of C<sub>air</sub> to assimilated seagrass carbon was calculated by a two-carbon-source mixing model using the  $\Delta^{14}\text{C}$  values of DIC ( $\Delta^{14}\text{C}_{\text{DIC}}$ ), C<sub>air</sub> ( $\Delta^{14}\text{C}_{\text{air}}$ ), and the seagrass ( $\Delta^{14}\text{C}_{\text{seagrass}}$ ) at each of four stations as follows:

$$\text{C}_{\text{air}}(\text{\% contribution}) = (\Delta^{14}\text{C}_{\text{seagrass}} - \Delta^{14}\text{C}_{\text{DIC}}) / (\Delta^{14}\text{C}_{\text{air}} - \Delta^{14}\text{C}_{\text{DIC}}) \cdot 100. \quad (6)$$


$\Delta^{14}\text{C}_{\text{air}}$  was estimated from the  $\Delta^{14}\text{C}$  value of the sampled terrestrial plants ( $\Delta^{14}\text{C} = +17.2 \text{‰}$ ). The  $\Delta^{14}\text{C}$  values of DIC as the carbon source for *Z. marina* in the mixing model were estimated from the linear model fitted with the ANCOVA.

## 3 Results and discussion

Our radiocarbon isotopic analysis shows quantitatively that the seagrass *Z. marina* uses C<sub>air</sub> in a shallow lagoon (Fig. 2a). In May and July 2014,  $\Delta^{14}\text{C}_{\text{seagrass}}$  was significantly higher



**Figure 2.** (a) Spatial distribution of the  $\Delta^{14}\text{C}$  values of dissolved inorganic carbon (DIC) (blue open circles) and seagrass (red open circles) along the salinity gradient in May and July 2014 in Furen Lagoon, Japan. Blue and red solid lines represent the linear models fitted with analyses of covariance (ANCOVA) examined for DIC and seagrass, respectively. (b) Spatial distribution of the relative contribution of  $\text{C}_{\text{air}}$  to total inorganic carbon assimilated by seagrass along the salinity gradient, as calculated by the two-carbon-source mixing model. (c) Spatial distribution of the  $\delta^{13}\text{C}$  values of DIC (blue open circles) and seagrass (red open circles) along the salinity gradient. (d) Spatial distribution of the  $\delta^{13}\text{C}$  values of bicarbonate ion ( $\text{HCO}_3^-$ ; green open circles) and aqueous  $\text{CO}_2$  [ $\text{CO}_2(\text{aq})$ ] (purple open circles) along the salinity gradient. The dashed line indicates the isotopic signature of atmospheric  $\text{CO}_2$  ( $\text{C}_{\text{air}}$ ).




**Figure 3.** Spatial distribution of the  $\Delta^{14}\text{C}$  values of dissolved inorganic carbon (DIC; blue) and seagrass (red) along the salinity gradient in September (open circles) and November (open triangles) 2014 in Furen Lagoon, Japan. The dashed line indicates the  $\Delta^{14}\text{C}$  of atmospheric  $\text{CO}_2$  ( $\Delta^{14}\text{C}_{\text{air}}$ ).

than  $\Delta^{14}\text{C}_{\text{DIC}}$  even if the effects of salinity were considered (ANCOVA,  $P < 0.001$ ), and the interaction term was not significant (ANCOVA,  $P > 0.05$ ). Our results indicate that the changes in  $\Delta^{14}\text{C}_{\text{DIC}}$  are regulated mostly by mixing between high- $\Delta^{14}\text{C}$  river water and low- $\Delta^{14}\text{C}$  seawater: the seagrass uses aquatic DIC as the main carbon source, as expected from previous studies (Campbell and Fourqurean, 2013; Hemminga and Duarte, 2000; Invers et al., 2001). The  $\Delta^{14}\text{C}_{\text{seagrass}}$  reflects  $\Delta^{14}\text{C}_{\text{DIC}}$  from May to July because *Z. marina* leaves start to grow in early May when sea ice is thawing at the study site, with the turnover time of leaves being 30–90 days (mean, 60 days; Hosokawa et al., 2009). Furthermore, the negative relationship between salinity and  $\Delta^{14}\text{C}_{\text{seagrass}}$  cannot be explained by any residual contamination from terrestrial organic carbon on the leaves because the terrestrial POC was  $^{14}\text{C}$ -depleted (mean  $\Delta^{14}\text{C}$  of terrestrial POC,  $-74.7 \pm 23.4$  ‰).

The significantly higher values in  $\Delta^{14}\text{C}_{\text{seagrass}}$  than  $\Delta^{14}\text{C}_{\text{DIC}}$  shows that the seagrass assimilates  $^{14}\text{C}$ -rich  $\text{C}_{\text{air}}$  ( $\Delta^{14}\text{C}$  around 17 ‰; Fig. 2a). The two-carbon-source mixing model indicated that the seagrass assimilated 0–40 % (mean  $\pm$ SD,  $17 \pm 12$  %) of its inorganic carbon as  $\text{C}_{\text{air}}$ ; the contribution was  $20 \pm 12$  % in the low-salinity zone (salinity, 12–15) and  $13 \pm 12$  % in the high-salinity zone (salinity, 25–29; Fig. 2b). The contribution of  $\text{C}_{\text{air}}$  as a carbon source varied greatly even between samples from the same station (Fig. 2b). Because we did not determine the exposure time of each shoot in this study, we are unable to quantify any relationship between the contribution of  $\text{C}_{\text{air}}$  and air exposure time; however, the exposure time would mediate the assimilation of  $\text{C}_{\text{air}}$  (Clavier et al., 2011).

As  $\Delta^{14}\text{C}_{\text{DIC}}$  was significantly lower than  $\Delta^{14}\text{C}_{\text{air}}$ , the contribution of  $\text{C}_{\text{air}}$  can be determined for May and July 2014 (Fig. 2a). This radiocarbon isotopic approach would be useful in the high latitudes of the Pacific Ocean where surface






**Figure 4.** Seasonal dynamics of ocean currents affecting the oceanic boundary of Furen Lagoon. Red circle shows the location of Furen Lagoon. (a) From mid-November to August, the oceanic boundary is the Oyashio ( $\Delta^{14}\text{C}_{\text{DIC}}$ ,  $-60$  to  $-10$ ‰; Aramaki et al., 2001). (b) From September to early November, the Soya warm current ( $\Delta^{14}\text{C}_{\text{DIC}}$ ,  $50$  to  $70$ ‰; Aramaki et al., 2007) intruded into the boundary.

seawater is  $^{14}\text{C}$ -depleted ( $\Delta^{14}\text{C}_{\text{DIC}} < 0$ ‰; Talley, 2007). In contrast, the  $\Delta^{14}\text{C}_{\text{DIC}}$  in surface seawater is generally higher than  $\Delta^{14}\text{C}_{\text{air}}$  in other regions of the Pacific Ocean because of bomb-derived  $^{14}\text{C}$  (Talley, 2007).

In any case, the  $\Delta^{14}\text{C}$  approach is potentially applicable to other regions by using the  $\Delta^{14}\text{C}$  gradient. However, the seasonal dynamics of  $\Delta^{14}\text{C}_{\text{DIC}}$  would affect the application of this approach because it is only applicable when the  $\Delta^{14}\text{C}$  values for end-members (seawater DIC, freshwater DIC, and  $\text{C}_{\text{air}}$ ) are distinct (not overlapping) as they were in May and July 2014 during this study. We could not use the  $\Delta^{14}\text{C}$  approach to quantify the  $\text{C}_{\text{air}}$  contribution in September or November 2014 in Furen Lagoon because the  $\Delta^{14}\text{C}_{\text{DIC}}$  of seawater increased to near  $\Delta^{14}\text{C}_{\text{air}}$  and there was overlap between the two (Fig. 3). The overlapping in the range of values, induced by variations in the  $\Delta^{14}\text{C}_{\text{DIC}}$  of seawater, was likely caused by the dynamics of the Oyashio (mean  $\Delta^{14}\text{C}_{\text{DIC}}$ ,  $-41$ ‰; Aramaki et al., 2001, 2007) and the Soya warm current (mean  $\Delta^{14}\text{C}_{\text{DIC}}$ ,  $52$ ‰; Aramaki et al., 2007; Kumamoto et al., 1998; Figs. 1 and 4). According to the dis-



**Figure 5.** (a) Distant and (b) close-up views of the seagrass leaves exposed to the air during low tide in Furen Lagoon, Japan. (c) Conceptual diagram of the uptake of atmospheric CO<sub>2</sub> ( $\text{C}_{\text{air}}$ ) across the surface-water film on the seagrass leaves during low tide (left), and the uptake of DIC during high tide (right).

tribution of sea surface temperature derived from the Moderate Resolution Imaging Spectroradiometers (MODIS) images in 2014 (Fig. 4; <http://oceancolor.gsfc.nasa.gov/cms/>), the oceanic boundary of Furen Lagoon was the Oyashio throughout the year except from late summer to autumn when the Soya warm current intrudes into the boundary (Oguma et al., 2008; Takizawa, 1982). The oceanic end-member  $\Delta^{14}\text{C}_{\text{DIC}}$  would reflect the value of the Oyashio from January to August 2014, when the Soya warm current did not reach the oceanic boundary of Furen Lagoon

(Fig. 4). The oceanic end-member  $\Delta^{14}\text{C}_{\text{DIC}}$  would, therefore, not overlap with  $\Delta^{14}\text{C}_{\text{air}}$  during January to August 2014, which includes the whole period of a 1–3-month seagrass leaf growing prior to sampling, i.e., February to July, indicating that the uptake of  $\text{C}_{\text{air}}$  by the seagrass is a robust estimate during the period. Even if the sporadic upwelling had occurred during the study period, our determination of the  $\text{C}_{\text{air}}$  contribution here would be underestimated because the  $\Delta^{14}\text{C}_{\text{DIC}}$  of the upwelling deep-sea water is lower than that of surface water (Aramaki et al., 2001, 2007). Nevertheless, the applicability of the  $\Delta^{14}\text{C}$  technique is dependent on the  $\Delta^{14}\text{C}$  dynamics of end-members.

Our  $\Delta^{14}\text{C}$  analysis considerably reduces the limitations and uncertainties of conventional methods such as that using only  $\delta^{13}\text{C}$  (Clavier et al., 2011; Cooper and McRoy, 1988; Raven et al., 2002). In particular, the use of  $\Delta^{14}\text{C}$  has the advantage of avoiding effects of isotopic fractionation (Stuiver and Polach, 1977); the use of  $\delta^{13}\text{C}$  does not and therefore generates large uncertainties. The  $\delta^{13}\text{C}$  of the seagrass was low ( $-14.0 \pm 2.4\text{‰}$ ) in the low-salinity zone (salinity, 12–15) and high ( $-8.8 \pm 1.9\text{‰}$ ) in the high-salinity zone (salinity, 25–29; Fig. 2c). There were significant correlations between salinity and  $\delta^{13}\text{C}$  of DIC,  $\text{HCO}_3^-$ ,  $\text{CO}_2(\text{aq})$  and the seagrass (Pearson's correlation coefficient:  $P < 0.001$ ; Fig. 2c, d). As the  $\delta^{13}\text{C}$  of  $\text{HCO}_3^-$  was isotopically distinct from  $\delta^{13}\text{C}$  of both  $\text{CO}_2(\text{aq})$  and  $\text{C}_{\text{air}}$  (Fig. 2d) and as *Z. marina* also uses  $\text{HCO}_3^-$  as a carbon source under low- $\text{CO}_2(\text{aq})$  conditions (Beer and Rehnberg, 1997), the  $\delta^{13}\text{C}$  of the seagrass should change depending on the contribution of  $\text{HCO}_3^-$  as a carbon source (Campbell and Fourqurean, 2009; Hemminga and Mateo, 1996; Raven et al., 2002). However, it is not possible to distinguish the contribution of  $\text{C}_{\text{air}}$  from that of other carbon sources because the  $\delta^{13}\text{C}$  of  $\text{C}_{\text{air}}$  overlapped those of both  $\text{HCO}_3^-$  and  $\text{CO}_2(\text{aq})$  (Fig. 2d). Furthermore,  $\delta^{13}\text{C}$  of both  $\text{HCO}_3^-$  and  $\text{CO}_2(\text{aq})$  changes through mixing between low- $\delta^{13}\text{C}$  river water and high- $\delta^{13}\text{C}$  seawater in brackish areas (Fig. 2d; Hemminga and Mateo, 1996; Simenstad and Wissmar, 1985).

In any case, there are large uncertainties when using  $\delta^{13}\text{C}$  to quantitatively estimate the contribution of  $\text{C}_{\text{air}}$  as a carbon source because the isotopic fractionation that occurs in the steps between the carbon source and organic plant compounds changes depending on the photosynthetic carbon demand (Hemminga and Mateo, 1996; Raven et al., 2002). The radiocarbon isotopic approach can avoid the uncertainties derived from both the chemical species of DIC and the isotopic fractionation factor in carbon assimilation.

The seagrass leaves assimilated  $\text{C}_{\text{air}}$  when exposed to air during low tide (Fig. 5).  $\text{CO}_2$  exchange between the air and water would occur at the very thin film of water on the air-exposed seagrass leaves (Fig. 5c), likely enhancing the passive uptake of  $\text{C}_{\text{air}}$  by diffusion. Our high estimate of the  $\text{C}_{\text{air}}$  contribution (mean, 17 %) was unexpected because prior works suggest that photosynthetic rates of seagrasses in in-

tertidal zones decrease during air exposure (Clavier et al., 2011), particularly in cases of desiccation (Leuschner et al., 1998). However, the leaves of subtidal seagrass are never desiccated because of the presence of the thin film of water, which reduces the negative effects of air exposure (i.e., desiccation).

The net ecosystem production of seagrass meadows is a key factor determining whether they are sinks or sources of  $\text{C}_{\text{air}}$  (Maher and Eyre, 2012; Tokoro et al., 2014; Watanabe and Kuwae, 2015). Previously, however, such an exchange of  $\text{CO}_2$  has been thought to occur only via the air–water interface with subsequent exchange with seagrasses as DIC. This study using radiocarbon isotope analysis demonstrates the assimilation of modern  $\text{C}_{\text{air}}$  by seagrass. Moreover, our radiocarbon isotopic approach has potential for application to other photoautotrophs living near the air–water interface, such as intertidal macroalgae and amphibious macrophytes. Other applications may include determining the origin of the DIC source (e.g., terrestrial or oceanic) in deeper seagrass systems. However, adequate determinations will require separation and stability in the end-member values (e.g., in oceanographic contexts and in the dynamics of  $\Delta^{14}\text{C}$  in coastal waters). The relative contribution of gas exchange via the air–seagrass water film to the total exchange is still unknown. To understand the role of seagrass meadows in the global carbon cycle, it will be necessary in future studies to precisely measure  $\text{CO}_2$  exchanges at both the air–water and air–seagrass water-film interfaces.

*Author contributions.* K. Watanabe and T. Kuwae designed this study, K. Watanabe carried out the field surveys and analyzed the data, and K. Watanabe and T. Kuwae wrote the paper.

*Acknowledgements.* We thank H. Moki, E. Miyoshi, S. Montani, and the members of Hokkaido University for help in field surveys, T. Tokoro for helpful comments, and K. Sakihara and A. Okuno for chemical analysis. This study was supported by a Canon Foundation grant, a Grant-in-Aid for Challenging Exploratory Researches (no. 24656316 and 26630251) from the Japan Society for the Promotion of Science, and a Core Research for Evolutional Science and Technology (CREST) Program of the Japan Science and Technology Agency.

Edited by: C. Heinze

## References

- Aramaki, T., Watanabe, S., Kuji, T., and Wakatsuchi, M.: The Okhotsk-Pacific seawater exchange in the viewpoint of vertical profiles of radiocarbon around the Bussol' Strait, *Geophys. Res. Lett.*, 28, 3971–3974, 2001.
- Aramaki, T., Senjyu, T., Togawa, O., Ootosaka, S., Suzuki, T., Kitamura, T., Amano, H., and Volkov, Y. N.: Circulation in the northern Japan Sea studied chiefly with radiocarbon, *Radiocarbon*, 49, 915–924, 2007.
- Beer, S. and Koch, E.: Photosynthesis of marine macroalgae and seagrasses in globally changing CO<sub>2</sub> environments, *Mar. Ecol.-Progr. Ser.*, 141, 199–204, 1996.
- Beer, S. and Rehnberg, J.: The acquisition of inorganic carbon by the seagrass *Zostera marina*, *Aquat. Bot.*, 56, 277–283, 1997.
- Beer, S., Björk, M., Hellblom, F., and Axelsson, L.: Inorganic carbon utilization in marine angiosperms (seagrasses), *Funct. Plant Biol.*, 29, 349–354, 2002.
- Campbell, J. E. and Fourqurean, J. W.: Interspecific variation in the elemental and stable isotopic content of seagrasses in South Florida, *Mar. Ecol.-Progr. Ser.*, 387, 109–123, 2009.
- Campbell, J. E. and Fourqurean, J. W.: Mechanisms of bicarbonate use influence the photosynthetic carbon dioxide sensitivity of tropical seagrasses, *Limnol. Oceanogr.*, 58, 839–848, 2013.
- Clavier, J., Chauvaud, L., Carlier, A., Amice, E., Van der Geest, M., Labrosse, P., Diagne, A., and Hily, C.: Aerial and underwater carbon metabolism of a *Zostera noltii* seagrass bed in the Banc d'Arguin, Mauritania, *Aquat. Bot.*, 95, 24–30, 2011.
- Cooper, L. W. and McRoy, C. P.: Stable carbon isotope ratio variations in marine macrophytes along intertidal gradients, *Oecologia*, 77, 238–241, 1988.
- Goñi, M. A., O'Connor, A. E., Kuzyk, Z. Z., Yunker, M. B., Gobeil, C., and Macdonald, R. W.: Distribution and sources of organic matter in surface marine sediments across the North American Arctic margin, *J. Geophys. Res.-Oceans*, 118, 4017–4035, 2013.
- Hellblom, F., Beer, S., Björk, M., and Axelsson, L.: A buffer sensitive inorganic carbon utilization system in *Zostera marina*, *Aquat. Bot.*, 69, 55–62, 2001.
- Hemminga, M. A. and Duarte, C. M.: Seagrass architectural features, in: *Seagrass Ecology*, edited by: Hemminga, M. A. and Duarte, C. M., Cambridge University Press, Cambridge, 27–64, 2000.
- Hemminga, M. A. and Mateo, M. A.: Stable carbon isotopes in seagrasses: variability in ratios and use in ecological studies, *Mar. Ecol.-Progr. Ser.*, 140, 285–298, 1996.
- Hosokawa, S., Nakamura, Y., and Kuwae, T.: Temperature induces shorter leaf life span in an aquatic plant, *Oikos*, 118, 1158–1163, 2009.
- Invers, O., Zimmerman, R. C., Alberte, R. S., Perez, M., and Romero, J.: Inorganic carbon sources for seagrass photosynthesis: an experimental evaluation of bicarbonate use in species inhabiting temperate waters, *J. Exp. Mar. Biol. Ecol.*, 265, 203–217, 2001.
- Ishikawa, N. F., Uchida, M., Shibata, Y., and Tayasu, I.: Carbon storage reservoirs in watersheds support stream food webs via periphyton production, *Ecology*, 95, 1264–1271, 2014.
- Ji, Y. and Tanaka, J.: Effect of desiccation on the photosynthesis of seaweeds from the intertidal zone in Honshu, Japan, *Phycol. Res.*, 50, 145–153, 2002.
- Jiang, Z., Huang, X., Zhang, J., Zhou, C., Lian, Z., and Ni, Z.: The effects of air exposure on the desiccation rate and photosynthetic activity of *Thalassia hemprichii* and *Enhalus acoroides*, *Mar. Biol.*, 161, 1051–1061, 2014.
- Kumamoto, Y., Yoneda, M., Shibata, Y., Kume, H., Tanaka, A., Uehiro, T., and Morita, M.: Direct observation of the rapid turnover of the Japan Sea bottom water by means of AMS radiocarbon measurement, *Geophys. Res. Lett.*, 25, 651–654, 1998.
- Larkum, A. W. D. and Den Hartog, C.: Evolution and biogeography of seagrasses, in: *Biology of seagrasses. Aquatic Plant Studies*, edited by: Larkum, A. W. D., McComb, A. J., and Shepherd, S. A., Elsevier, Amsterdam, 112–156, 1989.
- Leuschner, C. and Rees, U.: CO<sub>2</sub> gas exchange of two intertidal seagrass species, *Zostera marina* L. and *Zostera noltii* Hornem., during emersion, *Aquat. Bot.*, 45, 53–62, 1993.
- Leuschner, C., Landwehr, S., and Mehlig, U.: Limitation of carbon assimilation of intertidal *Zostera noltii* and *Z. marina* by desiccation at low tide, *Aquat. Bot.*, 62, 171–176, 1998.
- Maher, D. T. and Eyre, B. D.: Carbon budgets for three autotrophic Australian estuaries: Implications for global estimates of the coastal air-water CO<sub>2</sub> flux, *Global Biogeochem. Cy.*, 26, GB1032, doi:10.1029/2011GB004075, 2012.
- Oguma, S., Ono, T., Kusaka, A., Kasai, H., Kawasaki, Y., and Azumaya, T.: Isotopic tracers for water masses in the coastal regions of eastern Hokkaido, *J. Oceanogr.*, 64, 525–539, 2008.
- Plummer, L. N. and Busenberg, E.: The solubilities of calcite, aragonite and vaterite in CO<sub>2</sub>-H<sub>2</sub>O solutions between 0 and 90 °C, and an evaluation of the aqueous model for the system CaCO<sub>3</sub>-CO<sub>2</sub>-H<sub>2</sub>O, *Geochim. Cosmochim. Ac.*, 46, 1011–1040, 1982.
- Raven, J. A., Johnston, A. M., Kübler, J. E., Korb, R., McInroy, S. G., Handley, L. L., Scrimgeour, C. M., Walker, D. I., Beardall, J., Vanderklift, M., Fredriksen, S., and Dunton, K. H.: Mechanistic interpretation of carbon isotope discrimination by marine macroalgae and seagrasses, *Funct. Plant Biol.*, 29, 355–378, 2002.
- Simenstad, C. A. and Wissmar, R. C.: δ<sup>13</sup>C evidence of the origins and fates of organic carbon in estuarine and nearshore food webs, *Mar. Ecol.-Progr. Ser.*, 22, 141–152, 1985.
- Stuiver, M. and Braziunas, T. F.: Modeling atmospheric <sup>14</sup>C influences and <sup>14</sup>C ages of marine samples to 10 000 BC, *Radiocarbon*, 35, 137–189, 1993.
- Stuiver, M. and Polach, H. A.: Discussion: reporting of <sup>14</sup>C data, *Radiocarbon*, 19, 355–363, 1977.
- Takizawa, T.: Characteristics of the Soya warm current in the Okhotsk Sea, *J. Oceanogr. Soc. Japan*, 38, 281–292, 1982.
- Talley, L. D.: Hydrographic Atlas of the World Ocean Circulation Experiment (WOCE), Volume 2: Pacific Ocean, edited by: Sparrow, M., Chapman, P., and Gould, J., International WOCE Project Office, Southampton, 2007.
- Tokoro, T., Hosokawa, S., Miyoshi, E., Tada, K., Watanabe, K., Montani, S., Kayanne, H., and Kuwae, T.: Net uptake of atmospheric CO<sub>2</sub> by coastal submerged aquatic vegetation, *Global Change Biol.*, 20, 1873–1884, 2014.
- Watanabe, K. and Kuwae, T.: How organic carbon derived from multiple sources contributes to carbon sequestration processes in a shallow coastal system?, *Global Change Biol.*, 21, 2612–2623, 2015.

- Zeebe, R. E. and Wolf-Gladrow, D.: CO<sub>2</sub> in seawater: equilibrium, kinetics, and isotopes, in: Elsevier Oceanography Series 65, edited by: Halpern, D., Elsevier, Amsterdam, 346 pp., 2001.
- Zhang, J., Quay, P. D., and Wilbur, D. O.: Carbon isotope fractionation during gas-water exchange and dissolution of CO<sub>2</sub>, *Geochim. Cosmochim. Ac.*, 59, 107–114, 1995.
- Zimmerman, R. C., Kohrs, D. G., Steller, D. L., and Alberte, R. S.: Carbon partitioning in eelgrass: regulation by photosynthesis and the response to daily light–dark cycles, *Plant Physiol.*, 108, 1665–1671, 1995.