Biogeosciences, 5, 1739, 2008 www.biogeosciences.net/5/1739/2008/ © Author(s) 2008. This work is distributed under the Creative Commons Attribution 3.0 License.



Corrigendum to

"Peatlands and the carbon cycle: from local processes to global implications a synthesis" published in Biogeosciences, 5, 1475–1491, 2008

J. Limpens¹, F. Berendse¹, C. Blodau², J. G. Canadell³, C. Freeman⁴, J. Holden⁵, N. Roulet⁶, H. Rydin⁷, and G. Schaepman-Strub¹

In the introduction of the paper we provide several quantities of the amount of carbon stored in peatlands. The quantities indicated in 10¹² should in fact say 10¹⁵, and therefore they are not Teragrams (Tg) but Petagrams (Pg). All other quantities mentioned in the paper are in the correct unit.

Therefore the text in the introduction should read: Despite covering less than 3% of the Earth's land surface, boreal and subarctic peatlands store between 270 and 370 Pg C (1 Pg=1015 g) as peat (Turunen et al., 2002), which would amount to 34-46% of the 796 Pg C currently held in the atmosphere as CO_2 (IPCC, 2007). Tropical peatland C stores are estimated to be around 52 Pg C, with very large uncertainties (8–258 Pg C, Hooijer et al., 2006).



Correspondence to: J. Limpens (juul.limpens@wur.nl)

¹Nature Conservation and Plant Ecology Group, Wageningen University, Wageningen, The Netherlands

²Limnological Research Station and Department of Hydrology, University of Bayreuth, 95440 Bayreuth, Germany

³Global Carbon Project, CSIRO Marine and Atmospheric Research, GPO Box 3023, Canberra, ACT 2601, Australia

⁴School of Biological Sciences, Bangor University, Wales, LL57 2UW, UK

⁵School of Geography, University of Leeds, Leeds, LS2 9JT, UK

⁶Department of Geography, McGill University, 805, Sherbrooke Street West, Montréal, Québec, H3A 2K6, Canada

⁷Department of Plant Ecology, Evolutionary Biology Centre, Uppsala University, Villavägen 14, 752 36 Uppsala, Sweden