

Corrigendum to

“Incorporation of crop phenology in Simple Biosphere Model (SiBcrop) to improve land-atmosphere carbon exchanges from croplands” published in Biogeosciences, 6, 969–986, 2009

E. Lokupitiya¹, S. Denning¹, K. Paustian^{2,3}, I. Baker¹, K. Schaefer⁴, S. Verma⁵, T. Meyers⁶, C. J. Bernacchi⁷, A. Suyker⁵, and M. Fischer⁸

¹Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, USA

²Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO 80523, USA

³Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, USA

⁴National Snow and Ice Data Center, University of Colorado, Boulder, CO 80309, USA

⁵School of Natural Resources, University of Nebraska-Lincoln, Lincoln, NE 68583-0978, USA

⁶NOAA/ARL/ATDD, Oak Ridge, TN 37830-2456, USA

⁷Department of Plant Biology, University of Illinois at Urbana-Champaign, Champaign, IL 61820, USA

⁸Lawrence Berkeley National Laboratory, Environmental Energy Technologies Division,
Atmospheric Science Department, Berkeley, CA 94720, USA

In the above mentioned manuscript a mistake in Fig. 11 occurred. The corrected version of the figure is as follows.




Fig. 11. Evaluation of predicted NEE against the observed data at a rainfed site in Mead, NE with maize and soybean in rotation. (a) monthly comparison with original model; (b) monthly comparison with new model; (c) sub hourly comparison with new model in 2001; (d) diurnal variation in the sub hourly NEE in early July in 2001.



Correspondence to: E. Lokupitiya
(erandi@atmos.colostate.edu)