

Corrigendum to

“Estimating the suspended sediment yield in a river network by means of geomorphic parameters and regression relationships” published in Hydrol. Earth Syst. Sci., 12, 177–191, 2008

S. Grauso¹, G. Fattoruso², C. Crocetti³, and A. Montanari³

¹Dipartimento Ambiente, Cambiamenti Globali e Sviluppo Sostenibile, ENEA Centro Ricerche Casaccia, Roma, Italy

²Dipartimento Ambiente, Cambiamenti Globali e Sviluppo Sostenibile, ENEA Centro Ricerche Portici (Napoli), Italy

³Faculty of Engineering, University of Bologna, Italy

Due to an unexpected editing error, data referring to drainage density (D_d) and index of hierarchical anomaly (Δ_a), contained in Table 1 of the above cited article, are wrong.

The correct data, compliant with those actually reported in the reference work of Ciccacci et al. (1987), are here showed:

Table 1. Drainage density (D_d), hierarchical anomaly index (Δ_a) and contributing area (A) for the watersheds considered by Ciccacci et al. (1987).

Watershed	D_d [km ⁻¹]	Δ_a	A [km ²]
Trebbia	4.90	1.59	226
Enza	5.62	1.18	670
Idice	5.61	0.86	397
Senio	4.17	1.32	269
Foglia	4.99	1.39	603
Orcia	4.80	1.34	580
Tavo	3.14	1.38	109
Volturno	3.50	1.62	2015
C. S. Maria	2.21	1.14	60
Triolo	2.60	0.91	54
Casanova	2.70	0.36	52
Salsola	2.60	1.15	43
Vulgano	2.44	0.94	94
Celone	2.80	0.76	86
Venosa	2.76	1.08	261
Atella	3.65	1.17	158
Agri (Grumento)	2.42	1.1	278
Agri (Tarangelo)	3.06	1.4	507
Delia	2.59	0.57	140
Gornalunga	4.40	1.13	232
Range	2.21–5.62	0.36–1.62	43–2015



Correspondence to: S. Grauso
(sergio.grauso@enea.it)