



**Improving ETKF
using the
second-order
information**

G. Wu et al.

Improving the ensemble transform Kalman filter using a second-order Taylor approximation of the nonlinear observation operator

G. Wu¹, X. Zheng¹, L. Wang², X. Liang³, S. Zhang¹, and X. Zhang¹

¹State Key Laboratory of Remote Sensing Science, College of Global Change and Earth System Science, Beijing Normal University, Beijing, China

²Department of Statistics, University of Manitoba, Winnipeg, Canada

³National Meteorological Information Center, China Meteorological Administration, Beijing, China

Received: 28 January 2014 – Accepted: 12 March 2014 – Published: 11 April 2014

Correspondence to: X. Zheng (x.zheng@bnu.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union & American Geophysical Union.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The observation operators for indirect observations are often nonlinear. For example, radiative transfer codes (e.g., RTTOV, CRTM, Saunders et al., 1999; Han et al., 2006) can be treated as observation operators by mapping air temperature and moisture to the microwave radio brightness temperature (McNally, 2009). Because the relationship of these observations with modelled variables may be strongly nonlinear (Liou, 2002), and because the observation errors may be spatially correlated (Miyoshi et al., 2013), data assimilation schemes have to be appropriately designed to address such indirect observations.

Most data assimilation methods are fundamentally based on linear theory but have different responses to departures from linearity (Lawson and Hansen, 2004). Conceptually, variational data assimilation schemes (VAR, e.g., Parrish and Derber, 1992; Courtier et al., 1994; Lorenc, 2003) can assimilate data with nonlinear observation operators and spatially correlated observation errors. However, a drawback of VAR is that it has to calculate the adjoint of a dynamical model, which is not an easy task in practice. Moreover, VAR does not give a direct estimate of the background error covariance matrix, which is crucial for the performance of any data assimilation scheme.

The Ensemble Kalman Filter (EnKF) scheme has a strategy to optimize forecast error statistics without using the adjoint of the dynamical model (e.g., Evensen, 1994a, 1994b; Burgess et al., 1998; Anderson and Anderson, 1999; Wang and Bishop, 2003; Wu et al., 2013). It is also conceptually applicable to data assimilation with nonlinear observation operators. However, it has been demonstrated that when the observation operator is strongly nonlinear, using the linear approximation of the observation operator to derive the error covariance evolution equation can result in an oversimplified closure and dubious performance of the EnKF (e.g., Miller et al., 1994; Evensen, 1997; Yang et al., 2012).

The Ensemble Transform Kalman Filter (ETKF) was first introduced in atmospheric assimilation by Bishop and Toth (1999) and Bishop et al. (2001). Wang and Bishop (2003) transformed the forecast perturbations into analysis perturbations by multiplying a transformation matrix. They also proposed an efficient way to construct the transform

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

matrix through eigenvector decomposition of a matrix with the ensemble size. Hunt et al. (2007) extended the ETKF method to address a general nonlinear observation operator using the cost function, similar to Lorenc (2003). However, Hunt et al. (2007) minimized the weight vector of the ensemble analysis state instead of the analysis state, as in Lorenc (2003). In addition to the reduction of the computational cost compared with EnKF, another advantage of the ETKF proposed by Hunt et al. (2007) is that it can assimilate observations with strongly nonlinear observation operators (Chen et al., 2009) and with spatially correlated observation errors (Stewart et al., 2013), similar to VAR.

However, there are still problems associated with the ETKF when the observation operator is strongly nonlinear. First, the current ETKF is based on the minimization of a cost function similar to that in VAR for nonlinear observation operators (Hunt et al., 2007). The direct calculation for the minima may be computationally expensive because the nonlinear operators and their tangent-linear operators have to be iteratively calculated. Using the linear approximation of the nonlinear observation operators (e.g. Hunt et al., 2007) can effectively reduce the computational burden, but at the cost of increasing analysis error. Second, tangent-linear approximation of the observation operator is used for the forecast error inflation in the ETKF (e.g., Li et al., 2009). If the observation operators are strongly nonlinear, the inflation factors and hence the forecast error covariance matrices may be estimated erroneously in this way, leading to an eventual increase in the analysis error.

In this study, we propose two alternative approaches to improving assimilation quality when the observation operator is strongly nonlinear. First, in an effort to reduce computational cost without significantly reducing estimation quality, we use the second-order Taylor expansion of the observation operator to estimate both the inflation factors and the analysis states. Second, for the case where the inflation factor is constant in space, we propose a new forecast error inflation method that can address a general nonlinear observation operator without using the tangent-linear operator.

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The potential use of the second-order information has been noted by some authors. For example, Hunt et al. (2007) noted that the second-order derivatives of the objective function might be used to estimate the covariance of analysis weight, which is an important step in ETKF with a nonlinear observation operator. Moreover, Le Dimet et al. (2002) and Daescu and Navon (2007) noted that the second-order information in nonlinear variational data assimilation is important to the issue of solution uniqueness.

In the conventional ETKF scheme, the linear approximation of nonlinear observation operators is used for the purpose of reducing the computational cost compared with conventional methods of searching the minima of nonlinear cost functions (Hunt et al., 2007). This study also aims to investigate the changes of analysis errors when a nonlinear observation operator is substituted by its first-order and second-order Taylor approximations. We focus on the formulation of the forecast error inflation method in the case of a nonlinear observation operator and on improved accuracy with the second-order vs. the first-order approximation and the linear approximation. Further studies on the performance of the proposed schemes in practical data assimilations are needed and should be performed in the future.

The remainder of the paper is organized as follows. Our modified ETKF schemes are described in Sect. 2. The assimilation results on a Lorenz-96 model with a nonlinear observation system are presented in Sect. 3. The discussions are given in Sect. 4, and conclusions are presented in Sect. 5.

2 Methodology

2.1 ETKF with forecast error inflation

Hunt et al. (2007) gave a comprehensive description of the ETKF with a nonlinear observation operator, but without procedures for forecast error inflation. In this section, we propose an inflation scheme to address a general nonlinear observation operator.

Using the notations of Ide et al. (1997), a nonlinear discrete-time forecast and observation system can be written as

$$\mathbf{x}_i^f = M_{i-1}(\mathbf{x}_{i-1}^a) + \boldsymbol{\eta}_i, \quad (1)$$

$$\mathbf{y}_i^o = H_i(\mathbf{x}_i^f) + \boldsymbol{\varepsilon}_i, \quad (2)$$

where i is the time step index; $\mathbf{x}_i^f = \{x_i^f(1), x_i^f(2), \dots, x_i^f(n)\}^T$ is the n -dimensional true state vector; $\mathbf{x}_{i-1}^a = \{x_{i-1}^a(1), x_{i-1}^a(2), \dots, x_{i-1}^a(n)\}^T$ is the n -dimensional analysis state vector, which is an estimate of \mathbf{x}_{i-1}^f ; M_i is the nonlinear forecast operator; $\mathbf{y}_i^o = \{y_i^o(1), y_i^o(2), \dots, y_i^o(p_i)\}^T$ is the p_i -dimensional observation vector; $H_i = \{h_i(1), h_i(2), \dots, h_i(p_i)\}^T$ is the nonlinear observation operator, where $h_i(k)$ is a n -dimensional multivariate function; and $\boldsymbol{\eta}_i$ and $\boldsymbol{\varepsilon}_i$ are the forecast and observation error vectors, which are assumed to be statistically independent of each other, time-uncorrelated, and to have mean zero and covariance matrices \mathbf{P}_i and \mathbf{R}_i , respectively. The detailed procedure of the ETKF with a nonlinear observation operator (Hunt et al., 2007) with the proposed inflation scheme is as follows.

Step 1. Calculate the j th perturbed forecast state at time i as

$$\mathbf{x}_{i,j}^f = M_{i-1}(\mathbf{x}_{i-1,j}^a), \quad (3)$$

where $\mathbf{x}_{i-1,j}^a$ is the j th perturbed analysis state at time $i-1$. Then, the mean forecast state is defined as

$$\mathbf{x}_i^f = \frac{1}{m} \sum_{j=1}^m \mathbf{x}_{i,j}^f, \quad (4)$$

where m is the total number of ensemble members.

Step 2. Assume the forecast errors to be in the form $\sqrt{\lambda_i}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)$, ($j = 1, 2, \dots, m$), where the inflation factor λ_i can be estimated by minimizing the objective function

$$L_i(\lambda) = \text{Tr} \left[\left(\mathbf{d}_i \mathbf{d}_i^T - \mathbf{C}_i(\lambda) - \mathbf{I} \right) \left(\mathbf{d}_i \mathbf{d}_i^T - \mathbf{C}_i(\lambda) - \mathbf{I} \right)^T \right]. \quad (5)$$

**Improving ETKF
using the
second-order
information**

G. Wu et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Here, \mathbf{I} is the $m \times m$ identity matrix,

$$\mathbf{d}_i = \mathbf{R}_i^{-1/2} \left(\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) \right). \quad (6)$$

is the innovation vector normalized by the square root of the observation error covariance matrix (Wang and Bishop, 2003), and

$$\mathbf{C}_i(\lambda) \equiv \frac{1}{m-1} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left(H_i(\mathbf{x}_i^f + \sqrt{\lambda}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)) - H_i(\mathbf{x}_i^f) \right) \right. \\ \left. \left(H_i(\mathbf{x}_i^f + \sqrt{\lambda}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)) - H_i(\mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right]. \quad (7)$$

(See Appendix A for details).

Step 3. Calculate the analysis state as

$$\mathbf{x}_i^a = \mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}_i^a \quad (8)$$

where

$$\mathbf{X}_i^f = (\mathbf{x}_{i,1}^f - \mathbf{x}_i^f, \mathbf{x}_{i,2}^f - \mathbf{x}_i^f, \dots, \mathbf{x}_{i,m}^f - \mathbf{x}_i^f) \quad (9)$$

and \mathbf{w}_i^a is estimated by minimizing the objective function

$$J_i(\mathbf{w}) = \frac{1}{2}(m-1)\mathbf{w}^T \mathbf{w} + \frac{1}{2} \left[\mathbf{y}_i^o - H_i \left(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right) \right]^T \\ \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i \left(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right) \right]. \quad (10)$$

Step 4. Calculate a perturbed analysis state as

$$\mathbf{x}_{i,j}^a = \mathbf{x}_i^a + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}_{i,j}^a \quad (11)$$

where $\mathbf{W}_{i,j}^a$ is the j th column of the matrix $\mathbf{W}_i^a = \sqrt{m-1}(\ddot{J}_{i|\mathbf{w}_i^a})^{-1/2}$ and $\ddot{J}_{i|\mathbf{w}_i^a}$ is the second-order derivative of $J_i(\mathbf{w})$ at \mathbf{w}_i^a (see Appendix B for details). Lastly, set $i = i + 1$ and return to Step 1 for the next iteration.

For estimating the inflation factor, Li et al. (2009) proposed a scheme, for which the tangent-linear operator of the observation operator (see Sect. 2.2.1 for the definition) is required. In the effort to reduce computational cost of searching the minima of the objective function (Eq. 10), Hunt et al. (2007) suggested the following linear approximation,

$$H_i \left(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right) \approx H_i(\mathbf{x}_i^f) + \mathbf{Y}_i^f \mathbf{w} \quad (12)$$

where

$$\mathbf{Y}_i^f = \left(H_i \left(\sqrt{\hat{\lambda}_i} (\mathbf{x}_{i,1}^f - \mathbf{x}_i^f) + \mathbf{x}_i^f \right) - H_i(\mathbf{x}_i^f), H_i \left(\sqrt{\hat{\lambda}_i} (\mathbf{x}_{i,2}^f - \mathbf{x}_i^f) + \mathbf{x}_i^f \right) - H_i(\mathbf{x}_i^f), \dots, H_i \left(\sqrt{\hat{\lambda}_i} (\mathbf{x}_{i,m}^f - \mathbf{x}_i^f) + \mathbf{x}_i^f \right) - H_i(\mathbf{x}_i^f) \right). \quad (13)$$

In this study, this traditional ETKF approach is validated against the other approaches.

2.2 Simplified estimation methods in special cases

To compute the variational minimization in Eq. (10) operationally, one can directly compute the explicit solution of the minima and iterate the process as in the 4D-Var outer loop (Lorenc, 2003; Liu et al., 2008). However, doing so still requires repeatedly calculating the nonlinear function $H_i(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w})$ and its tangent-linear operator (see Sect. 2.2.1 for the definition) which depend on \mathbf{w} and \mathbf{x}_i^f . In this subsection, we show an alternative procedure when the observation operator H_i can be approximated by its Taylor expansions.

2.2.1 First-order Taylor approximation for H_i

The first-order Taylor approximation for H_i at the forecast state vector \mathbf{x}_i^f is defined as

$$H_i(\mathbf{x}_i^t) \approx H_i(\mathbf{x}_i^f) + \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}(\mathbf{x}_i^t - \mathbf{x}_i^f), \quad (14)$$

5 where

$$\dot{\mathbf{H}}_{i|\mathbf{x}_i^f} = \left(\begin{array}{ccc} \frac{\partial h_i(1)}{\partial x_i(1)} & \cdots & \frac{\partial h_i(1)}{\partial x_i(n)} \\ \vdots & \ddots & \vdots \\ \frac{\partial h_i(p_i)}{\partial x_i(1)} & \cdots & \frac{\partial h_i(p_i)}{\partial x_i(n)} \end{array} \right) \bigg|_{\mathbf{x}_i = \mathbf{x}_i^f} \quad (15)$$

is the first-order derivative of H_i evaluated at the forecast state \mathbf{x}_i^f (tangent-linear operator). Then, λ_i can be estimated by minimizing the quadratic function

$$10 \quad L_{1,i}(\lambda) = \text{Tr} \left[(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \mathbf{I}) (\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \mathbf{I})^T \right]. \quad (16)$$

The analytic solution is

$$15 \quad \hat{\lambda}_i = \frac{\text{Tr} \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} (\mathbf{d}_i \mathbf{d}_i^T - \mathbf{I})^T \right]}{\text{Tr} \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} \right]}, \quad (17)$$

where

$$\hat{\mathbf{P}}_i = \mathbf{X}_i^f (\mathbf{X}_i^f)^T / (m - 1). \quad (18)$$

Similarly, \mathbf{w}_i^a can be estimated by minimizing the multivariate quadratic function

$$J_{1,i}(\mathbf{w}) = \frac{1}{2}(m-1)\mathbf{w}^T \mathbf{w} + \frac{1}{2} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} \right]^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} \right] \quad (19)$$

5 and the analytic solution is

$$\mathbf{w}_i^a = ((m-1)\mathbf{I} + \hat{\lambda}(\mathbf{X}_i^f)^T \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f)^{-1} \sqrt{\hat{\lambda}_i} (\mathbf{X}_i^f)^T \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^f)). \quad (20)$$

(see Appendix C for details).

2.2.2 Second-order Taylor approximation for H_i

10 The second-order Taylor approximation for H_i at \mathbf{x}_i^f is defined as

$$H_i(\mathbf{x}_i^t) \approx H_i(\mathbf{x}_i^f) + \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} (\mathbf{x}_i^t - \mathbf{x}_i^f) + \frac{1}{2} ((\mathbf{x}_i^t - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} (\mathbf{x}_i^t - \mathbf{x}_i^f)), \quad (21)$$

where $\ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \equiv \{\ddot{\mathbf{H}}_{i|\mathbf{x}_i^f}(1), \dots, \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f}(\rho_i)\}^T$ is the second-order derivative of H_i at \mathbf{x}_i^f ,

$$\ddot{\mathbf{H}}_{i|\mathbf{x}_i^f}(k) \equiv \left(\begin{array}{ccc} \frac{\partial^2 h_i(k)}{\partial x_i(1) \partial x_i(1)} & \cdots & \frac{\partial^2 h_i(k)}{\partial x_i(1) \partial x_i(n)} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 h_i(k)}{\partial x_i(n) \partial x_i(1)} & \cdots & \frac{\partial^2 h_i(k)}{\partial x_i(n) \partial x_i(n)} \end{array} \right) \bigg|_{\mathbf{x}_i = \mathbf{x}_i^f} \quad k = 1, \dots, \rho_i, \quad (22)$$

15

and \otimes is the outer product operator, i.e., for two arbitrary n -dimensional vectors \mathbf{x} and \mathbf{y} ,

$$\mathbf{x}^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{y} = \{\mathbf{x}^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f}(1) \mathbf{y}, \dots, \mathbf{x}^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f}(\rho_i) \mathbf{y}\}^T, \quad (23)$$

is a p_i -dimensional vector. Then, λ_i can be estimated by minimizing the polynomial objective function of $\lambda^{1/2}$

$$L_{2,i}(\lambda) = \text{Tr} \left[\left(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \lambda^{3/2} \mathbf{C}_{1,i} - \lambda^{3/2} \mathbf{C}_{1,i}^T - \lambda^2 \mathbf{C}_{2,i} - \mathbf{I} \right) \right. \\ \left. \left(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \lambda^{3/2} \mathbf{C}_{1,i} - \lambda^{3/2} \mathbf{C}_{1,i}^T - \lambda^2 \mathbf{C}_{2,i} - \mathbf{I} \right)^T \right], \quad (24)$$

where

$$\mathbf{C}_{1,i} = \frac{1}{2(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda_i} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right. \\ \left. \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right], \quad (25)$$

and

$$\mathbf{C}_{2,i} = \frac{1}{4(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \right. \\ \left. \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right]. \quad (26)$$

Moreover, \mathbf{w}_i^a can be estimated by minimizing the multivariate polynomial objective function

$$J_{2,i}(\mathbf{w}) \approx \frac{1}{2} (m-1) \mathbf{w}^T \mathbf{w} \\ + \frac{1}{2} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} - \frac{\hat{\lambda}_i}{2} \left((\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) \right) \right]^T \\ \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} - \frac{\hat{\lambda}_i}{2} \left((\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) \right) \right] \quad (27)$$

Improving ETKF
using the
second-order
information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(see Appendix D for details).

2.3 Validation statistics

In the following experiments, the “true” state \mathbf{x}_i^t is known by experimental design and is non-dimensional. In this case, we can use the Root Mean Square Error of the Analysis state (A-RMSE) to evaluate the accuracy of the assimilation results. The A-RMSE at the i th step is defined as

$$\text{A-RMSE} = \sqrt{\frac{1}{n} \|\mathbf{x}_i^a - \mathbf{x}_i^t\|^2}, \quad (28)$$

where $\|\cdot\|$ denotes the Euclidean norm and n is the dimension of the state vector. A smaller A-RMSE indicates a better performance of the assimilation scheme.

Following Anderson (2007) and Liang et al. (2012), the Root Mean Square Error of the Forecast state (F-RMSE) and the Spread of the Forecast state (F-Spread) at the i th step are defined as

$$\text{F-RMSE} = \sqrt{\frac{1}{n} \|\mathbf{x}_i^f - \mathbf{x}_i^t\|^2}. \quad (29)$$

and

$$\text{F-Spread} = \sqrt{\frac{1}{n(m-1)} \sum_{j=1}^m \|\mathbf{x}_{i,j}^f - \mathbf{x}_i^f\|^2}. \quad (30)$$

Roughly speaking, if $\mathbf{x}_{i,j}^f$ and \mathbf{x}_i^f are identically distributed with a mean value of \mathbf{x}_i^f , then F-RMSE and F-Spread should be consistent with each other. This is more likely the case if the model error is small. In general, the F-RMSE can be decomposed into an F-Spread component and a model error component, so it is larger than F-Spread (see Appendix B of Wu et al. (2013) for a detailed proof).

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Experiments with the Lorenz-96 model

In Sect. 2.1, we outlined the general ETKF assimilation scheme with Second-order Least Squares (SLS) error covariance matrix inflation. In Sect. 2.2, we proposed simplified estimation methods for two special cases: when H_i is tangent-linear (Sect. 2.2.1) and when H_i can be approximated by the second-order Taylor expansion (Sect. 2.2.2). In this section, we apply these assimilation schemes to the Lorenz-96 model (Lorenz, 1996) with model errors and a nonlinear observation system because it is a nonlinear dynamical system with properties relevant to realistic forecast problems.

3.1 Description of the dynamic and observation system

The Lorenz-96 model (Lorenz, 1996) is a strongly nonlinear dynamical system with quadratic nonlinearity governed by the equation

$$\frac{dX_k}{dt} = (X_{k+1} - X_{k-2})X_{k-1} - X_k + F, \quad (31)$$

where $k = 1, 2, \dots, K$ ($K = 40$, so there are 40 variables). We apply the cyclic boundary conditions $X_{-1} = X_{K-1}$, $X_0 = X_K$, $X_{K+1} = X_1$. The dynamics of Eq. (31) are “atmosphere-like” in that the three terms on the right-hand side consist of a nonlinear advection-like term, a damping term and an external forcing term, respectively. These terms can be thought of as a given atmospheric quantity (e.g., zonal wind speed) distributed on a latitude circle.

We solve Eq. (31) using the fourth-order Runge–Kutta time integration scheme (Butcher, 2003) with a time step of 0.05 non-dimensional units to derive the true state. This is roughly equivalent to 6 h in real time, assuming that the characteristic time-scale of the dissipation in the atmosphere is 5 days (Lorenz, 1996). In our assimilation schemes, we set $F = 8$ so that the leading Lyapunov exponent implies an error-doubling time of approximately 8 time steps (i.e., 0.4 non-dimensional time units) and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the fractal dimension of the attractor is 27.1 (Lorenz and Emanuel, 1998). The initial condition is chosen to be $X_k = F$ when $k \neq 20$ and $X_{20} = 1.001F$.

Because the microwave brightness temperature is an exponential function of soil temperature, we use the exponential observation function to mimic the radiative transfer model in this study. Suppose the synthetic observation generated at the k th model grid point is

$$y_j^o(k) = x_j^t(k) \exp \{ \alpha x_j^t(k) \} + \varepsilon_j(k), \quad (32)$$

where $k = 1, \dots, p_j$, and $\varepsilon_j = \{ \varepsilon_j(1), \varepsilon_j(2), \dots, \varepsilon_j(p_j) \}^T$ is the observation error vector with a mean of zero and covariance matrix \mathbf{R}_j . Here, α is a parameter controlling the nonlinearity of the observation operator, and $\alpha = 0$ corresponds to the linear case. All the 40 model variables are observed in our experiments. Suppose the observation errors are spatially correlated. The leading-diagonal elements of \mathbf{R}_j are $\sigma_0^2 = 1$, and the off-diagonal elements at site pair (j, k) are

$$\mathbf{R}_j(j, k) = \sigma_0^2 \times 0.5^{\min(|j-k|, 40-|j-k|)}, \quad (33)$$

With the exponential observation function and spatially correlated observation errors, the proposed scheme may potentially be applied to assimilate remote sensing observations and radiance data.

We added model errors in the Lorenz-96 model because they are inevitable in real dynamic systems. The model is a forced dissipative model with a parameter F that controls the strength of the forcing (Eq. 31). It behaves quite differently, with different values of F , and it produces chaotic systems with integer values of F larger than 3. Thus, we used various values of F to simulate a wide range of model errors while retaining $F = 8$ when generating the “true” state. These observations were then assimilated with $F = 4, 5, \dots, 12$. We simulated observations every 4 time steps for 100 000 steps to ensure robust results (Sakov and Oke, 2008; Oke et al., 2009). The ensemble size is 30.

Improving ETKF using the second-order information

G. Wu et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



3.2 Assimilation results

In this section, we examine the following five data assimilation methods corresponding to five different treatments of nonlinearity in inflation factor estimation and optimization:

ETKF: Traditional ETKF described in the last paragraph of Sect. 2.1.

5 TT: Tangent-linear approximation in both inflation (Eq. 17) and optimization (Eq. 20)

TN: Tangent-linear approximation in inflation (Eq. 17) and nonlinearity in optimization (Eq. 10)

SS: Second-order approximation in both inflation (Eq. 24) and optimization (Eq. 27)

NN: Nonlinearity in both inflation (Eq. 5) and optimization (Eq. 10).

10 The corresponding time-mean A-RMSEs of these assimilation schemes with $\alpha = 0.1$ and $F = 4, 5, \dots, 12$, over 100 000 time steps are plotted in Fig. 1. First, the figure clearly shows that for each estimation method, the A-RMSE increases as F becomes increasingly distant from the true value of 8.

Moreover, method NN has a smaller A-RMSE uniformly over all values of F than method TN, indicating that the proposed nonlinear inflation estimation (Eq. 5) performs better than the tangent-linear inflation scheme (Eq. 17). On the other hand, the A-RMSEs of methods SS and TN are close and smaller than that of method TT, suggesting that the second-order Taylor approximation method is comparable to the partial nonlinear method and is better than the first-order Taylor approximation method. Lastly, the traditional ETKF method has the largest A-RMSE, which implies that although the linear approximation is computationally more efficient, it may introduce larger analysis error.

25 For the Lorenz-96 model with large error ($F = 12$), the time-mean A-RMSEs and F-RMSEs of the five methods are given in Table 1 as well as the time-mean values of the objective functions. It can be seen that the full nonlinear method (NN) has both the smallest A-RMSE and F-RMSE, while the traditional linear approximation method (ETKF) has the largest RMSEs. The second-order Taylor approximation method (SS) performs similarly to the partial nonlinear method (TN), but better than the first-order

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Taylor approximation method (TT). In all cases, a smaller error corresponds to a smaller value of the objective function L .

To investigate the consistency between F-RMSE and F-Spread, we present the time-mean values of the five methods for cases $F = 12$ and $F = 8$ in Tables 2 and 3, respectively, as well as the ratios of F-RMSE over F-Spread. It is easy to see that in all cases, the F-RMSEs are larger than F-Spreads, and therefore, all ratios are greater than 1. However, the ratio of the full nonlinear method (NN) is the smallest, while the ratio of the linear approximation method is the largest. The ratio of the second-order approximation method (SS) is comparable to that of the partial nonlinear method (TN), but smaller than that of the first-order approximation method (TT). This suggests that the ensemble perturbed predictions are the most (least) reasonable for method NN (ETKF). Moreover, the ratios with $F = 8$ are much closer to 1 than those with $F = 12$ because the model error with $F = 12$ is much larger than that with $F = 8$ (see Sect. 2.3).

3.3 Impacts of Taylor approximations

In Sect. 3.2, we see that the A-RMSEs derived from the five ETKF assimilation schemes are close when F is close to the true value of 8 but are different when F departs from 8. This effect may depend on how well the Taylor expansions approximate the nonlinear observation operator H_i .

For example, the Taylor expansion of the k th component of observation operator $H_i(\mathbf{x}) = \mathbf{x} \exp\{\alpha \mathbf{x}\}$ (Eq. 32) with $\alpha = 0.1$ around the forecast state $x_i^f(k)$ is

$$\begin{aligned} x_i^t(k) \exp\{0.1x_i^t(k)\} &= x_i^f(k) \exp\{0.1x_i^f(k)\} \\ &+ (1 + 0.1x_i^f(k)) \exp\{0.1x_i^f(k)\} (x_i^t(k) - x_i^f(k)) \\ &+ (0.2 + 0.01x_i^f(k)) \exp\{0.1x_i^f(k)\} (x_i^t(k) - x_i^f(k))^2 + \dots \end{aligned} \quad (34)$$

To verify how well the Taylor expansions approximate the nonlinear observation operator H_i , we calculate the ratios of the Taylor expansion residuals over

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Improving ETKF
using the
second-order
information**

G. Wu et al.

[Title Page](#)

Abstract	Introduction
Conclusions	References
Tables	Figures

⏪	⏩
◀	▶
Back	Close

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

$x_i^t(k) \exp\{0.1x_i^t(k)\}$. If a ratio falls outside the interval $[-0.1, 0.1]$, then the corresponding residual cannot be regarded as being of a higher order infinitesimal and, therefore, cannot be ignored. Therefore, a larger proportion of the ratios falling outside the interval $[-0.1, 0.1]$ indicates a worse Taylor expansion and vice versa.

5 The proportions of the ratios that fall outside the interval $[-0.1, 0.1]$ are plotted in Fig. 2, which shows that when $F = 8$, the proportions are 0.0169 and 0.0006 for the first-order and second-order Taylor expansions, respectively. This result indicates that at almost all times and locations, both the first-order and second-order Taylor ex-
 10 pansions are good approximations of $x_i^t(k) \exp\{0.1x_i^t(k)\}$. However, when $F = 12$, at approximately 47% (19%) of the times and locations, $x_i^t(k) \exp\{0.1x_i^t(k)\}$ cannot be adequately approximated by its first (second) order Taylor expansion. Therefore, the A-RMSEs derived by the five ETKF schemes are quite different. This example also indicates that the success of the Taylor approximation method depends on both the
 15 smoothness of H_i and the range of forecast states. It seems that for the same strongly nonlinear observation operator, the larger the model error, the less success of the Taylor approximation.

4 Discussions

4.1 Inflation

It is widely recognized that the initial estimates of ensemble forecast errors should be
 20 inflated to improve assimilated results. To date, however, all of the existing adaptive inflation schemes in ETKF are based on the assumption that the observation operator is linear or tangent-linear (e.g., Li et al., 2009; Miyoshi, 2011). In this study, a method to estimate the multiplicative inflation factors is proposed for general nonlinear obser-
 25 vation operators.

Historically, in systems such as the Met Office ETKF (Flowerdew and Bowler, 2011), the need for inflation arises primarily due to spurious correlations that cause the raw



Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

analysis ensemble to be severely under-spread even when the background ensemble is well-spread. In this case, therefore, inflation must be applied to the analysis ensemble to correctly respond to the actual analysis uncertainty in the nonlinear forecast step. Inflation of the background ensemble may be more appropriate when the inflation primarily represents forecast model error, although stochastic physics or additive inflation may be more appropriate in this case (Hamill and Whitaker, 2005; Wu et al., 2013).

Our choice to inflate the background ensemble is crucial to the ability of finding a direct nonlinear solution for Eqs. (5)–(7) because of the way the inflation factor appears in these equations. Our objective function for estimating the multiplicative inflation factors is the second-order distance between the expectations of the squared innovation and its realization, which also makes the rms spread equal to the rms error (e.g., Palmer et al., 2006; Wang and Bishop, 2003; Flowerdew and Bowler, 2011).

The proposed nonlinear method is tested using the Lorenz-96 model with nonlinear observation systems (Sect. 3.2). The resulting A-RMSEs are clearly smaller than those of the first-order Taylor approximation in the estimation of the inflation factor. This indicates that the proposed full nonlinear inflation method is better than the first-order Taylor approximation inflation method in the case of nonlinear observation operators (i.e., method NN is better than method TN). In addition, the F-RMSE and the F-Spread of the proposed nonlinear method are more consistent than those of the first-order Taylor approximation method.

The proposed inflation method works well in the case where observation errors are spatially correlated. Some data assimilation schemes assume the observation error covariance matrix to be diagonal for simplicity and ease of computation (e.g., Anderson, 2007, 2009). However, because satellite observations often contain significantly correlated errors, the observation error covariance matrix has nonzero off-diagonal entries (Miyoshi et al., 2013). The inflation method proposed in this study can be applied to assimilate such observations.

In many practical experiments, the inflation factor is constant in time and is chosen by trial and error to give the assimilation with the most favourable statistics (e.g. Anderson

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and Anderson, 1999). For testing the empirical tuning method, the most accurate approach (i.e. estimate the minima of the objective function, Eq. 10), and the statistics root mean square errors of analysis-minus-observation and forecast minus-observation are used to tune the inflation factor. The A-RMSEs are estimated as 2.97 and 2.85 respectively which are larger than that of method SS (2.29). The ratios of F-RMSE to F-Spread are estimated as 3.14 and 3.45 respectively, which are also larger than that of method SS (1.80). All these facts indicate that the empirical estimation method for the inflation factor is not as good as our proposed method in this experimentation.

4.2 Second-order Taylor approximation

In Sect. 3.2, we showed that the ETKF scheme equipped with our proposed nonlinear inflation method leads to the smallest A-RMSE in all ETKF schemes analysed in this study. However, this ETKF scheme requires repeated calculation of the nonlinear observation functions $H_i(\mathbf{x}_i^f + \sqrt{\lambda}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f))$ and $H_i(\mathbf{x}_i^f + \sqrt{\hat{\lambda}}\mathbf{X}_i^f\mathbf{w})$ when minimizing the objective functions $L_j(\lambda)$ and $J_j(\mathbf{w})$, which can be computationally expensive. To reduce the computational cost, a commonly used approach is to substitute H_i by its tangent-linear operator (i.e., first-order Taylor expansion). However, this approach comes at the cost of losing estimation quality, as we have shown in this study.

As an effort to strike a balance between the estimation quality and computational cost, the nonlinear observation operator H_i in the objective functions $L_j(\lambda)$ and $J_j(\mathbf{w})$ is substituted by its second-order Taylor expansion. This is because (1) the second-order Taylor expansion is a better approximation of H_i than its tangent-linear operator; (2) with second-order Taylor expansion, the inflation factor λ and the weight vector \mathbf{w} are concentrated out of H_i , so the objective functions (Eqs. 24 and 27) become polynomials, for which a minima is easier to derive; and (3) the second-order derivative of H_i is required for estimating ensemble analysis states (Eq. 11) in the ETKF scheme, so its computation is not an additional task.

Improving ETKF using the second-order information

G. Wu et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

The accuracy of the ETKF scheme with the second-order Taylor approximation is examined in Sect. 3.2. The results suggest that the scheme is more accurate than the ETKF scheme based on the first-order Taylor approximation and is comparable with the scheme based on nonlinear optimization and tangent-linear multiplicative inflation. However, it is less accurate than the nonlinear optimization and nonlinear inflation estimation ETKF scheme proposed in this study. On the other hand, both schemes have similar F-RMSE over F-Spread ratios.

Despite the advantage that the objective functions (Eqs. 24 and 27) are easier to minimize, the computational cost of the ETKF with the second-order Taylor approximation may increase from computing $(\mathbf{X}_i^f \mathbf{w})^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f, k} \mathbf{X}_i^f \mathbf{w}$. Because the most typical nonlinear observation operator in numerical weather prediction is the radiative transfer model RTTOV, the related computational issue is discussed and is documented in Appendix E. In fact, unlike forecast operators, the observation operators are usually localized, and therefore, the computation of $(\mathbf{X}_i^f \mathbf{w})^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f, k} \mathbf{X}_i^f \mathbf{w}$ is still feasible.

In addition, there are other ways to address this problem. For example, in the deterministic variational framework, Met Office re-linearizes the observation operator every 10 iterations (Rawlins et al., 2007), and ECMWF uses a nonlinear outer loop. Both approaches retain the efficiency of a tangent-linear approximation in the inner loop, while allowing for nonlinearity at a higher level. To better understand the efficacy of the ETKF scheme with second-order Taylor approximation, a more careful comparison with alternative assimilation schemes is necessary. We plan to face this challenge in the near future.

4.3 Caveats

This study assumes the inflation factor to be constant in space, but this is apparently not the case in many practical applications, specifically when observations are sparse. Applying the same inflation value to all state variables may overinflate the forecast errors of the state variables without observations (Hamill and Whitaker, 2005; Anderson,



2009; Miyoshi et al., 2010; Miyoshi and Kunii, 2012). If the forecast model has a large error, a multiplicative inflation may not be effective enough to improve the assimilation results. In this case, the additive inflation and localization technique may be applied to further improve the assimilation quality (Wu et al., 2013).

5 This study also assumes that the analysis increment can be expressed as a linear combination of ensemble forecast errors (Eq. 8). This assumption is true if the observation operator is tangent-linear, but the nonlinear observation operator can affect the combination of possible increments that produce the optimal analysis (Yang et al., 2012). However, our examples demonstrate that the proposed ETKF methods can still
10 work well when the observation operators are not tangent-linear.

At the last, but not the least, the results concluded in this study are related to the Lorenz-96 experiment. It may not be regarded as general rules. However, they can serve as counter examples to validate some ideas.

5 Conclusions

15 In this study, a new approach to inflating the ensemble forecast errors is proposed for the ETKF with a nonlinear observation operator. For an idealized model, it is shown that the proposed inflation approach can reduce analysis error compared with the tangent-linear multiplicative inflation, despite it being computationally more expensive. An ETKF scheme with the second-order Taylor approximation is also proposed. In terms of analysis error, the scheme is better than the first-order Taylor approximation ETKF scheme
20 and traditional ETKF scheme, specifically when the model error is larger. However, it is comparable to the scheme based on nonlinear optimization and tangent-linear multiplicative inflation. Finally, the proposed ETKF scheme with nonlinear optimization and nonlinear inflation was found to be the best among all of the schemes presented in this
25 study.

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the future studies, we plan to further investigate the computational efficiency of the proposed ETKF schemes and to validate them using more sophisticated dynamic models and observation systems.

Appendix A

5 Derivation of Eq. (6)

The estimation of the inflation factors λ is based on the innovation statistic normalized by the square root of the observation error covariance matrix

$$\begin{aligned} \mathbf{d}_i &= \mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^f)) \\ &= \mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) + \mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)), \end{aligned} \quad (\text{A1})$$

where \mathbf{y}_i^o , \mathbf{x}_i^f and \mathbf{x}_i^t are the observation, forecast and true state vector at the i th time step, respectively, and H_i is the observation operator. The covariance matrix of the random vector \mathbf{d}_i can be expressed as a second-order regression equation (Wang and Leblanc, 2008):

$$\begin{aligned} \mathbf{d}_i \mathbf{d}_i^T &= E \left[(\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) + \mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))) \right. \\ &\quad \left. (\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) + \mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)))^T \right] + \Xi, \end{aligned} \quad (\text{A2})$$

where E is the expectation operator and Ξ is a zero-mean error matrix. The expectation in Eq. (A2) has the decomposition

$$\begin{aligned} E \left[(\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) + \mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))) \right. \\ \left. (\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) + \mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)))^T \right] \end{aligned}$$

$$\begin{aligned}
&= E \left[\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t))^T \mathbf{R}_i^{-1/2} \right] \\
&\quad + E \left[\mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)) (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))^T \mathbf{R}_i^{-1/2} \right] \\
&\quad + E \left[\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))^T \mathbf{R}_i^{-1/2} \right] \\
&\quad + E \left[\mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)) (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t))^T \mathbf{R}_i^{-1/2} \right]. \tag{A3}
\end{aligned}$$

Assuming the forecast and observation errors are statistically independent, we have

$$\begin{aligned}
&E \left[\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))^T \mathbf{R}_i^{-1/2} \right] \\
&= \mathbf{R}_i^{-1/2} E \left[(\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))^T \right] \mathbf{R}_i^{-1/2} = \mathbf{0}, \tag{A4}
\end{aligned}$$

$$\begin{aligned}
&E \left[\mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)) (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t))^T \mathbf{R}_i^{-1/2} \right] \\
&= \mathbf{R}_i^{-1/2} E \left[(H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)) (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t))^T \right] \mathbf{R}_i^{-1/2} = \mathbf{0}. \tag{A5}
\end{aligned}$$

From Eq. (2), $\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)$ is the observation error at the i th time step, and hence,

$$\begin{aligned}
&E \left[\mathbf{R}_i^{-1/2} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t))^T \mathbf{R}_i^{-1/2} \right] \\
&= \mathbf{R}_i^{-1/2} E \left[(\mathbf{y}_i^o - H_i(\mathbf{x}_i^t)) (\mathbf{y}_i^o - H_i(\mathbf{x}_i^t))^T \right] \mathbf{R}_i^{-1/2} \\
&= \mathbf{R}_i^{-1/2} \mathbf{R}_i \mathbf{R}_i^{-1/2} \\
&= \mathbf{I}. \tag{A6}
\end{aligned}$$

In a perfect system, truth would be statistically indistinguishable from one of the ensemble forecast states, but in a real system this is not guaranteed. Hence, we use an inflation factor to adjust the ensemble forecast states $\mathbf{x}_{i,j}^f$ to $\mathbf{x}_i^f + \sqrt{\lambda}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)$, ($j = 1, \dots, m$).

**Improving ETKF
using the
second-order
information**

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Because the ensemble forecast states may be regarded as sample points of \mathbf{x}_i^t (Anderson, 2007), we have

$$\begin{aligned}
 & E \left[\mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f)) (H_i(\mathbf{x}_i^t) - H_i(\mathbf{x}_i^f))^T \mathbf{R}_i^{-1/2} \right] \\
 &= \frac{1}{m-1} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} (H_i(\mathbf{x}_i^f + \sqrt{\lambda}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)) - H_i(\mathbf{x}_i^f)) \right. \\
 & \quad \left. (H_i(\mathbf{x}_i^f + \sqrt{\lambda}(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)) - H_i(\mathbf{x}_i^f))^T \mathbf{R}_i^{-1/2} \right] \\
 &\equiv \mathbf{C}_i(\lambda).
 \end{aligned} \tag{A7}$$

Substituting Eqs. (A3)–(A7) into Eq. (A2), we have

$$\mathbf{d}_i \mathbf{d}_i^T = \mathbf{C}_i(\lambda) + \mathbf{I} + \Xi. \tag{A8}$$

It follows that the second-order moment statistic of error Ξ can be expressed as

$$\begin{aligned}
 \text{Tr} [\Xi \Xi^T] &= \text{Tr} \left[(\mathbf{d}_i \mathbf{d}_i^T - \mathbf{C}_i(\lambda) - \mathbf{I}) (\mathbf{d}_i \mathbf{d}_i^T - \mathbf{C}_i(\lambda) - \mathbf{I})^T \right] \\
 &\equiv L_i(\lambda).
 \end{aligned} \tag{A9}$$

Appendix B

Derivation of $J_{i|\mathbf{w}}$ and $\tilde{J}_{i|\mathbf{w}}$

The first-order derivative of the objective function $J_i(\mathbf{w})$ (Eq. 10) is

$$\mathbf{j}_i(\mathbf{w}) = (m-1)\mathbf{w} - \sqrt{\hat{\lambda}_i}(\mathbf{X}_i^f)^T \hat{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i}\mathbf{X}_i^f\mathbf{w}}^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i \left(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i}\mathbf{X}_i^f\mathbf{w} \right) \right], \tag{B1}$$

**Improving ETKF
using the
second-order
information**

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where

$$\dot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}} = \left(\begin{array}{ccc} \frac{\partial h_i(1)}{\partial x_i(1)} & \cdots & \frac{\partial h_i(1)}{\partial x_i(n)} \\ \vdots & \ddots & \vdots \\ \frac{\partial h_i(p_i)}{\partial x_i(1)} & \cdots & \frac{\partial h_i(p_i)}{\partial x_i(n)} \end{array} \right) \Bigg|_{\mathbf{x}_i = \mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}} \quad (\text{B2})$$

is the first-order derivative of H_i evaluated at $\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}$. Then, the second-order derivative of $J_i(\mathbf{w})$ is

$$\ddot{J}_i(\mathbf{w}) = (m-1)\mathbf{I} + \hat{\lambda}_i (\mathbf{X}_i^f)^T \dot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}}^T \mathbf{R}_i^{-1} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}} \mathbf{X}_i^f - \hat{\lambda}_i \mathbf{A}, \quad (\text{B3})$$

where \mathbf{A} is an $m \times m$ matrix with the (k, l) entry

$$\left((\mathbf{x}_{i,k}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f + \mathbf{X}_i^f \mathbf{w}} \otimes (\mathbf{x}_{i,l}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i \left(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right) \right]. \quad (\text{B4})$$

The notation “ \otimes ” denotes an outer product operator of the block matrix defined in Eq. (23). $\ddot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}}$ is the second-order derivative of H_i at $\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}$, that is,

$$\ddot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}} \equiv \begin{pmatrix} \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}, 1} \\ \vdots \\ \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}, p_i} \end{pmatrix},$$

$$\ddot{\mathbf{H}}_{i|\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}, k} \equiv \left(\begin{array}{ccc} \frac{\partial^2 h_i(k)}{\partial x_i(1) \partial x_i(1)} & \cdots & \frac{\partial^2 h_i(k)}{\partial x_i(1) \partial x_i(n)} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 h_i(k)}{\partial x_i(n) \partial x_i(1)} & \cdots & \frac{\partial^2 h_i(k)}{\partial x_i(n) \partial x_i(n)} \end{array} \right) \Bigg|_{\mathbf{x}_i = \mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}}, \quad k = 1, \dots, p_i. \quad (\text{B5})$$

Appendix C

Details of the first-order approximation method in Sect. 2.2.1

Suppose H_i can be approximated by its first-order Taylor expansion at \mathbf{x}_i^f ,

$$H_i \left(\mathbf{x}_i^f + \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \approx H_i(\mathbf{x}_i^f) + \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f). \quad (\text{C1})$$

The term $\mathbf{C}_i(\lambda)$ in Eq. (6) can be simplified to

$$\begin{aligned} \mathbf{C}_i(\lambda) &\equiv \frac{1}{m-1} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left(H_i \left(\mathbf{x}_i^f + \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) - H_i(\mathbf{x}_i^f) \right) \right. \\ &\quad \left. \left(H_i \left(\mathbf{x}_i^f + \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) - H_i(\mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right] \\ &= \frac{1}{m-1} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left(\dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \left(\dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right] \\ &= \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \frac{1}{m-1} \sum_{j=1}^m \left[(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \right] \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} \\ &= \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2}. \end{aligned}$$

It follows that the objective function $L_i(\lambda)$ of Eq. (5) can be simplified to

$$L_{1,i}(\lambda) = \text{Tr} \left[\left(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \mathbf{I} \right) \left(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \mathbf{I} \right)^T \right]. \quad (\text{C2})$$

Because $L_{1,i}(\lambda)$ is a quadratic function of λ with positive quadratic coefficients, the inflation factor can be easily expressed as

$$\hat{\lambda}_i = \frac{\text{Tr} \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} (\mathbf{d}_i \mathbf{d}_i^T - \mathbf{I})^T \right]}{\text{Tr} \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} \right]}. \quad (\text{C3})$$

5 Similarly,

$$H_i \left(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right) \approx H_i(\mathbf{x}_i^f) + \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w}. \quad (\text{C4})$$

Substituting Eq. (C3) into Eq. (8), we can simplify the objective function $J_i(\mathbf{w})$ to

$$J_{1,i}(\mathbf{w}) = \frac{1}{2} (m-1) \mathbf{w}^T \mathbf{w} + \frac{1}{2} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} \right]^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} \right]. \quad (\text{C5})$$

The first-order derivative of $J_{1,i}(\mathbf{w})$ is

$$J_{1,i}(\mathbf{w}) = (m-1) \mathbf{w} - \left(\sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \right)^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} \right] \\ = (m-1) \mathbf{w} - \sqrt{\hat{\lambda}_i} (\mathbf{X}_i^f)^T \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} \right]. \quad (\text{C6})$$

Setting Eq. (C6) to zero and solving it leads to

$$\mathbf{w}_i^a = ((m-1) \mathbf{I} + \hat{\lambda}_i \mathbf{X}_i^{fT} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f)^{-1} \sqrt{\hat{\lambda}_i} \mathbf{X}_i^{fT} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} (\mathbf{y}_i^o - H_i(\mathbf{x}_i^f)). \quad (\text{C7})$$

Lastly, the second-order derivative of $J_{1,i}(\mathbf{w})$ is

$$J_{1,i}(\mathbf{w}) = (m-1) \mathbf{I} + \hat{\lambda}_i \mathbf{X}_i^{fT} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f. \quad (\text{C8})$$

**Improving ETKF
using the
second-order
information**

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Appendix D

Details of the second-order approximation method in Sect. 2.2.2

Suppose H_i can be approximated by its second-order Taylor expansion at \mathbf{x}_i^f ,

$$H_i \left(\mathbf{x}_i^f + \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \approx H_i(\mathbf{x}_i^f) + \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) + \frac{1}{2} \lambda \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right), \quad (\text{D1})$$

The notation “ \otimes ” is defined as in Eq. (23). The term $\mathbf{C}_i(\lambda)$ in Eq. (7) can be simplified to

$$\begin{aligned} \mathbf{C}_i(\lambda) &\equiv \frac{1}{m-1} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left(H_i \left(\mathbf{x}_i^f + \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) - H_i(\mathbf{x}_i^f) \right) \right. \\ &\quad \left. \left(H_i \left(\mathbf{x}_i^f + \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) - H_i(\mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right] \\ &= \frac{1}{m-1} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left(\dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) + \frac{1}{2} \lambda \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \right) \right. \\ &\quad \left. \cdot \left(\dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\lambda} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) + \frac{1}{2} \lambda \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \right)^T \mathbf{R}_i^{-1/2} \right] \\ &= \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \frac{1}{m-1} \sum_{j=1}^m \left[(\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \right] \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} \\ &\quad + \frac{\lambda^{3/2}}{2(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right] \\ &\quad + \frac{\lambda^{3/2}}{2(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} \right] \end{aligned}$$

$$\begin{aligned}
& + \frac{\lambda^2}{4(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \right. \\
& \quad \left. \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right] \\
& = \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \lambda^{3/2} \mathbf{C}_{1,i} - \lambda^{3/2} \mathbf{C}_{1,i}^T - \lambda^2 \mathbf{C}_{2,i}. \tag{D2}
\end{aligned}$$

5 where

$$\mathbf{C}_{i,1} = \frac{1}{2(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right], \tag{D3}$$

and

$$\begin{aligned}
\mathbf{C}_{i,2} = \frac{1}{4(m-1)} \sum_{j=1}^m \left[\mathbf{R}_i^{-1/2} \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right) \right. \\
\left. \left((\mathbf{x}_{i,j}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,j}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1/2} \right] \tag{D4}
\end{aligned}$$

are $p_i \times p_i$ matrices, which are independent of λ .

It follows that the objective function $L_i(\lambda)$ of Eq. (5) can be expressed as

$$\begin{aligned}
L_{2,i}(\lambda) = \text{Tr} \left[\left(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \lambda^{3/2} \mathbf{C}_{1,i} - \lambda^{3/2} \mathbf{C}_{1,i}^T - \lambda^2 \mathbf{C}_{2,i} - \mathbf{I} \right) \right. \\
\left. \cdot \left(\mathbf{d}_i \mathbf{d}_i^T - \lambda \mathbf{R}_i^{-1/2} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \hat{\mathbf{P}}_i \dot{\mathbf{H}}_{i|\mathbf{x}_i^f}^T \mathbf{R}_i^{-1/2} - \lambda^{3/2} \mathbf{C}_{1,i} - \lambda^{3/2} \mathbf{C}_{1,i}^T - \lambda^2 \mathbf{C}_{2,i} - \mathbf{I} \right)^T \right], \tag{D5}
\end{aligned}$$

which is a fourth-order algebraic equation of the inflation factor λ . Similarly,

$$H_i(\mathbf{x}_i^f + \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w}) \approx H_i(\mathbf{x}_i^f) + \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} + \frac{1}{2} \left(\left(\sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes \left(\sqrt{\hat{\lambda}_i} \mathbf{X}_i^f \mathbf{w} \right) \right). \tag{D6}$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Substituting Eq. (D6) into Eq. (10), we can simplify the objective function $J_i(\mathbf{w})$ to

$$J_{2,i}(\mathbf{w}) = \frac{1}{2}(m-1)\mathbf{w}^T \mathbf{w} + \frac{1}{2} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} - \frac{\hat{\lambda}_i}{2} \left((\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) \right) \right]^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} - \frac{\hat{\lambda}_i}{2} \left((\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) \right) \right]. \quad (\text{D7})$$

The first-order derivative of $J_{2,i}(\mathbf{w})$ is

$$J_{2,i}(\mathbf{w}) = (m-1)\mathbf{w} - \left[\sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f + \hat{\lambda}_i \mathbf{B}_1 \right]^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} - \frac{\hat{\lambda}_i}{2} \left((\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) \right) \right], \quad (\text{D8})$$

where \mathbf{B}_1 is a $p_i \times m$ matrix with the (k, l) entry $\mathbf{X}_{i,l}^{f,T} \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f, k} \mathbf{X}_i^f \mathbf{w}$.

The second-order derivative of $J_{2,i}(\mathbf{w})$ is

$$\ddot{J}_{2,i}(\mathbf{w}) = (m-1)\mathbf{I} + \left[\sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f + \hat{\lambda}_i \mathbf{B}_1 \right]^T \mathbf{R}_i^{-1} \left[\sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f + \hat{\lambda}_i \mathbf{B}_1 \right] - \hat{\lambda}_i \mathbf{B}_2. \quad (\text{D9})$$

where \mathbf{B}_2 is an $m \times m$ matrix with the (k, l) entry

$$\left((\mathbf{x}_{i,k}^f - \mathbf{x}_i^f)^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{x}_{i,l}^f - \mathbf{x}_i^f) \right)^T \mathbf{R}_i^{-1} \left[\mathbf{y}_i^o - H_i(\mathbf{x}_i^f) - \sqrt{\hat{\lambda}_i} \dot{\mathbf{H}}_{i|\mathbf{x}_i^f} \mathbf{X}_i^f \mathbf{w} - \frac{\hat{\lambda}_i}{2} \left((\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) \right) \right].$$

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Computational feasibility

We take the radiative transfer model (RTTOV) as an example of observation operators in numerical weather prediction to discuss the computational feasibility of the ETKF with second-order approximation assimilation method. Generally speaking, the ensemble size m is from tens to hundreds, the dimension of observations (including gauge observations and AMSU brightness temperature) p_i is hundreds of thousands, and the dimension of state vector n is tens of millions. If the storage and the number of multiplications for computing any array are not in the dimension of $n \times n$, $n \times p_i$ or $p_i \times p_i$, the computation should be feasible.

In our proposed ETKF with second-order approximation, the most expensive part is in computing the array

$$(\mathbf{X}_i^f \mathbf{w})^T \otimes \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f} \otimes (\mathbf{X}_i^f \mathbf{w}) = \left\{ (\mathbf{X}_i^f \mathbf{w})^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f,1} \mathbf{X}_i^f \mathbf{w}, \dots, (\mathbf{X}_i^f \mathbf{w})^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f,p_i} \mathbf{X}_i^f \mathbf{w} \right\}. \quad (\text{E1})$$

Therefore, we only discuss the problems related to the computation of $(\mathbf{X}_i^f \mathbf{w})^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f,k} \mathbf{X}_i^f \mathbf{w}$.

E1 Storage problems

By the matrix multiplication rule,

$$(\mathbf{X}_i^f \mathbf{w})^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f,k} \mathbf{X}_i^f \mathbf{w} = \mathbf{w}^T \left((\mathbf{X}_i^f)^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f,k} \mathbf{X}_i^f \right) \mathbf{w}, \quad (\text{E2})$$

where the matrix in the middle of the right hand-side of Eq. (E2)

$$(\mathbf{X}_i^f)^T \ddot{\mathbf{H}}_{i|\mathbf{x}_i^f,k} \mathbf{X}_i^f \quad (\text{E3})$$

is of dimension $m \times m$, because subscript k runs from 1 to p_i , the size of the array in Eq. (E1) is $m \times m \times p_i$. Therefore, there is no storage problem to save this array.

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



E2 The computational cost of Eq. (E3)

Usually, $mn(m+n)$ times multiplication are required to compute a matrix such as the one in Eq. (E3). However, in the case of the RTTOV observation operator, $\ddot{\mathbf{H}}_{i|x_i^f, k}$ is a sparse matrix with a large number of zeros and the non-zero part has a simple regular structure. This is because an MSU brightness temperature measurement on a grid point (denoted by $y_i^o(k)$) is only related to the meteorological state variables on the transmission route. Suppose the meteorological model has 50 layers and 6 types of variables, the number of state variables on the transmission route of the MSU brightness temperature $y_i^o(k)$ is approximately $s = 300$. For the variables not on the transmission route, the corresponding entries in $\ddot{\mathbf{H}}_{i|x_i^f, k}$ (Eq. 22) are zero. Therefore, the computation of Eq. (E3) only requires $ms(m+s)/2$ times of multiplication, which is clearly feasible.

Acknowledgements. This work was supported by the National Program on Key Basic Research Project of China (Grant Nos. 2010CB951604 and 2010CB950703), the Open Fund of State Key Laboratory of Remote Sensing Science (Grant No. OFSLRSS201418), the Natural Sciences and Engineering Research Council of Canada (NSERC), and the Fundamental Research Funds for the Central Universities (No. 2012LYB39). We thank the editor and the anonymous referees for improving this paper.

References

- Anderson, J. L.: An adaptive covariance inflation error correction algorithm for ensemble filters, *Tellus A*, 59, 210–224, 2007.
- Anderson, J. L.: Spatially and temporally varying adaptive covariance inflation for ensemble filters, *Tellus A*, 61, 72–83, 2009.
- Anderson, J. L. and Anderson, S. L.: A Monte Carlo implementation of the non-linear filtering problem to produce ensemble assimilations and forecasts, *Mon. Weather Rev.*, 127, 2741–2758, 1999.

**Improving ETKF
using the
second-order
information**

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bishop, C. H. and Toth, Z.: Ensemble transformation and adaptive observations, *J. Atmos. Sci.*, 56, 1748–1765, 1999.
- Bishop, C. H., Etherton, J., and Majumdar, J.: Adaptive sampling with the ensemble transform Kalman filter. Part I: Theoretical aspects, *Mon. Weather Rev.*, 129, 420–436, 2001.
- 5 Burgers, G., van Leeuwen, P. J., and Evensen, G.: Analysis scheme in the ensemble Kalman Filter, *Mon. Weather Rev.*, 126, 1719–1724, 1998.
- Butcher, J. C.: *Numerical Methods for Ordinary Differential Equations*, ISBN 0471967580, John Wiley & Sons, England, 425 pp., 2003.
- Chen, Y., Oliver, D. S., and Zhang, D. X.: Data assimilation for nonlinear problems by ensemble
10 Kalman filter with reparameterization, *J. Petrol. Sci. Eng.*, 66, 1–14, 2009.
- Courtier, P., Thépaut, J. N., and Hollingsworth, A.: A strategy for operational implementation of 4D-Var, using an incremental approach, *Q. J. Roy. Meteor. Soc.*, 120, 1367–1387, 1994.
- Daescu, D. N. and Navon, I. M.: Efficiency of a POD-based reduced second-order adjoint model in 4D-Var data assimilation, *Int. J. Numer. Meth. Fl.*, 53, 985–1004, 2007.
- 15 Evensen, G.: Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics, *J. Geophys. Res.*, 99, 10143–10162, 1994a.
- Evensen, G.: Inverse methods and data assimilation in nonlinear ocean models, *Physica D*, 77, 108–129, 1994b.
- 20 Evensen, G.: Advanced data assimilation for strongly nonlinear dynamics, *Mon. Weather Rev.*, 125, 1342–1354, 1997.
- Flowerdew, J. and Bowler, N.: Improving the use of observations to calibrate ensemble spread, *Q. J. Roy. Meteor. Soc.*, 137, 467–482, 2011.
- Hamill, T. M. and Whitaker, J. S.: Accounting for the error due to unresolved scales in ensemble data assimilation: a comparison of different approaches, *Mon. Weather Rev.*, 133, 3132–
25 3147, 2005.
- Han, Y., van Delst, P., Liu, Q. H., Weng, F. Z., Yan, B. H., Treadon, R., and Derber, J.: Community Radiative Transfer Model (CRTM) – Version 1, NOAA NESDIS Technical Report, NOAA Technical Report NESDIS 122., Washington, D.C., 122 pp., 2006.
- 30 Hunt, B. R., Kostelich, J., and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: a local ensemble transform Kalman filter, *Physica D*, 230, 112–126, 2007.
- Ide, K., Courtier, P., Michael, G., and Lorenc, A. C.: Unified notation for data assimilation: operational, sequential and variational, *J. Meteorol. Soc. Jpn.*, 75, 181–189, 1997.

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Lawson, W. G. and Hansen, J. A.: Implications of stochastic and deterministic filters as ensemble-based data assimilation methods in varying regimes of error growth, *Mon. Weather Rev.*, 132, 1966–1981, 2004.

Le Dimet, F. X., Navon, I. M., and Daescu, D. N.: Second-order information in data assimilation, *Mon. Weather Rev.*, 130, 629–648, 2002.

Li, H., Kalnay, E., and Miyoshi, T.: Simultaneous estimation of covariance inflation and observation errors within an ensemble Kalman filter, *Q. J. Roy. Meteor. Soc.*, 135, 523–533, 2009.

Liang, X., Zheng, X. G., Zhang, S. P., Wu, G. C., Dai, Y. J., and Li, Y.: Maximum likelihood estimation of inflation factors on forecast error covariance matrix for ensemble Kalman filter assimilation, *Q. J. Roy. Meteor. Soc.*, 138, 263–273, 2012.

Liou, K. N.: *An Introduction to Atmospheric Radiation*, Academic Press, London, 583 pp., 2002.

Liu, C., Xiao, Q., and Wang, B.: An ensemble-based four-dimensional variational data assimilation scheme. Part 1: Technical formulation and preliminary test, *Mon. Weather Rev.*, 136, 3363–3373, 2008.

Lorenc, A. C.: The potential of the ensemble Kalman filter for NWP – a comparison with 4D-Var, *Q. J. Roy. Meteor. Soc.*, 129, 3183–3203, 2003.

Lorenz, E. N.: Predictability – a problem partly solved, in: *Proc. Seminar on Predictability*, ECMWF, Shinfield Park, Reading, Berkshire, UK, 40–58, 1996.

Lorenz, E. N. and Emanuel, K. A.: Optimal sites for supplementary weather observations: simulation with a small model, *J. Atmos. Sci.*, 55, 399–414, 1998.

McNally, A.: The direct assimilation of cloud-affected satellite infrared radiances in the ECMWF 4D-Var, *Q. J. Roy. Meteor. Soc.*, 135, 1214–1229, 2009.

Miller, R. N., Ghil, M., and Gauthiez, F.: Advanced data assimilation in strongly nonlinear dynamical systems, *J. Atmos. Sci.*, 51, 1037–1056, 1994.

Miyoshi, T.: The Gaussian approach to adaptive covariance inflation and its implementation with the local ensemble transform Kalman filter, *Mon. Weather Rev.*, 139, 1519–1535, 2011.

Miyoshi, T. and Kunii, M.: The Local Ensemble Transform Kalman Filter with the Weather Research and Forecasting Model: experiments with real observations, *Pure Appl. Geophys.*, 169, 321–333, 2012.

Miyoshi, T., Sato, Y., and Kadowaki, T.: Ensemble Kalman filter and 4D-Var inter-comparison with the Japanese operational global analysis and prediction system, *Mon. Weather Rev.*, 138, 2846–2866, 2010.

Improving ETKF using the second-order information

G. Wu et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Miyoshi, T., Kalnay, E., and Li, H.: Estimating and including observation error correlations in data assimilation, *Inverse Probl. Sci. En.*, 21, 387–398, 2013.
- Oke, P. R., Sakov, P., and Schulz, E.: A comparison of shelf observation platforms for assimilation in an eddy-resolving ocean model, *Dynam. Atmos. Oceans.*, 48, 121–142, 2009.
- 5 Palmer, T., Buizza, R., Hagedorn, R., Lawrence, A., Leutbecher, M., and Smith, L.: Ensemble prediction: a pedagogical perspective, *ECMWF Newsletter*, 106, 10–17, 2006.
- Parrish, D. F. and Deber, J. C.: The National meteorological center's spectral statistical-interpolation analysis system, *Mon. Weather Rev.*, 120, 1747–1763, 1992.
- Rawlins, F., Ballard, S. P., Bovis, K. J., Clayton, A. M., Li, D., Inverarity, G. W., Lorenc, A. C., and
 10 Payne, T. J.: The Met Office global four-dimensional variational data assimilation scheme, *Q. J. Roy. Meteor. Soc.*, 133, 347–362, 2007.
- Reichle, R. H.: Data assimilation methods in the earth sciences, *Adv. Water Resour.*, 31, 1411–1418, 2008.
- Sakov, P. and Oke, P. R.: A deterministic formulation of the ensemble Kalman filter: an alternative to ensemble square root filters, *Tellus A*, 60, 361–371, 2008.
- Saunders, R., Matricardi, M., and Brunel, P.: An improved fast radiative transfer model for assimilation of satellite radiance observations, *Q. J. Roy. Meteor. Soc.*, 125, 1407–1425, 1999.
- Stewart, L. M., Dance, S., and Nichols, N. K.: Data assimilation with correlated observation errors: experiments with a 1D shallow water model, *Tellus A*, 65, 19546, doi:10.3402/tellusa.v65i0.19546, 2013.
- 20 Wang, L. and Leblanc, A.: Second-order nonlinear least squares estimation, *Ann. I. Stat. Math.*, 60, 883–900, 2008.
- Wang, X. and Bishop, C. H.: A comparison of breeding and ensemble transform Kalman filter ensemble forecast schemes, *J. Atmos. Sci.*, 60, 1140–1158, 2003.
- 25 Wu, G. C., Zheng, X. G., Wang, L. Q., Zhang, S. P., Liang, X., and Li, Y.: A new structure of error covariance matrices and their adaptive estimation in EnKF assimilation, *Q. J. Roy. Meteor. Soc.*, 139, 795–804, 2013.
- Yang, S. C., Kalnay, E., and Hunt, B.: Handling nonlinearity in an Ensemble Kalman Filter: experiments with the Three-Variable Lorenz Model, *Mon. Weather Rev.*, 140, 2628–2646, 30 2012.

Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. The time-mean values of A-RMSE, F-RMSE and objective function (second-order distance of the squared innovation statistic to its expectation, Eq. 5) in the five ETKF schemes for Lorenz-96 model with forcing parameter $F = 12$ and parameter of observation operator $\alpha = 0.1$. ETKF: Traditional ETKF described in the last paragraph of Sect. 2.1; TT: tangent-linear approximation in both inflation (Eq. 17) and optimization (Eq. 20); TN: tangent-linear approximation in inflation (Eq. 17) and nonlinearity in optimization (Eq. 10); SS: second-order Taylor approximation in both inflation (Eq. 24) and optimization (Eq. 27); NN: nonlinearity in both inflation (Eq. 5) and optimization (Eq. 10).

Scheme	ETKF	TT	TN	SS	NN
A-RMSE	2.74	2.50	2.25	2.29	2.08
F-RMSE	3.20	3.00	2.77	2.66	2.52
L	49 700 074	17 078 480	8 768 825	9 177 962	8 458 902

Improving ETKF using the second-order information

G. Wu et al.

Table 2. The time-mean values of F-RMSE, F-Spread and the ratio of F-RMSE over F-Spread in the four ETKF schemes for Lorenz-96 model with forcing parameter $F = 12$ and parameter of observation operator $\alpha = 0.1$.

Scheme	ETKF	TT	TN	SS	NN
F-RMSE	3.20	3.00	2.77	2.66	2.52
F-Spread	1.06	1.45	1.46	1.48	1.45
F-RMSE/F-Spread	3.02	2.07	1.90	1.80	1.74

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improving ETKF using the second-order information

G. Wu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Similar to Table 2, but with $F = 8$.

Scheme	ETKF	TT	TN	SS	NN
F-RMSE	0.30	0.29	0.26	0.27	0.23
F-Spread	0.20	0.22	0.21	0.22	0.21
F-RMSE/F-Spread	1.50	1.32	1.24	1.18	1.09

Improving ETKF using the second-order information

G. Wu et al.




Fig. 1. Time-mean values of the A-RMSE as a function of forcing F for different assimilation methods on Lorenz-96 model. ETKF: traditional ETKF described in the last paragraph of Sect. 2.1; TT: tangent-linear approximation in both inflation (Eq. 17) and optimization (Eq. 20) (thin solid line); TN: tangent-linear approximation in inflation (Eq. 17) and nonlinearity in optimization (Eq. 10) (dashed line); SS: second-order Taylor approximation in both inflation (Eq. 24) and optimization (Eq. 27) (dotted line); NN: nonlinearity in both inflation (Eq. 5) and optimization (Eq. 10) (thick solid line). The ensemble size is 30.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improving ETKF using the second-order information

G. Wu et al.




Fig. 2. The proportions of residual ratios of the first-order and second-order Taylor expansions over the nonlinear observation operator $\mathbf{x}_{i,k}^t \exp\{0.1\mathbf{x}_{i,k}^t\}$ that fall outside the interval $[-0.1, 0.1]$, as a function of forcing F .

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
