



**Oscillations in a
simple
climate–vegetation
model**

J. Rombouts and M. Ghil

Oscillations in a simple climate–vegetation model

J. Rombouts¹ and M. Ghil^{2,3}

¹Centre for Complexity Science, University of Warwick, Coventry, UK

²Geosciences Department and Environmental Research & Teaching Institute,
Ecole Normale Supérieure, Paris, France

³Atmospheric & Oceanic Sciences Department and Institute of Geophysics & Planetary
Physics, University of California, Los Angeles, CA, USA

Received: 21 November 2014 – Accepted: 27 November 2014 – Published: 2 February 2015

Correspondence to: J. Rombouts (j.b.l.rombouts@warwick.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union & the
American Geophysical Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We formulate and analyze a simple dynamical systems model for climate–vegetation interaction. The planet we consider consists of a large ocean and a land surface on which vegetation can grow. The temperature affects vegetation growth on land and the amount of sea ice on the ocean. Conversely, vegetation and sea ice change the albedo of the planet, which in turn changes its energy balance and hence the temperature evolution. Our highly idealized, conceptual model is governed by two nonlinear, coupled ordinary differential equations, one for global temperature, the other for vegetation cover. The model exhibits either bistability between a vegetated and a desert state or oscillatory behavior. The oscillations arise through a Hopf bifurcation off the vegetated state, when the death rate of vegetation is low enough. These oscillations are anharmonic and exhibit a sawtooth shape that is characteristic of relaxation oscillations, as well as suggestive of the sharp deglaciations of the Quaternary.

Our model’s behavior can be compared, on the one hand, with the bistability of even simpler, Daisyworld-style climate–vegetation models. On the other hand, it can be integrated into the hierarchy of models trying to simulate and explain oscillatory behavior in the climate system. Rigorous mathematical results are obtained that link the nature of the feedbacks with the nature and the stability of the solutions. The relevance of model results to climate variability on various time scales is discussed.

1 Introduction

1.1 Background

Climate has an important effect on vegetation. Plant growth is affected by temperature, carbon dioxide (CO₂) levels and nutrient availability. The space available for growth matters as well: ice-covered parts of land are not suitable for it. The interaction also works the other way around, though: vegetation plays an important role in climate reg-

NPGD

2, 145–178, 2015

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to include vegetation in our models to obtain a better understanding of climate evolution and variability.

The simplest climate models are conceptual models, which are usually governed by a small number of ODEs. These models do not claim to be realistic – in the sense of making precise quantitative predictions, but allow one to study basic underlying mechanisms. They are also useful for exploring qualitative changes in the climate system’s behavior, commonly known in dynamical systems theory as bifurcations (Ghil and Childress, 1987; Dijkstra and Ghil, 2005), and related more recently to the broader and fuzzier concept of “tipping points” (Lenton et al., 2008). Bifurcations or tipping points correspond to situations in which a small parameter change can have a large effect on the behavior of the whole system.

Studying conceptual models can also provide guidance in interpreting results from larger, more detailed models (Brovkin et al., 1998, 2003; Ghil and Robertson, 2000). Early applications of dynamical systems theory and of numerical bifurcation studies to the climate system included H. Stommel’s bistability study of the thermohaline circulation (Stommel, 1961) and E. N. Lorenz’s study of chaos in a simple convection model (Lorenz, 1963). Another such application was to explain Quaternary glaciation cycles as climatic-oscillator solutions (Källén et al., 1979; Le Treut and Ghil, 1983; Saltzman, 1983; Ghil and Childress, 1987; Ghil, 1994; Crucifix, 2012).

In the latter idealized climate-oscillator models, vegetation is not usually included. There are, however, some simple models that explore the interaction between climate and vegetation, and we will briefly review some noteworthy ones in the next section.

1.2 Simple ODE models for climate–vegetation feedbacks

A pioneering model dealing with vegetation and climate was Daisyworld (Watson and Lovelock, 1983). This model illustrates how vegetation may act to regulate planetary temperature through the albedo feedback, for a wide range of parameter values. The varying parameter is in this case the incoming solar radiation. The model has been thoroughly studied and extended; see the review of Wood et al. (2008).

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Svirezhev and von Bloh (1996, 1997) introduced another set of simple, spatially zero-dimensional (0-D) models for vegetation-climate interactions. They include an ODE for temperature evolution, absent from Daisyworld, but look at only one type of vegetation, whereas Daisyworld has two. In their two-ODE model (Svirezhev and von Bloh, 1996), they find multiple steady states, a feature that regularly occurs across a hierarchy of climate–vegetation models, from the simplest (Dekker et al., 2007; Janssen et al., 2008; Aleina et al., 2013) to more complex (Brovkin et al., 1998; Claussen, 1998; Irizarry-Ortiz et al., 2003; Renssen et al., 2003) ones. In these models, vegetation and temperature are often coupled with precipitation, which provides an additional feedback mechanism.

Beyond the possibility of multiple steady states, it is of interest to examine in the simplest conceptual models the possibility of internal, self-sustained oscillations. Oscillatory behavior has been observed in Daisyworld-like models, for example when an explicit temperature equation is added (Nevison et al., 1999; Fernando Salazar and Poveda, 2009), or when delays are introduced (De Gregorio et al., 1992).

So far, though, simple, 0-D climate–vegetation models seem not to have included an ocean component. Earth's oceans constitute about 70 % of the area of the planet, and are a very important factor in determining its climate (Stommel, 1961; Saltzman, 1983; Ghil, 1994; Dijkstra and Ghil, 2005). The global ocean is often included in 0-D paleoclimate models for glacial cycles (Källèn et al., 1979; Le Treut and Ghil, 1983; Ghil and Childress, 1987; Gildor and Tziperman, 2001; Crucifix, 2012). Our model will include an ocean and the associated sea ice, and will combine certain features of Daisyworld-like models and climatic-oscillator models.

The purpose of this paper is to explore the solution space of such a model and to relate its behavior to other models and to observations. In the next section we formulate the model, and the numerical results are presented in Sect. 3. The model's behavior and its relation to more detailed models and to observations are discussed in Sect. 4.

2 Model description

The climate system contains several subsystems, all working together to produce highly nonlinear behavior through its many feedback mechanisms. Some of the simplest and most important feedback effects act via the planetary albedo: darker areas – like the vegetated ones – absorb more solar energy and thus warm the planet, while lighter areas – such as those covered by snow and ice – tend to cool it. The ice-albedo feedback has been included in energy balance models (EBMs) of climate long ago (Budyko, 1969; Sellers, 1969), and a vegetation-albedo feedback regulates Daisyworld’s behavior. The latter will constitute the main mechanism in our present model.

The model’s two governing equations are given below:

$$C_T \frac{dT}{dt} = (1 - \alpha(T, A))Q_0 - R_o(T), \quad (1a)$$

$$\frac{dA}{dt} = \beta(T)A(1 - A) - \gamma A. \quad (1b)$$

The variable T denotes global average temperature, while A denotes the fraction of land that is covered by vegetation.

Temperature T changes in Eq. (1a) as a result of the balance between incoming and outgoing energy. The parameter Q_0 is the incoming solar energy, which is equal here to 342.5 Wm^{-2} . The function $\alpha = \alpha(T, A)$ denotes the dependence of the albedo on T and A , and it is given by

$$\alpha = (1 - \rho)\alpha_o(T) + \rho(\alpha_v A + \alpha_g(1 - A)), \quad (2)$$

where ρ is the fraction of the planet that is land, taken to be 0.3 like on Earth.

The values α_o , α_v and α_g denote the albedo of the ocean, of the vegetation and of bare ground, respectively. The latter two are constant, and an essential ingredient of our model is that $\alpha_v < \alpha_g$, i.e. that forests and savannas are darker and absorb more energy than bare ground.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The albedo of the ocean will be taken as a function of global temperature T , to account for the possible presence of sea ice. We will use a ramp function, as in Sellers (1969) and Ghil (1976). This function is given by the following equation:

$$\alpha_o(T) = \begin{cases} \alpha_{\max} & \text{if } T \leq T_{\alpha,\ell}, \\ \alpha_{\max} + \frac{\alpha_{\min} - \alpha_{\max}}{T_{\alpha,u} - T_{\alpha,\ell}}(T - T_{\alpha,\ell}) & \text{if } T_{\alpha,\ell} < T \leq T_{\alpha,u}, \\ \alpha_{\min} & \text{if } T_{\alpha,u} < T. \end{cases} \quad (3)$$

5 Here $\alpha_{\max} = 0.85$ for the ice-covered ocean, $\alpha_{\min} = 0.25$ for the ice-free ocean, $T_{\alpha,\ell} = 263\text{K}$ and $T_{\alpha,u} = 300\text{K}$. The value of $T_{\alpha,u}$ is rather high – almost 27°C – which means that a tiny bit of sea ice will be present even for very high global temperature. Figure 1 shows a schematic representation of our planet's surface features and a plot of the ocean albedo versus temperature is shown in Fig. 2.

10 The function $R_o(T)$ denotes the energy outgoing from the planet. Several EBMs have used a modified version of the quartic Stefan-Boltzmann law (Sellers, 1969; Ghil, 1976) while other such models have used a linearization thereof (Budyko, 1969). We will also use here a linear dependence of outgoing radiation on temperature, but radiation increases in it more slowly with temperature than in the Stefan-Boltzmann case. This
 15 slower increase takes into account the fact that increasing temperature entrains increasing CO_2 levels and thus an increased greenhouse effect, which tends to decrease the outgoing radiation.

The form of $R_o(T)$ we choose is

$$R_o(T) = B_0 + B_1(T - T_{\text{opt}}), \quad (4)$$

20 where B_0 and B_1 are constants, while T_{opt} is the optimal growth temperature for the vegetation. There is a huge uncertainty in the values of these parameters, especially in B_1 , the linearized effect of temperature on radiative forcing (Stocker et al., 2013), since it all depends on which effects are taken into account and which are not (Zaliapin and Ghil, 2010).

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We will not attempt to estimate the most realistic values of these or other model parameters – since that is not the role of such simple, conceptual models (Ghil, 2001) – but for definiteness we will use $B_0 = 200$ and $B_1 = 2.5$. Their exact value does not play an important role in affecting the model’s behavior; still, we note that – over the range of temperatures we are interested in – they correspond rather well to a linearization of an outgoing energy term of the form $g\sigma T^4$ with $g = 0.55$, where g is a constant denoting the grayness of the atmosphere (see Fig. 3). Such a parametrization of outgoing energy – Stefan–Boltzmann law with a multiplicative factor – is often used, but usually g is allowed to depend on another variable, such as temperature (Sellers, 1969) or CO_2 level (Svirezhev and von Bloh, 1997), rather than being merely a constant, as in Fig. 3 here.

Vegetation cover A in Eq. (1b) grows logistically, with a temperature-dependent growth rate $\beta(T)$. The shape of the function β follows Daisyworld and related models, to wit,

$$\beta(T) = \max\left\{0, 1 - k(T - T_{\text{opt}})^2\right\}. \quad (5)$$

This means that the growth rate is zero, except for a certain interval, in which the dependence is parabolic with a maximum at T_{opt} , which is 283 K in our model.

The parameter γ is the mean death rate of the global vegetation A , and its value is, at first, set to 0.1 yr^{-1} . Table 1 lists the definition of the model parameters and their nominal values. In Sect. 3.2, γ will serve as a control parameter in our exploration of model behavior.

3 Analytical and numerical results

3.1 Multiple equilibria and their stability

We start, as usual, by looking at the fixed points of the system (Eq. 1), and will use, again as usual, interchangeably the terms steady states, stationary solutions and equilibria. One obvious fixed point is given by $A = A_0 = 0$, with temperature T_0 given by the

solution of $Q_0(1 - \alpha(T_0, 0)) = R_o(T_0)$. For our parameter values, a solution T_0 only exists when $T < T_{\alpha, \ell}$ and it equals $T_0 \simeq 243\text{K}$.

This result implies that a planet without vegetation will necessarily drop into a very cold “snowball-Earth” state. Note that this result differs from that of EBMs with no vegetation, in which two stable fixed points co-exist for sufficiently high Q_0 (North et al., 1981; Ghil and Childress, 1987).

A quick check shows that, for $p = 0$, i.e. for an “aqua-planet,” three fixed points are found for $A = 0$, as in classical EBMs: (i) a snowball Earth – called a “deep freeze” in the 1980s, before geological evidence for it was found; (ii) a high global temperature, comparable to Earth’s present climate; and (iii) an intermediate one, between the former two, and considerably closer to the present climate than to the subfreezing one. Hence it is the inclusion of the relatively high ground albedo for $p > 0$ that pushes our climate–vegetation model to a low temperature, if no vegetation is present.

In the presence of vegetation, $A \neq 0$, a fixed point (T_+, A_+) must satisfy

$$A_+ = 1 - \frac{\gamma}{\beta(T_+)}, \quad (6)$$

according to the vegetation equation (Eq. 1b). From the temperature equation (Eq. 1a), we see furthermore that, at a fixed point, one also has

$$A_+ = \frac{1}{\alpha_v - \alpha_g} \left[\frac{1}{p} \left(1 - \frac{R_o(T_+)}{Q_0} - (1 - p)\alpha_o(T_+) \right) - \alpha_g \right]. \quad (7)$$

The two curves in the phase plane (T, A) of the model that are given by Eqs. (6) and (7) are called the *null isoclines* – or *nullclines*, for short – of the system and they are plotted in Fig. 4. They intersect in two points, aside from the previously discussed bare-land solution; hence we have two additional equilibria for a vegetated planet.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To study the stability of these two fixed points, we consider the Jacobian matrix $\mathbf{J} = \mathbf{J}(T, A)$ for the system 1,

$$\mathbf{J} = \begin{pmatrix} a & \frac{-1}{C_T} Q_0 \frac{\partial \alpha}{\partial A} \\ \frac{\partial \beta}{\partial T} A(1 - A) & \beta(T)(1 - 2A) - \gamma \end{pmatrix}, \quad (8)$$

where

$$a = \frac{-1}{C_T} \left[Q_0 \frac{\partial \alpha}{\partial T} + \frac{\partial R_o}{\partial T} \right]. \quad (9)$$

For the no-vegetation fixed point $A = 0$, the eigenvalues of $\mathbf{J}(T, 0)$ are a and $\beta(T) - \gamma$. These eigenvalues are both negative, since T is so low at this fixed point that $\partial \alpha / \partial T = 0$ and $\beta(T) = 0$. Hence this non-vegetated state is always linearly stable.

Let us now look at the other two fixed points, shown in Fig. 4, where $A > 0$. Note that since $A = 1 - \gamma / \beta(T)$, the temperature must be in the range where $\beta(T) > 0$. With our choice of parameters, this range also lies in the interesting range between $T_{\alpha, \ell}$ and $T_{\alpha, u}$, within which we have sea ice feedback; it corresponds to the decreasing part of the temperature nullcline, given by Eq. (7), and shown as the dashed red line in Fig. 5.

The leftmost of these two fixed points, where $A > 0$, is a saddle. This can be seen from numerical computations, but we can also deduce it analytically. In the following, we will use β' instead of $\partial \beta / \partial T$, since the function β depends on temperature alone, and no confusion can arise. We have the following proposition:

Proposition 1. *A climate–vegetation fixed point (T_+, A_+) , where $A_+ > 0$, is a saddle, and hence unstable, if the vegetation growth rate $\beta(T)$ is an increasing function of global temperature T , $\beta'(T_+) > 0$.*

The proof is given in Appendix A, and it does not depend on any particular parameter values, only on their relative signs; nor is the exact parametrization of the growth rate β used.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The stable and unstable manifolds of the saddle were also computed numerically and they are plotted in Fig. 5. The stable manifold acts as a separatrix of the phase-plane flow, and it is almost vertical. This means that initial states with lower temperatures than that of the saddle will lead to the stable steady state ($T_0, A_0 = 0$) that is characterized by freezing temperatures and no vegetation, while initial states with higher temperatures will lead to the steady state with at least some vegetation present.

For the stability of the latter, rightmost fixed point, Proposition 1 above says nothing, since $\beta'(T) < 0$ there. Numerical computation shows that the eigenvalues of the Jacobian $\mathbf{J}(T, A)$ at this point are complex conjugate and equal to $\lambda_{\pm} = -0.006 \pm 0.028i$. Hence the fixed point is a focus and the real part $\lambda_r = \text{tr}(\mathbf{J})/2$, where $\text{tr}(\mathbf{J})$ is the trace of the Jacobian matrix, is negative; thus the focus is stable.

We can compute the trace of \mathbf{J} analytically:

$$\text{tr}(\mathbf{J}) = a + \beta(T)(1 - 2A) - \gamma = a - \beta(T) + \gamma = a + x. \quad (10)$$

The term a on the right-hand side is larger than zero, but the term $x = \gamma - \beta(T)$ compensates and, for the standard parameter values in Table 1, $\text{tr}(\mathbf{J})$ is negative, implying the stability of the focus. One suspects, however, that, as some of these parameter values change, the focus may lose its stability.

We investigate next this loss of stability and the more complex model behavior to which it leads.

3.2 Hopf bifurcation and oscillatory behavior

3.2.1 Analytical results

We study now in greater detail the stability of the fixed point with non-zero vegetation (T_+, A_+). The result is stated in the following proposition and the proof thereof is given in Appendix A, after the one for Proposition 1.

Proposition 2. A model fixed point (T_+, A_+) that lies in the region with active sea ice feedback ($T_{\alpha,\ell} < T_+ < T_{\alpha,u}$) is a focus if and only if the following inequality holds:

$$(1 - y)^2 < 4\eta(y)y, \quad (11)$$

where $y = (y - \beta(T_+))/a$, $\eta(y)$ is (up to a sign change) the ratio of the derivatives of the two nullclines and a is given by Eq. (9).

Given the inequality (Eq. 11), the focus is stable if $y < -1$.

The proof in the Appendix relies simply on rewriting the classic conditions on the Jacobian.

We know that $y < 0$, since $x < 0$ and a is positive, and also that $\eta(y)$ is always negative. Since for all negative y we have

$$(1 - y)^2 > 4y,$$

a necessary condition to have a focus is that $|\eta| > 1$, i.e. that the ratio of the derivatives of the nullclines should be large enough in absolute value.

The proof of Proposition 1 in the Appendix implies, however, that this condition is already necessary for a fixed point to *not* be a saddle. Since the temperature nullcline is a piecewise linear curve, its derivative is a constant and a focus can only occur when the vegetation nullcline is steep enough. This condition, in turn, corresponds to the vegetation cover A_+ being small enough, as can be seen in the phase plane plotted in Fig. 5. It follows that a Hopf bifurcation and, therefore, oscillations can only occur around a model state with low vegetation. Since we did not use the actual value of any of the parameters, this result is quite general.

As shown in the caption of Fig. 5, the temperature value for the vegetated fixed point is $T_+ = 297.9$. The corresponding value of y is -1.96 . The ratio $\eta(y)$ of the nullcline slopes equals -9.6 , hence $|\eta(y)| > 1$ and thus the point could be a focus. Substituting in Eq. (11) yields $8.75 < 75.17$, which shows that the inequality is satisfied and that this point is a focus. Since $y = -1.96 < -1$, the focus is stable.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



3.2.2 Numerical results

As announced at the end of Sect. 2, we now vary the the death rate γ of the vegetation and follow the change in model solutions. For decreasing γ , i.e. for longer-lived vegetation, the focus loses its stability and the associated Hopf bifurcation gives rise to a limit cycle, as shown in the bifurcation diagram of Fig. 6. This bifurcation occurs for $\gamma = 0.02572$, i.e. when the overturning time $1/\gamma$ of the vegetation is about 40 years, which corresponds definitely to trees rather than grasses.

We can also see in Fig. 6 that, when γ is higher than 0.41, non-zero vegetation can no longer be a steady-state solution. At $\gamma \simeq 0.41$, a saddle-node bifurcation gives rise to a stable (blue curve) and an unstable (red curve) branch of steady states, in addition to the stable branch $A \equiv 0$ studied in the previous subsection and not shown in the figure.

We mentioned in Sect. 3.1 above that the parameter change needs to make $\text{tr}(\mathbf{J})$ positive in order to change the stability. This also works if, instead of decreasing γ , we decrease the thermal heat capacity C_T . It is the product γC_T that determines the behavior of the system, not the two parameters separately.

This product is a ratio of time scales, with C_T a typical time scale for temperature adjustments, while $1/\gamma$ is a typical time scale for vegetation. The oscillations occur, therefore, when the ratio of these two time scales satisfies a certain inequality. Such a condition was found to hold for other Hopf bifurcations, where the time scales need to match. For instance, in the temperature–ice sheet oscillator of Källèn et al. (1979), the characteristic time of ice sheet evolution C_L should be large enough in comparison with the time scale C_T associated with the thermal inertia of the oceans; see Ghil and Tavantzis (1983).

Nevison et al. (1999) also obtained certain results that are somewhat similar to ours. Their paper explores a Daisyworld with an equation for temperature evolution. The system exhibits self-sustained oscillations for a certain range of C_T and γ , just like in our model. The albedo feedback in this model is due to two types of daisies, black

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(absorbing) and white (reflecting) ones. The bimodality of the albedos in both models leads us to suspect that the role of sea ice in our model corresponds roughly to that of the white daisies in the model by Nevison and colleagues.

This analogy shouldn't be pushed too far, however. White daisy growth is favored by warmer temperatures, whereas sea ice appears at low temperatures. A comparison between our Fig. 7 and Fig. 2 in Nevison et al. (1999) shows that the behavior of black daisies and temperature is analogous to our vegetation and temperature, while increase in sea ice and white daisies have a different phase. The former occurs with decreasing temperature, while the latter happens as temperature increases.

Another interesting difference between the two models lies in the value of γ for which oscillations are obtained. We have $1/\gamma \simeq 40$ yr, while Nevison and colleagues obtain the oscillations when $1/\gamma$ is of the order of a few months, with a similar value of C_T . The difference probably resides in the nature of the feedback: in our model, the ocean and sea ice play a role, with both operating on seasonal and interannual time scales. In the Nevison et al. (1999) model, it is two kinds of daisies on an all-land planet that interact, both with much shorter time scales.

Oscillatory behavior is also observed in the Daisyworld variant studied by De Gregorio et al. (1992). In their paper, an explicit delay is introduced into the Daisyworld system. When the delay is large enough, oscillations are observed. This again shows the importance of time scales in these oscillations. The mechanism by which the oscillations arise in the delayed system of De Gregorio and colleagues is also shown to be a Hopf bifurcation, like in delayed, spatially dependent EBMs (Bhattacharya et al., 1982; Roques et al., 2014) and in certain simple and intermediate models of the El Niño–Southern Oscillation (ENSO) (Jin et al., 1994; Tziperman et al., 1994; Ghil et al., 2008, and references therein).

This again shows the importance of time scales: only when the effect of temperature on the growth rate of the vegetation is substantially delayed, the system exhibits internal oscillations. The discussion on time scales in Daisyworld is the subject of a recent

paper by Weaver and Dyke (2012), where the persistency of regulatory behavior under a separation of timescales is examined.

Figure 7 (upper panel) shows plots of temperature (red curve) and vegetation cover (blue curve) versus time. The oscillations have an amplitude of a few degrees for the temperature, but are very large for the vegetation, going from almost no vegetation to 70 % of the land being covered by plants. In addition, the vegetation plot shows a sawtooth-like shape, with the vegetation cover almost vanishing for long times, after which it shoots back up.

Such sawtooth-like behavior characterizes the ice volume evolution during Quaternary glaciations; see, for instance, Ghil and Childress (1987, Ch. 11 and references therein). This behavior has been modeled using relaxation oscillations, cf. Le Treut and Ghil (1983), Ghil (1994) and Crucifix (2012), among others. In these studies, the sawtooth shape is due to a quick melting and slow build-up of the ice sheets, whereas in Gildor and Tziperman (2001), the sea ice plays a key role. In our model, it is the surge in vegetation growth accompanying a temperature rise that provides the characteristically asymmetric nature of the oscillation.

Given the importance of sea ice feedback in our model, sea ice extent is plotted in Fig. 7b. In the temperature interval in which the sea ice feedback is active, the sea ice extent depends linearly on T , so the shape of the curve in this panel is similar to that of temperature in the upper panel, with the directions of waxing and waning reversed. There is a slow expansion of sea ice and rapid melting, which resembles the behavior of land ice volume during glacial-interglacial cycles. This highly nonlinear oscillation in our model indicates that it might be of interest to include vegetation in “ice age oscillators,” possibly combining it with a simple carbon cycle. As far as we know, this has not been done for ODE models. Certain results of more detailed models, however, do show that vegetation plays an important role in glacial cycles (Meissner et al., 2003; Horton et al., 2010).

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

4 Conclusions

4.1 Discussion

While the sea ice albedo feedback plays a crucial role in our model's oscillatory behavior, the amplitude of the oscillations in sea ice extent is quite small, from 0 to a mere 6%. Gildor and Tziperman (2001) first proposed sea ice as determining factor in the inception of glacial cycles, but our model's oscillations have a much shorter period than the dominant 100 000 year cycle of the latter. Note that our climate–vegetation model differs from that of Gildor and Tziperman (2001) by the absence of the much slower and more massive continental ice sheets, and it differs from the model of Svirezhev and von Bloh (1996) by including the ocean and the sea ice, as well as by the less crucial parametrization of the outgoing radiation. The latter authors actually prove the absence of limit cycles in their climate–vegetation model by using the Bendixson-Dulac criterion (e.g. Andronov et al., 1966). This proof reinforces our argument that sea ice plays an import role in intrinsic climate variability on long time scales.

Because of Proposition 1 and the shape of the growth curve, it is essential that the fixed point for which the Hopf bifurcation occurs have a temperature higher than T_{opt} , since $\beta'(T)$ needs to be negative, and lower than $T_{\alpha,u}$, so that the sea ice feedback be on. The former condition implies that the optimal growth temperature for plants needs to be relatively low, while the latter condition requires that sea ice be present in polar regions for relatively high global temperatures.

Such distinctions between local and global variables are hard to make in highly idealized, ODE models, and it is clear that there is a need for pursuing this type of analysis in more detailed, spatially dependent models, either with several climate zones or with continuous dependence on latitude. Such an extension would allow the model, for example, to have vegetated taiga and tundra interact with sea ice in higher latitudes and desert vegetation interact with atmospheric and oceanic temperatures in lower latitudes.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spatially dependent versions of Daisyworld have already been studied and yielded interesting results. Adams et al. (2003) observed patten formation in a spatially one-dimensional Daisyworld. In their model, vegetation in different latitude belts interacts with global temperature. Patterns are also seen by Biton and Gildor (2012), whose one-dimensional Daisyworld is forced by seasonal variations in solar radiation. Both of these studies suggest that pursuing our model analysis in higher dimension, while still including ocean and sea ice, would be of interest.

Earth system models of intermediate complexity provide some evidence that vegetation plays a role in Quaternary climate variations. In particular, the albedo changes that accompany shifts in vegetation act to amplify orbital forcing (Claussen, 2009). In addition, more detailed models, such as those by Zeng et al. (1999) and Zeng and Neelin (2000) conclude that vegetation in the Sahel region interacts strongly with global temperatures on shorter time scales.

The concept of a hierarchy of models – already fairly well accepted in climate modeling (Schneider and Dickinson, 1974; Ghil and Robertson, 2000; Ghil, 2001) – suggests precisely this kind of further exploration of more detailed models. The present paper clearly shows that intrinsic variability is possible in a climate–vegetation system, at least in its simplest possible form – either as bistability or as self-sustained periodic oscillations – and that the combined effect of vegetation-albedo and of ice-albedo feedbacks can give rise to an oscillation-producing mechanism.

The conditions $\beta'(T) < 0$ and $\partial\alpha/\partial T < 0$ on these two feedbacks are necessary in our simple model for the vegetated fixed point to become unstable and lead to intrinsic variability. We note that limit cycles are also observed when changing for example the parametrization of outgoing radiation into a modified Stefan-Boltzmann law, or if we change the parameters $T_{\alpha,u}$ and T_{opt} . The important thing is that the conditions above on the derivatives be satisfied.

In our model, the feedback that produces oscillations is negative: increasing temperature lowers vegetation cover, which in turn lowers energy absorption and decreases temperature. Due to the large number of interacting processes in nature, the effective

sign of the feedback is not known directly from observations, and it might be different in different areas (Claussen, 2009; Meir et al., 2006). One can think of different models, too, in which another component or physical process than sea ice and the associated feedback provide the requisite oscillatory mechanism.

5 One such process is the hydrological cycle. Aleina et al. (2013) and Brovkin et al. (1998), for instance, have included it already in conceptual climate–vegetation models but observed no oscillations. A study by Fernando Salazar and Poveda (2009) did observe oscillations in a model where Daisyworld is extended with clouds and a basic hydrological cycle. The authors concluded that interaction between clouds and vegeta-
10 tion can improve the circumstances for life on the planet. Clouds are known to play an important role in changing the radiation balance of the planet, but their exact influence is poorly known (IPCC, 2007).

In addition to the influence of vegetation on the hydrological cycle, there is a whole range of other feedback effects between vegetation and several components of the
15 climate system. Of particular importance here is the biogeochemical feedback, as opposed to the biogeophysical effects, such as albedo change, that we have explored. Biogeochemical effects include the influence of vegetation on the carbon cycle, and therefore on the greenhouse effect and the radiation balance of the Earth.

A more detailed model than the one studied here could include a simplified version
20 of these effects. Lovelock and Kump (1994), for instance, have examined the regulatory effect of vegetation, both on land and in the ocean, under different conditions. They conclude that vegetation is involved in regulation for low temperatures, but the regulatory system collapses at high temperatures.

Another interesting model extension that comes to mind is the inclusion of oceanic
25 vegetation, i.e. phytoplankton. Plankton interacts with climate in different ways, through its possible effect on cloud formation (Ayers and Cainey, 2007), but also through its albedo.

**Oscillations in a
simple
climate–vegetation
model**

J. Rombouts and M. Ghil

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



4.2 Summary

We described a simple dynamical systems model for climate–vegetation interaction. The model planet has a large ocean, which can be covered by sea ice, and a land area, which can be covered by vegetation. The system variables are temperature and vegetation cover, and the coupling between those is given by the growth rate of the latter and by the albedo, both of which are temperature-dependent.

The model is similar to Daisyworld and related models, the main difference being the inclusion of an ocean. Our model is also related to EBMs and, in this respect, the novelty lies in the inclusion of vegetation as a main variable.

The model exhibits two stable states, one with and one without vegetation. The vegetated state can lose its stability through a Hopf bifurcation and give rise to a limit cycle. This happens when the typical time scale for vegetation overturn becomes high enough. The influence of the model's sea ice in reinforcing the albedo feedback is essential to the presence of oscillatory behavior. The oscillations observed are anharmonic and have a sawtooth shape, reminiscent of deglaciations during the Quaternary. We have obtained analytical results on the role of the feedbacks in determining the system's behavior and compared the resulting oscillations with other models.

Although some parameter values in our highly idealized model are not entirely realistic, the results add to the evidence that vegetation, in combination with other feedback effects, can play an important role in affecting climate. The model studied herein is also interesting because it is one of the simplest ODE models for climate–vegetation interactions that exhibits oscillatory behavior. This intrinsic variability is a basic manifestation of how vegetation affects climate, and constitutes an example of how complex behavior arises in the Earth system even at the lowermost levels of the modeling hierarchy. Once demonstrated and understood at such a level, one can ask next whether similar behavior does persist in more detailed models, and whether it does reflect the actual behavior of the natural climate–vegetation system.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Appendix: Proof of Propositions 1 and 2

In this Appendix, we prove the propositions from the main text. Recall the definition

$$a = -C_T^{-1}(Q_0 \partial \alpha / \partial T + \partial R_o / \partial T), \quad (\text{A1})$$

and let $b = -C_T^{-1} Q_0 \partial \alpha / \partial A$. From now on, we focus on the temperature interval where $\beta(T)$ is positive and, therefore, where the sea ice feedback is active. For these values of T , both a and b are positive constants, because of the linear dependence of the albedo on T and A and of the outgoing energy R_o on T .

For a fixed point (T_+, A_+) , the Jacobian in Eq. (8) can be rewritten as

$$\mathbf{J} = \begin{pmatrix} a & b \\ -\frac{\beta'(T)}{\beta^2(T)} \gamma x & x \end{pmatrix}, \quad (\text{A2})$$

where $x = \gamma - \beta(T)$. The determinant of the Jacobian is then

$$\det(\mathbf{J}) = ax + b \frac{\beta'(T)}{\beta^2(T)} \gamma x = x \left(a + b \frac{\beta'(T)}{\beta^2(T)} \gamma \right).$$

Since the fixed point (T_+, A_+) lies on the A -nullcline, we know that $A_+ = 1 - \gamma / \beta(T_+)$. Because $0 < A < 1$, we deduce that $x < 0$. and we have

$$\det(\mathbf{J}) < 0 \Leftrightarrow -\frac{a}{b} < \frac{\beta'(T) \gamma}{\beta(T)^2}. \quad (\text{A3})$$

This proves Proposition 1, since, when $\beta'(T) > 0$, the above inequality is always satisfied. Therefore $\det(\mathbf{J}) < 0$, which implies the fixed point is a saddle and thus unstable.

We will now use the same notation to prove Proposition 2, which we repeat here for the reader's convenience.

Proposition 2. A model fixed point (T_+, A_+) that lies in the region with active sea ice feedback ($T_{\alpha,\ell} < T_+ < T_{\alpha,u}$) is a focus if and only if the following inequality holds:

$$(1 - y)^2 < 4\eta(y)y, \quad (\text{A4})$$

where $y = (\gamma - \beta(T_+))/a$, $\eta(y)$ is (up to a sign change) the ratio of the derivatives of the two nullclines and a is given by Eq. (A1).

Given the inequality (Eq. A4), The focus is stable if $y < -1$.

Proof. In order to have a focus, the Jacobian matrix \mathbf{J} has to have eigenvalues with non-zero imaginary part. This translates into the condition

$$\text{tr}(\mathbf{J})^2 - 4\det(\mathbf{J}) < 0. \quad (\text{A5})$$

We substitute the values into \mathbf{J} above, while using our new constants a and b and the variable x , and get the inequalities

$$(a+x)^2 - 4 \left(ax - b \frac{\beta'}{\beta^2} \gamma x \right) < 0 \Leftrightarrow (a-x)^2 < 4 \frac{\beta'}{\beta^2} \gamma b x \Leftrightarrow (1-y)^2 < 4\eta(y)y, \quad (\text{A6})$$

where $y = x/a$ and $\eta(y) = \beta' b \gamma / (\beta^2 a)$. To show that the function η actually is, up to a change of sign, the ratio of the slopes of the nullclines, is a matter of computation: a quick calculation shows that the slope of the T -isocline is equal to $-a/b$, cf. Eq. (7), and that of the A -isocline is $\gamma\beta'(T)/\beta(T)^2$, cf. Eq. (6).

The statement about the stability of the focus follows from the fact that a sufficient condition for the focus to be stable is that $\text{tr}(\mathbf{J}) < 0$. This translates into $a + x < 0$, or $y < -1$. \square

Acknowledgements. It is a pleasure to thank K. Pakdaman for useful discussions on the presentation of this work. J. Rombouts was supported through an Erasmus Mundus scholarship of the European Union and M. Ghil benefited from partial support of the European Union's ENSEMBLES project. The paper is the result of research carried out during J. Rombouts's stay at the Ecole Normale Supérieure and the Ecole Polytechnique in Paris, as part of an Erasmus Mundus programme.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Adams, B., Carr, J., Lenton, T. M., and White, A.: One-dimensional daisyworld: spatial interactions and pattern formation, *J. Theor. Biol.*, 223, 505–513, doi:10.1016/S0022-5193(03)00139-5, 2003. 161
- 5 Aleina, F. C., Baudena, M., D'Andrea, F., and Provenzale, A.: Multiple equilibria on planet Dune: climate–vegetation dynamics on a sandy planet, *Tellus B*, 65, 17662, doi:10.3402/tellusb.v65i0.17662, 2013. 149, 162
- Andronov, A., Vitt, A., and Khaikin, A.: *Theory of Oscillators*, Pergamon, Oxford, 1966. 160
- Ayers, G. P. and Caine, J. M.: The CLAW hypothesis: a review of the major developments, *Environ. Chem.*, 4, 366–374, 2007. 147, 162
- 10 Bhattacharya, K., Ghil, M., and Vulis, I. L.: Internal variability of an energy-balance model with delayed albedo effects, *J. Atmos. Sci.*, 39, 1747–1773, 1982. 158
- Biton, E. and Gildor, H.: The seasonal effect in one-dimensional Daisyworld, *J. Theor. Biol.*, 314, 145–156, doi:10.1016/j.jtbi.2012.08.043, 2012. 161
- 15 Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests, *Science*, 320, 1444–1449, doi:10.1126/science.1155121, 2008. 147
- Brovkin, V., Claussen, M., Petoukhov, V., and Ganopolski, A.: On the stability of the atmosphere–vegetation system in the Sahara/Sahel region, *J. Geophys. Res.-Atmos.*, 103, 31613–31624, doi:10.1029/1998JD200006, 1998. 148, 149, 162
- 20 Brovkin, V., Levis, S., Loutre, M.-F., Crucifix, M., Claussen, M., Ganopolski, A., Kubatzki, C., and Petoukhov, V.: Stability analysis of the climate–vegetation system in the northern high latitudes, *Climatic Change*, 57, 119–138, doi:10.1023/A:1022168609525, 2003. 147, 148
- Budyko, M. I.: The effect of solar radiation variations on the climate of the Earth, *Tellus*, 21, 611–619, 1969. 150, 151
- 25 Charlson, R., Lovelock, J., Andreae, M., and Warren, S.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655–661, doi:10.1038/326655A0, 1987. 147
- Charney, J. G.: Dynamics of deserts and drought in the Sahel, *Q. J. Roy. Meteorol. Soc.*, 101, 193–202, doi:10.1002/qj.49710142802, 1975. 147
- 30 Charney, J. G., Stone, P. H., and Quirk, W. J.: Drought in the Sahara: a biogeophysical feedback mechanism, *Science*, 187, 434–435, doi:10.1126/science.187.4175.434, 1975. 147

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Claussen, M.: On multiple solutions of the atmosphere–vegetation system in present-day climate, *Global Change Biol.*, 4, 549–559, doi:10.1046/j.1365-2486.1998.t01-1-00122.x, 1998. 149
- Claussen, M.: Late Quaternary vegetation-climate feedbacks, *Clim. Past*, 5, 203–216, doi:10.5194/cp-5-203-2009, 2009. 147, 161, 162
- Claussen, M., Kubatzki, C., Brovkin, V., Ganopolski, A., Hoelzmann, P., and Pachur, H.-J.: Simulation of an abrupt change in Saharan vegetation in the Mid-Holocene, *Geophys. Res. Lett.*, 26, 2037–2040, doi:10.1029/1999GL900494, 1999. 147
- Crucifix, M.: Oscillators and relaxation phenomena in Pleistocene climate theory, *Philos. T. Roy. Soc. A*, 370, 1140–1165, doi:10.1098/rsta.2011.0315, 2012. 148, 149, 159
- De Gregorio, S., Pielke, R. A., and Dalu, G. A.: A delayed biophysical system for the Earth's climate, *J. Nonlin. Sci.*, 2, 293–318, doi:10.1007/BF01208927, 1992. 149, 158
- Dekker, S. C., Rietkerk, M., and Bierkens, M. F. P.: Coupling microscale vegetation-soil water and macroscale vegetation-precipitation feedbacks in semiarid ecosystems, *Global Change Biol.*, 13, 671–678, doi:10.1111/j.1365-2486.2007.01327.x, 2007. 149
- Dijkstra, H. A. and Ghil, M.: Low-frequency variability of the large-scale ocean circulation: a dynamical systems approach, *Rev. Geophys.*, 43, RG3002, doi:10.1029/2002RG000122, 2005. 148, 149
- Ermentrout, B.: *Simulating, Analyzing, and Animating Dynamical Systems: a Guide to XPPAUT for Researchers and Students*, SIAM, Philadelphia, USA, available at: <http://www.math.pitt.edu/~bard/xpp/xpp.html> (last access: 20 August 2014), 2002. 177
- Fernando Salazar, J. and Poveda, G.: Role of a simplified hydrological cycle and clouds in regulating the climate-biota system of Daisyworld, *Tellus B*, 61, 483–497, doi:10.1111/j.1600-0889.2008.00411.x, 2009. 149, 162
- Ghil, M.: Climate stability for a Sellers-type model, *J. Atmos. Sci.*, 33, 3–20, 1976. 151
- Ghil, M.: Cryothermodynamics: the chaotic dynamics of paleoclimate, *Physica D*, 77, 130–159, doi:10.1016/0167-2789(94)90131-7, 1994. 148, 149, 159
- Ghil, M.: Hilbert problems for the geosciences in the 21st century, *Nonlin. Processes Geophys.*, 8, 211–211, doi:10.5194/npg-8-211-2001, 2001. 147, 152, 161
- Ghil, M. and Childress, S.: *Topics in Geophysical Fluid Dynamics: Atmospheric Dynamics, Dynamo Theory, and Climate Dynamics*, Applied Mathematical Sciences, vol. 60, Springer-Verlag, New York, 1987. 148, 149, 153, 159

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ghil, M. and Robertson, A. W.: Solving problems with GCMs: general circulation models and their role in the climate modeling hierarchy, in: *General Circulation Model Development: Past, Present and Future*, edited by: Randall, D., Academic Press, San Diego, 285–325, 2000. 147, 148, 161
- 5 Ghil, M. and Tavantzis, J.: Global Hopf bifurcation in a simple climate model, *SIAM J. Appl. Math.*, 43, 1019–1041, doi:10.1137/0143067, 1983. 157
- Ghil, M., Zaliapin, I., and Thompson, S.: A delay differential model of ENSO variability: parametric instability and the distribution of extremes, *Nonlin. Processes Geophys.*, 15, 417–433, doi:10.5194/npg-15-417-2008, 2008. 158
- 10 Gildor, H. and Tziperman, E.: A sea ice climate switch mechanism for the 100-kyr glacial cycles, *J. Geophys. Res.-Oceans*, 106, 9117–9133, doi:10.1029/1999JC000120, 2001. 149, 159, 160
- Horton, D. E., Poulsen, C. J., and Pollard, D.: Influence of high-latitude vegetation feedbacks on late Palaeozoic glacial cycles, *Nat. Geosci.*, 3, 572–577, doi:10.1038/ngeo922, 2010. 159
- 15 IPCC: *Climate Change 2007: The Physical Science Basis*, in: *Contribution of Working Group I to the Fourth Assessment Report of the IPCC*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, 2007. 162
- Irizarry-Ortiz, M. M., Wang, G., and Eltahir, E. A. B.: Role of the biosphere in the mid-Holocene climate of West Africa, *J. Geophys. Res.-Atmos.*, 108, 4042, doi:10.1029/2001JD000989, 2003. 149
- 20 Janssen, R. H. H., Meinders, M. B. J., Van Nes, E. H., and Scheffer, M.: Microscale vegetation–soil feedback boosts hysteresis in a regional vegetation–climate system, *Global Change Biol.*, 14, 1104–1112, doi:10.1111/j.1365-2486.2008.01540.x, 2008. 149
- 25 Jin, F.-F., Neelin, J. D., and Ghil, M.: El Niño on the Devil’s Staircase: annual subharmonic steps to chaos, *Science*, 264, 70–72, 1994. 158
- Källén, E., Crafoord, C., and Ghil, M.: Free Oscillations in a Climate Model with Ice-Sheet Dynamics, *J. Atmos. Sci.*, 36, 2292–2303, 1979. 148, 149, 157
- Le Treut, H. and Ghil, M.: Orbital forcing, climatic interactions, and glaciation cycles, *J. Geophys. Res.-Oceans*, 88, 5167–5190, doi:10.1029/JC088iC09p05167, 1983. 148, 149, 159
- 30 Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J.: Tipping elements in the Earth’s climate system, *P. Natl. Acad. Sci. USA*, 105, 1786–1793, doi:10.1073/pnas.0705414105, 2008. 148

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Lorenz, E. N.: Deterministic nonperiodic flow, *J. Atmos. Sci.*, 20, 130–141, 1963. 148

Lovelock, J. E. and Kump, L. R.: Failure of climate regulation in a geophysiological model, *Nature*, 369, 732–734, doi:10.1038/369732a0, 1994. 162

Meir, P., Cox, P., and Grace, J.: The influence of terrestrial ecosystems on climate, *Trends Ecol. Evol.*, 21, 254–260, doi:10.1016/j.tree.2006.03.005, 2006. 147, 162

Meissner, K. J., Weaver, A. J., Matthews, H. D., and Cox, P. M.: The role of land surface dynamics in glacial inception: a study with the UVic Earth System Model, *Clim. Dynam.*, 21, 515–537, doi:10.1007/s00382-003-0352-2, 2003. 159

Nevison, C., Gupta, V., and Klinger, L.: Self-sustained temperature oscillations on Daisyworld, *Tellus B*, 51, 806–814, doi:10.1034/j.1600-0889.1999.t01-3-00005.x, 1999. 149, 157, 158

North, G. R., Cahalan, R. F., and Coakley, J.: Energy balance climate models, *Rev. Geophys. Space GE*, 19, 91–121, 1981. 153

Renssen, H., Brovkin, V., Fichefet, T., and Goosse, H.: Holocene climate instability during the termination of the African Humid Period, *Geophys. Res. Lett.*, 30, 1184, doi:10.1029/2002GL016636, 2003. 149

Roques, L., Chekroun, M. D., Cristofol, M., Soubeyrand, S., and Ghil, M.: Parameter estimation for energy balance models with memory, *P. Roy. Soc. A*, 470, 20140349, 2014. 158

Saltzman, B.: Climatic systems analysis, *Adv. Geophys.*, 25, 173–233, 1983. 148, 149

Schneider, S. H. and Dickinson, R. E.: Climate modeling, *Rev. Geophys. Space GE*, 12, 447–493, 1974. 147, 161

Sellers, W. D.: A global climatic model based on the energy balance of the earth–atmosphere system, *J. Appl. Meteorol.*, 8, 392–400, doi:10.1175/1520-0450(1969)008<0392:AGCMBO>2.0.CO;2, 1969. 150, 151, 152

Shepon, A. and Gildor, H.: The lightning-biota climatic feedback, *Global Change Biol.*, 14, 440–450, doi:10.1111/j.1365-2486.2007.01501.x, 2008. 147

Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M. (Eds.): IPCC, 2013: Climate Change 2013: The Physical Science Basis, in: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 2013. 151

Stommel, H.: Thermohaline convection with two stable regimes of flow, *Tellus*, 13, 224–230, 1961. 148, 149

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Svirezhev, Y. M. and von Bloh, W.: A minimal model of interaction between climate and vegetation: qualitative approach, *Ecol. Model.*, 92, 89–99, doi:10.1016/0304-3800(95)00198-0, 1996. 149, 160

Svirezhev, Y. M. and von Bloh, W.: Climate, vegetation, and global carbon cycle: the simplest zero-dimensional model, *Ecol. Model.*, 101, 79–95, doi:10.1016/S0304-3800(97)01973-X, 1997. 149, 152

Tziperman, E., Stone, L., Cane, M., and Jarosh, H.: El Niño chaos: overlapping of resonances between the seasonal cycle and the Pacific ocean–atmosphere oscillator, *Science*, 264, 72–74, 1994. 158

Watson, A. J. and Lovelock, J. E.: Biological homeostasis of the global environment: the parable of Daisyworld, *Tellus B*, 35, 284–289, doi:10.1111/j.1600-0889.1983.tb00031.x, 1983. 148

Weaver, I. S. and Dyke, J. G.: The importance of timescales for the emergence of environmental self-regulation, *J. Theor. Biol.*, 313, 172–180, doi:10.1016/j.jtbi.2012.07.034, 2012. 159

Wood, A. J., Ackland, G. J., Dyke, J. G., Williams, H. T. P., and Lenton, T. M.: Daisyworld: a review, *Rev. Geophys.*, 46, RG1001, doi:10.1029/2006RG000217, 2008. 148

Zaliapin, I. and Ghil, M.: Another look at climate sensitivity, *Nonlin. Processes Geophys.*, 17, 113–122, doi:10.5194/npg-17-113-2010, 2010. 151

Zeng, N. and Neelin, J. D.: The role of vegetation–climate interaction and interannual variability in shaping the African savanna, *J. Climate*, 13, 2665–2670, 2000. 147, 161

Zeng, N., Neelin, J. D., Lau, K.-M., and Tucker, C. J.: Enhancement of interdecadal climate variability in the Sahel by vegetation interaction, *Science*, 286, 1537–1540, doi:10.1126/science.286.5444.1537, 1999. 147, 161

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Definition and values of the model parameters.

Symbol	Meaning	Value
C_T	Heat capacity	$500 \text{ W yr K}^{-1} \text{ m}^{-2}$
Q_0	Incoming solar energy	342.5 W m^{-2}
ρ	Fraction of land	0.3
α_v	Albedo of vegetation	0.1
α_g	Albedo of ground	0.4
α_{\max}	Albedo of ice-covered ocean	0.85
α_{\min}	Albedo of ice-free ocean	0.25
$T_{\alpha,\ell}$	Temperature below which ocean is ice-covered	263 K
$T_{\alpha,u}$	Temperature above which ocean is ice-free	300 K
B_0	Constant in outgoing radiation	200 W m^{-2}
B_1	Constant in outgoing radiation	$2.5 \text{ W K}^{-1} \text{ m}^{-2}$
T_{opt}	Optimal growth temperature	283 K
k	Parameter for width of growth curve	$0.004 \text{ yr}^{-1} \text{ K}^{-2}$
γ	Death rate of vegetation	0.1 yr^{-1}

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil




Figure 1. Schematic representation of the model planet's surface, including a fraction p of land and $1-p$ of ocean. The fraction of the ocean that is covered by sea ice depends on the global temperature T , while the land is covered by a fraction A of vegetation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil




Figure 2. Fraction of sea ice cover (blue) and ocean albedo (red) as a function of temperature T . The ocean albedo is a weighted average of the albedo of open water and of ice, and is therefore directly dependent on the fraction of sea ice.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. Outgoing energy as function of temperature for different parametrizations.

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil




Figure 4. The (T, A) phase plane of the model, with the two null isoclines given by Eq. (6) (red straight line) and Eq. (7) (blue curve), respectively. The fixed points for the vegetated states of our planet are the intersections of these two curves. The non-vegetated fixed point for $A \neq 0$ does not appear in this plot.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil




Figure 5. The model's phase plane with the two nullclines (dashed lines) and the three fixed points (open and filled dots); the two filled dots are stable, while the open one is unstable. The stable manifold of the saddle (open dot) is shown in black and the unstable one in magenta. The coordinates of the fixed points are: (242.04, 0) for the unvegetated state, and (280.8, 0.898) and (297.9, 0.129) for the unstable and stable vegetated states, respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil




Figure 6. Bifurcation diagram, with the vegetation A plotted vs. the death rate γ . The branches were computed using the software package XPPAUT (Ermentrout, 2002).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Oscillations in a simple climate–vegetation model

J. Rombouts and M. Ghil




Figure 7. Typical oscillatory model solution, shown here for $\gamma = 0.02$, i.e. $\gamma/C_T = 1/10$. **(a)** Temperature and vegetation cover; and **(b)** fraction of the ocean that is covered by sea ice. The period of these regular oscillations is a little under 1000 years.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)