

This discussion paper is/has been under review for the journal Nonlinear Processes in Geophysics (NPG). Please refer to the corresponding final paper in NPG if available.

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

The Pennsylvania State University, University Park, Pennsylvania, 16802, USA

Received: 30 October 2015 – Accepted: 22 November 2015 – Published: 21 December 2015

Correspondence to: J. A. Schulte (jas6367@psu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union & the American Geophysical Union.

NPGD

2, 1705–1737, 2015

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Methods for detecting and quantifying nonlinearities in nonstationary time series are introduced and developed. In particular, higher-order wavelet analysis was applied to an ideal time series and the Quasi-biennial Oscillation (QBO) time series. Multiple-testing problems inherent in wavelet analysis were addressed by controlling the false discovery rate. A new local autobicoherence spectrum facilitated the detection of local nonlinearities and the quantification of cycle geometry. The local autobicoherence spectrum of the QBO time series showed that the QBO time series contained a mode with a period of 28 months that was phase-coupled to a harmonic with a period of 14 months. An additional nonlinearly interacting triad was found among modes with periods of 10, 16, 26 months. Local biphase spectra determined that the nonlinear interactions were not quadratic and that the effect of the nonlinearities was to produce non-smoothly varying oscillations. The oscillations were found to be skewed so that negative QBO regimes were preferred, and also asymmetric in the sense that phase transitions between the easterly and westerly phases occurred more rapidly than those from westerly to easterly regimes.

1 Introduction

Spectral analysis is a tool for extracting embedded structures in a time series. In particular, Fourier analysis has been used extensively by researchers for extracting deterministic structures from time series but is incapable of detecting nonstationary features often present in geophysical time series. Wavelet analysis can extract transient features embedded in time series, with a wavelet power spectrum representing variance (power) of a time series as a function of time and period. Since the seminal work of Torrence and Compo (1998), wavelet analysis has been applied extensively to geophysical time series such as the indices for the North Atlantic Oscillation (Olsen et al., 2012), Arctic Oscillation (Jevrejeva et al., 2003), Pacific Decadal Oscillation (Macdonald and

NPGD

2, 1705–1737, 2015

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Case, 2005; Newmann et al., 2003), El-Niño/Southern Oscillation (ENSO; Torrence and Webster, 1999), Pacific-North American Pattern, and West Pacific pattern (Gan et al., 2007). The application of wavelet coherence and cross-wavelet analyses (Grinsted et al., 2004), moreover, has proven useful in relating geophysical time series to other time series (Jevrejeva et al., 2003; Gan et al., 2007; Labat, 2010; Lee and Lwiza, 2008).

Many statistical methods, including power and cross-spectral analyses, rely on the assumption that the variable in question is Gaussian distributed (King, 1996). For a linear system in which the output is proportional to the input, the first- and second-order moments, the mean and variance, can fully describe the distribution of a process. In the frequency domain, by analogy, the variable can be fully described by the power spectrum, the decomposition of variance as a function of frequency. Suppose, however, that the distribution is non-Gaussian so that higher-order moments such as skewness and kurtosis exist. In this case, the mean and variance, while useful, are unable to fully describe the distribution in question. In a time series context, non-Gaussian distributions can arise from nonlinear systems, systems for which the output is no longer simply proportional to the input. For a nonlinear system, if the input is the sum of two sinusoids with different frequency components the output will contain additional frequency components representing the sum and difference of the input frequencies (King, 1996). In such cases, it is necessary to examine the decomposition of higher-order moments in frequency space.

The frequency decomposition of the third-order moment, for example, results in a bispectrum or skewness function that measure deviations from Gaussianity (Nikias and Raghuveer, 1987; King, 1996). In fact, Hinich (1985) developed a bispectral test to determine if a time series is non-Gaussian and nonlinear. In some situations, higher-order nonlinearities such as cubic nonlinearities may exist, in which case the trispectrum or other polyspectra would have to be used (Collis et al., 1998).

Another advantage of higher-order spectral analysis is that the cycle geometry of oscillations, such as asymmetry with respect to a horizontal axis (skewed oscillation)

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



or with respect to a vertical axis (asymmetric oscillation) can be quantified using the biphase. A pure sine wave, for example, is neither skewed nor asymmetric, whereas a time series resembling a saw-tooth is asymmetric. Skewed and asymmetric cycle geometry can identify, for example, abrupt climatic shifts, sudden shifts in the climate system that exceed the magnitude of the background variability (King, 1996). Abrupt climate shifts have occurred numerous times in the past and have dire impacts on ecological and economic systems (Alley et al., 2005). An understanding of past abrupt climate shifts is essential to understanding future climate change and so there is a need to quantify nonlinearities present in climatic oscillations.

The Quasi-biennial Oscillation (QBO), as another example, has been shown to behave nonlinearly, transitioning from easterly phases to westerly phases more rapidly than from westerly to easterly phases (Lu et al., 2009). Another source of asymmetry in the QBO time series arises from the westerly shear zone descending more regularly than the easterly shear zone. Asymmetries in the QBO time series are not well-captured by linear methods such as linear principal component and singular spectrum analyses (Lu et al., 2009) but are better captured using, for example, nonlinear principal component analysis (Hamilton and Hsieh, 2002). Another example of a nonlinear time series is the sunspot cycle. Solar activity undergoes an 11 year oscillation characterized by asymmetric cycle geometry, with solar maxima generally rising faster than they fall, indicating the presence of nonlinearities (Moussas et al., 2005; Rusu, 2007). ENSO, a climate phenomenon with regional- to global-scale impacts, has also been shown to exhibit nonlinearities (Timmermann, 2003). The presence of nonlinearities and possible nonstationarities in the QBO, ENSO, and sunspot time series makes traditional Fourier and wavelet analysis inadequate for feature extraction, underscoring the need to develop methods for quantifying nonlinearities in a nonstationary geophysical setting.

The application of higher-order wavelet analysis has been rather limited compared to traditional wavelet analysis (van Millagan et al., 1995; Elsayed, 2006). One geophysical application of higher-order wavelet analysis is to oceanic waves (Elsayed, 2006),

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



local autobicoherence spectrum is introduced, eliminating the selection of a time interval on which to calculate nonlinear properties of time series. The new and existing methods are applied to an ideal time series and the QBO index. In Sect. 4, a new procedure for estimating confidence intervals of global autobicoherence quantities is developed to estimate uncertainties in the sample autobicoherence spectra. The application of the new procedure to the sample autobicoherence spectrum of the QBO time series is then used to further assess confidence in results.

2 Wavelet analysis

The idea behind wavelet analysis is to convolve a time series with a function satisfying certain conditions. Such functions are called wavelets, of which the most widely used is the Morlet wavelet, a sinusoid damped by a Gaussian envelope:

$$\psi_0(\eta) = \pi^{-1/4} e^{i\omega_0\eta} e^{-\frac{1}{2}\eta^2} \quad (1)$$

where ψ_0 is the Morlet wavelet, ω_0 is the dimensionless frequency, and η is the dimensionless time (Torrence and Compo, 1998; Grinsted et al., 2004). In practical applications, the convolution of the wavelet function with a time series $X = (x_n; n = 1, \dots, N)$ is calculated discretely using

$$W_n^X(s) = \sqrt{\frac{\delta t}{s}} \sum_{n'=1}^N x_{n'} \psi_0 \left[(n' - n) \frac{\delta t}{s} \right] \quad (2)$$

where δt is a uniform time step, s is scale, $\eta = s \times t$, and $W_n^X(s)$ is the wavelet transform. The wavelet power is given by $\left| W_n^X(s) \right|^2$ (Torrence and Compo, 1998; Grinsted et al., 2004). For the Morlet wavelet with $\omega_0 = 6$, the wavelet scale and the Fourier period λ are approximately equal ($\lambda = 1.03s$). A more detailed discussion of wavelet analysis can be found in Torrence and Compo (1998).

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Shown in Fig. 1a is the time series of the QBO index and shown in Fig. 1b is the corresponding wavelet power spectrum. The QBO data from 1950–2013 were obtained from the Climate Prediction Center. The QBO index is defined as the zonal average of the 30 hPa zonal wind at the equator. As such, a positive index indicates westerly winds and a negative index indicates easterly winds. The most salient feature of the time series is the rather regular periodicity of approximately 28 months. Also note the asymmetry between the negative and positive phase, with the negative phases generally being stronger. The periodic behavior of the QBO was corroborated by examining the wavelet power spectrum. A well-defined 28 month periodicity is evident, with the associated wavelet power changing little throughout the study period.

There are also secondary features located at a period of approximately 14 months, primarily from 1985 to 2013. The appearance of significant power at a period of 14 months also coincides with most of the largest negative phases of the QBO. Such a correspondence may not have been a coincidence; the 14 month mode and the 28 month mode may have interacted constructively to generate large negative events but interacted destructively to create smaller positive events. However, additional tools are needed to confirm if the periodicities are interacting and to understand how the interactions were related to the behavior of the QBO.

3 Higher-order wavelet analysis

3.1 Wavelet-based autobicoherence

Higher-order spectral analysis provides the opportunity to quantify nonlinearities and allows the detection of interacting oscillatory modes within a time series. More specifically, nonlinearities are quantified using bicoherence, a tool for measuring quadratic nonlinearities, where quadratic nonlinearities imply that for frequencies f_1 , f_2 , and f_3 and corresponding phases ϕ_1 , ϕ_2 , and ϕ_3 the sum rules

$$f_1 + f_2 = f_3 \quad (3)$$

and

$$\phi_1 + \phi_2 = \phi_3 \quad (4)$$

are satisfied. Whereas Eq. (3) implies frequency coupling, Eq. (4) implies phase coupling. To see from where Eqs. (3) and (4) originate, let

$$X(t) = \sin(2\pi f_1 t + \phi_1) + \sin(2\pi f_2 t + \phi_2) \quad (5)$$

be the input into a system whose output is related to the input by

$$Y(t) = X(t) + \varepsilon X(t)^2 + w(t). \quad (6)$$

The multiplicative factor ε is used to represent the contribution of the nonlinear component of the signal and $w(t)$ is Gaussian white noise. Note that if $\varepsilon = 0$, then the system is linear because the output contains the same frequency components as the input. The substitution of Eq. (5) into Eq. (6) results in

$$Y(t) = \sin(2\pi f_1 t + \phi_1) + \sin(2\pi f_2 t + \phi_2) + \frac{\varepsilon}{2} [1 - \cos(2(2\pi f_1 t + \phi_1)) - \cos(2(2\pi f_2 t + \phi_2)) + \cos(2\pi(f_2 - f_1)t + \phi_2 - \phi_1) - \cos(2\pi(f_1 + f_2)t + \phi_1 + \phi_2)] + w(t) \quad (7)$$

and thus the output has sinusoids with additional frequency components $2f_1$, $2f_2$, $f_2 - f_1$, and $f_2 + f_1$, which arise from the second term in right-hand side of Eq. (6).

Unlike the power spectrum, which is the Fourier transform of the second-order moment of a time series, the bispectrum is defined as the double Fourier transform of the third-order moment, or, more generally, the third-order cumulant, i.e.,

$$b_{xxx}(f_1, f_2) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} C(t_1, t_2) e^{-i2\pi(f_1 t_1 + f_2 t_2)} dt_1 dt_2, \quad (8)$$

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



T is a time interval, $W_x(s, t)$ is the wavelet transform of a time series X at scale s and time t , and $W_x^*(s, t)$ denotes the complex conjugate of $W_x(s, t)$. The wavelet-based autobicoherence measures the degree of quadratic phase coupling, where a peak at (s_1, s_2) indicates a nonlinear interaction among the scale components s_1 , s_2 , and s .

In practice, the autobicoherence is computed discretely so that Eq. (13) can be written as

$$\overline{W_b}(s_1, s_2) = \frac{|B_{xxx}^w(s_1, s_2)|^2}{\left(\sum_{n=n_1}^{n_2} |W_n^X(s_1) W_n^X(s_2)|^2 \right) \left(\sum_{n=n_1}^{n_2} |W_n^X(s)|^2 \right)}, \quad (16)$$

where

$$\begin{aligned} B_{xxx}^w(s_1, s_2) &= \sum_{n=n_1}^{n_2} W_n^{*X}(s) W_n^X(s_1) W_n^X(s_2) \\ &= \sum_{n=n_1}^{n_2} B_n^w(s_1, s_2), \end{aligned} \quad (17)$$

$n_1 \geq 1$, and $n_2 \leq N$. Note that if $n_1 = 1$ and $n_2 = N$, then Eq. (16) represents the global autobicoherence spectrum.

The Monte Carlo approach to pointwise significance testing is adopted in this paper and is similar to that used in wavelet coherence (Grinsted et al., 2014). To estimate the significance of wavelet-based autobicoherence at each point (s_1, s_2) , Monte Carlo methods are used to (1) generate a large ensemble of red-noise processes with the same lengths and lag-1 autocorrelation coefficients as the input time series and (2) compute for each randomly generated red-noise process the autobicoherence spectrum. From the ensemble of autobicoherence spectra, the $p = 100(1 - \alpha_p)$ percentile of the autobicoherence estimates is computed for every point (s_1, s_2) , where p

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



corresponds to the critical level of the test and α_p is the pointwise significance level of the test. Given the symmetry of the autobicoherence spectrum, the critical level of the test can be computed using only half of the autobicoherence estimates, reducing computational costs.

3.2 Multiple testing

Let α_p be the significance level of the pointwise significance test as described above and let K denote the number of autobicoherence estimates being tested, then there will be on average $\alpha_p K$ false positive results. A similar problem occurs in traditional wavelet analysis (Maraun et al., 2007; Schulte et al., 2015a). In the case of simultaneously testing multiple hypotheses, the number of false positive results can be reduced by applying, for example, the Bonferroni correction (Lehmann, 1986). However, this simple correction often results in many true positives being rejected and is especially permissive in the case of autocorrelated data (Maraun et al., 2004). Other procedures also exist, including the Walker p value adjustment procedure, which has more statistical power than the Bonferroni correction. An even more powerful method is the Benjamini and Hochberg (1995) procedure, which controls the false discovery rate (FDR), where the FDR is the expected proportion of the false rejections that are actually true. An advantage of this method, in addition to its statistical power, is that it takes into account the confidence with which local hypotheses are rejected and is robust even in the case of autocorrelated data (Wilks, 2002). Benjamini and Yekutieli (2001) developed a modified version of the Benjamini and Hochberg (1995) procedure that works for any dependency structure among the local test statistics and thus this procedure will be used in this paper to control the FDR.

The procedure can be described as follows: suppose that K local hypotheses were tested. Let $p_{(i)}$ denote the smallest of the K local p values, then, under the assumption that the K local tests are independent, the FDR can be controlled at the q level by rejecting those local tests for which $p_{(i)}$ is no greater than

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\begin{aligned} \rho_{\text{FDR}} &= [\rho_{(j)} : \rho_{(j)} \leq q (j/K)] \\ &= [\rho_{(j)} : \rho_{(j)} \leq \alpha_{\text{global}} (j/K)] \end{aligned} \quad (18)$$

so that the FDR level is equivalent to the global test level. For a local p value to be deemed significant using this procedure, it must be less than or equal to the largest p value for which Eq. (18) is satisfied. If no such local p values exist, then none are deemed insignificant, and, therefore, the global test hypothesis cannot be rejected. If the test statistics have an unknown dependency structure, q can be replaced with $q / \sum_{i=1}^K \frac{1}{i}$, though this substitution makes the procedure less powerful (Reiner et al., 2002). This modified method will be applied to autobicoherence spectra at the 0.05 level throughout this paper.

3.3 Wavelet-based autobicoherence of an idealized time series

To demonstrate the features of a time series that can be extracted using higher-order wavelet analysis, an idealized nonstationary time series will first be considered. Consider the quadratically nonlinear time series

$$X(t) = \cos(2\pi f t + \phi) + \gamma(t) \cos(4\pi f t + 2\phi) + w(t), \quad (19)$$

where f is frequency, $w(t)$ is Gaussian white noise, and $\gamma(t)$ is a time-dependent nonlinear coefficient given by

$$\gamma(t) = 0.001t. \quad (20)$$

Note that Eqs. (3) and (4) are satisfied because $f_1 + f_2 = 2f_1 = 2f_2$ and similarly for ϕ . The sinusoid with frequency $2f_1$ is said to be the harmonic of the primary frequency component with frequency f_2 , where the amplitude of the harmonic depends on $\gamma(t)$, the strength of the quadratic nonlinearity. $X(t)$ and the corresponding wavelet power

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimates is also impacted. In particular, smoothing the autobicoherence spectrum with respect to s_{\max} allowed extrema to be smoothed out, eliminating spuriously large autobicoherence. For this reason, all local autobicoherence spectra in this paper will be computed by smoothing with respect to s_{\max} .

The advantage of using Eq. (21) is that transient quadratic nonlinearities can now be detected and the need for choosing an integration time interval has been eliminated. If $s_1 = s_2$, then $(t, s_1, s_1) = (t, s_2, s_2) = (t, 2s)$ and thus, in the case of this diagonal slice, the local wavelet-based bicoherence spectrum is a two-dimensional representation of the degree of local quadratic nonlinearity. The vertical axis corresponds to the primary frequency and the horizontal axis corresponds to time. As a concrete example, a peak at (64, 64) would indicate that at time index $t = 50$ the oscillation with a fundamental period $\lambda = 1.03(2s) \approx 64$ is locally coupled to an oscillation with period $\lambda \approx 32$.

One can also compute a local biphas from the smoothed bispectrum by taking the four quadrant inverse tangent of the smoothed imaginary part divided by the smoothed real part. The local biphas, for example, was computed for the skewed time series shown in Fig. 2a. As expected, the biphas fluctuates regularly around 0° and the mean is 2° . The local biphas for the saw-toothed time series is shown in Fig. 3b. The biphas fluctuates about 90° and the mean biphas is 90° as expected.

The procedure for the estimation of the statistical significance of local autobicoherence is the following: generate red-noise time series with the same lag-1 autocorrelation coefficients as the input time series and use the local autobicoherence estimates outside the COI to generate a null distribution of $b_n^W(s_1, s_2)$. Note that the calculation only needs to be performed at a fixed time outside of the COI because red-noise is a stationary process, which produces a stationary background spectrum.

3.6 Local wavelet-based autobicoherence of an idealized time series

The local autobicoherence spectrum of $X(t)$ for (30, 30) is shown in Fig. 6b. Initially, there is no local autobicoherence that exceeds the 5% significance level. At $t = 250$ and $t = 500$, on the other hand, small regions of 5% significant autobicoher-

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ence emerge, indicating a transient nonlinear interaction. At $t = 500$ the nonlinearity is strong and results in a large region of significant local autobicoherence extending from $t = 500$ to the edge of the wavelet domain.

In order to determine if the peaks in autobicoherence are associated with a quadratic nonlinearity, it is important to compute the biphas, which is shown in Fig. 7b. From $t = 0$ to $t = 400$ there is an unstable phase relationships between the phase of the primary frequency component and its harmonic. Such a lack of phase coherence indicates a weak nonlinear interaction, which is consistent with how the autobicoherence is lower before $t = 400$. In contrast, after $t = 400$, the biphas becomes stable, changing little with time, indicating a consistent phase relationship between the primary frequency mode and its harmonic. It also noted that the biphas during this time fluctuates near 0° , which implies that the phase relationships arise from a quadratic nonlinearity. The near zero biphas is consistent with how $X(t)$ was constructed from the sum of two cosines with zero phase and also suggests that the interaction results in skewed cycle geometry, where positive values of the time series are preferred. Indeed, by inspection of Fig. 4a the oscillations initially appear to be sinusoidal, varying smoothly, whereas after $t = 400$ spikes begin to appear and $X(t)$ favors positive values.

3.7 Local wavelet-based autobicoherence of the QBO time series

The local autobicoherence spectrum of the QBO index at the point (28, 28) in the full autobicoherence spectrum is shown in Fig. 8. From 1950 to 1970 the magnitude of the autobicoherence fluctuated and consisted of one local significant peak at 1965. Significant autobicoherence was also found from 1975 to 1998, contrasting with the autobicoherence after 1998, which was not found to be significant until 2010.

To determine if the peaks indicated in the autobicoherence are associated with a quadratic nonlinearity, the local biphas was computed. Figure 8a shows the local biphas for the autobicoherence peak at (28, 28). For most of the study period, the biphas was found to vary considerably, particularly during the 1950–1970 and 1995–2013 periods. On the other hand, the biphas varied smoothly from 1970 to 1995, con-

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sistent with how the autobicoherence during that period was large and stable (Fig. 8a). Also, during that period the biphasic was nonzero; in fact, the mean biphasic during the period was -100° , suggesting that the phase coupling is not the result of a quadratic interaction. A biphasic of -100° indicated asymmetric geometry, which physically represents how phase transitions of the QBO occurred at different rates. Recall that it has already been discussed in the introduction that the QBO transitions from easterly phases to westerly phases more rapidly than from westerly to easterly phases (Lu et al., 2009). Another interesting feature is the general increase in the biphasic from 1970 to 1995. In the beginning of the time period, the biphasic was -180° and after 1980 the biphasic switched to -90° .

The local autobicoherence and biphasic corresponding to the peak (16, 26) was also computed (Fig. 9). The mean of the absolute value of the biphasic for the period 1950–2013 was 130° , indicating that the interaction among the modes with periods of 10, 16, 26 months resulted in skewed waveforms. In fact, because the biphasics were close to 180° the waveforms should have been skewed to negative values (Maccarone, 2013) and such skewness is evident by inspecting Fig. 1. Also note that some of the largest negative phases of the QBO occurred from 1995 to 2010, which coincided with the period of most significant autobicoherence as shown in Fig. 9a.

4 Block bootstrapping methods

4.1 Block bootstrapping autobicoherence

Bootstrapping is a widely used technique to estimate the variance or uncertainty of a sample estimate. For independent data one samples with replacement individual data points (Efron, 1979); for dependent data one must sample with replacement blocks of data to preserve the autocorrelation structure of the data (Kunsch, 1989). The latter technique is called block bootstrapping and should be used for variance estimation of global wavelet quantities, as wavelet coefficients are known to be autocorrelated in

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



both time and scale. The use of traditional bootstrapping techniques would result in confidence intervals that are too narrow. It is expected, however, that the choice of the bootstrapping technique is more critical at larger scales, as the decorrelation length of the mother wavelet increases with scale.

5 A brief overview of the procedure is provided below but a more detailed discussion can be found in Schulte et al. (2015b). To find the approximate $100(1 - \beta)\%$ confidence interval of an autobicoherence estimate, divide the set of wavelet coefficients at each scale into overlapping blocks. The lengths of the blocks at each scale should be the same and the randomly resampled blocks chosen should be the same at each scale to avoid randomizing the data. The concatenation of the blocks then results in a synthetic set of wavelet coefficients at each scale. The synthetic set of wavelet coefficients can then be used to calculate a bootstrap replicate of the autobicoherence. The iteration of the procedure 1000 times results in a distribution of bootstrap replicates from which a 95 % confidence interval can be obtained.

10 As noted by Schulte et al. (2015b), the appropriate block length to use can be determined by Monte Carlo methods. Using Monte Carlo methods, it was determined that a block length of $N^{0.6}$ was found to produce accurate confidence bounds while also producing the widest confidence intervals at all scales.

4.2 Application to ideal and climatic time series

20 Figure 5b shows the application of the block bootstrap procedure to the diagonal slice $s_1 = s_2 = 2s$ of the autobicoherence for the ideal case. The 95 % confidence intervals were also obtained using the ordinary bootstrap. A pronounced peak at $2s = 30$ was identified and represents the interaction between the primary frequency and its harmonic. By inspection of Fig. 5b, there is a clear difference between the widths of the confidence intervals obtained from the two bootstrapping procedures. For the ordinary bootstrap, the confidence intervals are narrow and the widths of the confidence intervals appear to be only weakly dependent on scale. On the other hand, the confidence intervals obtained using the block bootstrap procedure are wide, especially at large

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



scales, and the width of the confidence intervals depends strongly on scale, increasing from small scales to large scales. It is also noted that, whereas the block bootstrap procedure has deemed no spurious peaks as significant, the ordinary bootstrap procedure deemed two the spurious peaks at $2s = 14$ and $2s = 100$ as significant. The implementation of the block bootstrap procedure can therefore enhance confidence in results, facilitating the investigation of a deeper physical understanding.

The application of the block bootstrap procedure to the diagonal slice $s_1 = s_2 = 2s$ of the full autobicoherence spectrum of the QBO index is shown in Fig. 10. The 95% confidence intervals corresponding to the peaks (14, 14) and (28, 28) do not cross the 5% significance bound and thus one has more confidence that those peaks are significant. All other peaks have been deemed insignificant.

5 Summary

Higher-order wavelet analysis together with significance testing procedures were used to detect nonlinearities embedded in an ideal time series and the QBO time series. The autobicoherence spectrum of the QBO index revealed phase coupling of the 28 month mode with a higher frequency mode with period 14 months. A local autobicoherence spectrum of the QBO index showed that the strength of the nonlinearities varied temporally. Furthermore, the local biphase spectrum indicated that the nonlinear interaction resulted in waveforms that were both skewed and asymmetric, indicating that the strength of negative QBO events were stronger than positive events, and that transitions between events occurred at different rates.

Acknowledgements. Support for this research was provided by the National Science Foundation Physical Oceanography Program (award number 0961 423) and the Hudson River Foundation (award number GF/02/14).

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Alley, R. B., Marotzke, J., Nordhaus, W. D., Overpeck, J. T., Peteet, D. M., Pielke Jr., R. A., Pierrehumbert, R. T., Rhines, P. B., Stocker, T. F., Talley, L. D., and Wallace, J. M.: Abrupt climate change, *Science*, 299, 2005–2010, 2003.
- 5 Benjamini, Y. and Hochberg, Y.: Controlling the false discovery rate: a practical and powerful approach to multiple testing, *J. R. Stat. Soc.*, 57, 289–300, 1995.
- Benjamini, Y. and Yekutieli, D.: The control of the false discovery rate in multiple testing under dependency, *Ann. Stat.*, 29, 1165–1188, 2001.
- Collis, W. B., White, P. R., and Hammond, J. K.: Higher-order spectra: the bispectrum and trispectrum, *Mech. Syst. Signal Pr.*, 12, 375–394, 1998.
- 10 Efron, B.: Bootstrap Methods: another Look at the Jackknife, *Ann. Stat.*, 7, 1–26, 1979.
- Elgar, S. and Chandran, V.: Higher-order spectral analysis to detect nonlinear interactions in measured time series and an application to Chua's Circuit, *Int. J. Bifurcat. Chaos*, 3, 19–34, 1993.
- 15 Elgar, S. and Sebert, G.: Statistics of bicoherence and biphas, *J. Geophys. Res.*, 94, 10993–10998, 1989.
- Elsayed, M. A. K.: Wavelet bicoherence analysis of wind–wave interaction, *Ocean Eng.*, 33, 458–470, 2006.
- Gan, T. Y., Gobena, A. K., and Wang, Q.: Precipitation of Southwestern Canada: wavelet, scaling, multifractal analysis, and teleconnection to climate anomalies, *J. Geophys. Res.*, 112, D10110, doi:10.1029/2006JD007157, 2007.
- 20 Grinsted, A., Moore, J. C., and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, *Nonlin. Processes Geophys.*, 11, 561–566, doi:10.5194/npg-11-561-2004, 2004.
- 25 Hagelberg, T., Pisas, N., and Elgar, S.: Linear and nonlinear couplings between orbital forcing and the marine $\delta^{18}\text{O}$ record during the Late Neocene, *Paleoceanography*, 6, 729–746, 1991.
- Hamilton, K. and Hsieh, W. W.: Representation of the quasi-biennial oscillation in the tropical stratospheric wind by nonlinear principal component analysis, *J. Geophys. Res.*, 107, 4232, doi:10.1029/2001JD001250, 2002.
- 30 Higuchi, K., Huang, J., and Shabbar, A.: A wavelet characterization of the North Atlantic Oscillation variation and its relationship to the North Atlantic sea surface temperature, *Int. J. Climatol.*, 19, 1119–1129, 1999.

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jevrejeva, S., Moore, J. C., and Grinsted, A.: Influence of the Arctic Oscillation and El Niño–Southern Oscillation (ENSO) on ice conditions in the Baltic Sea: the wavelet approach, *J. Geophys. Res.*, 108, 4677, doi:10.1029/2003JD003417, 2003.
- King, T.: Quantifying nonlinearity and geometry in time series of climate, *Quaternary Sci. Rev.*, 15, 247–266, 1996.
- Kunsch, H. R.: The jackknife and the bootstrap for General Stationary Observations, *Ann. Stat.*, 17, 1217–1241, 1989.
- Labat, D.: Cross wavelet analyses of annual continental freshwater discharge and selected climate indices, *J. Hydrol.*, 385, 269–278, 2010.
- Lee, Y. J. and Lwiza, K. M.: Factors driving bottom salinity variability in the Chesapeake Bay, *Cont. Shelf Res.*, 28, 1352–1362, 2008.
- Lu, B. W., Pandolfo, L., and Hamilton, K.: Nonlinear representation of the Quasi-Biennial Oscillation, *J. Atmos. Sci.*, 66, 1886–1904, 2009.
- Maccarone, T. J.: The Biphase explained: understanding the asymmetries in coupled Fourier components of astronomical timeseries, *Mon. Not. R. Astron. Soc.*, 435, 3547, doi:10.1093/mnras/stu1824, 2013.
- MacDonald, G. M. and Case, R. A.: Variations in the Pacific Decadal Oscillation over the past millennium, *Geophys. Res. Lett.*, 32, L08703, doi:10.1029/2005GL022478, 2005.
- Maraun, D., Kurths, J., and Holschneider, M.: Nonstationary Gaussian processes in the wavelet domain: synthesis, estimation, and significance testing, *Phys. Rev. E*, 75, 016707, doi:10.1103/PhysRevE.75.016707, 2007.
- Moussas, X., Polygiannakis, J. M., Preka-Papadema, P., and Exarhos, G.: Solar cycles: a tutorial, *Adv. Space Res.*, 35, 725–738, 2005.
- Newman, M., Compo, G. P., and Alexander, M. A.: ENSO-forced variability of the Pacific decadal oscillation, *J. Climate*, 16, 3853–3857, 2003.
- Nidal, K. and Malik, A. S.: EEG/ERP Analysis: Methods and Applications, CRC Press, Boca Raton, Florida, 334 pp., 2013.
- Nikias, C. L. and Raghuvver, M. R.: Bispectrum Estimation: a Digital Signal Processing Framework, *Proc. IEEE*, 75, 869–891, 1987.
- Olsen, J., Anderson, J. N., and Knudsen, M. F.: Variability of the North Atlantic Oscillation over the past 5,200 years, *Nat. Geosci.*, 5, 808–812, 2012.
- Polygiannakis, J. M., Moussas, X., and Sonett, C. P.: A nonlinear RLC solar cycle model, *Sol. Phys.*, 163, 193–203, 1996.

**Wavelet analysis for
non-stationary,
non-linear time series**

J. A. Schulte




Figure 1. (a) The QBO index and (b) the corresponding wavelet power spectrum. Contours enclose regions of 5% statistical pointwise significance (Torrence and Compo, 1998). Light shading represents the cone of influence, the region in which edge effects cannot be ignored.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Wavelet analysis for
non-stationary,
non-linear time series**

J. A. Schulte




Figure 2. (a) A skewed time series and (b) its corresponding local biphase. The biphase close to zero indicates a nonlinear interaction resulting in a skewed oscillation. The biphase was calculated from the first three cosines in the summation described in the text. The large deviations from zero at the edges are the result of edge effects.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Wavelet analysis for
non-stationary,
non-linear time series**

J. A. Schulte




Figure 3. (a) A saw-toothed time series and (b) its corresponding local biphas. The biphas close to 90° indicates a nonlinear interaction resulting in an asymmetric waveform. The biphas was calculated from the first three cosines in the summation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Wavelet analysis for
non-stationary,
non-linear time series**

J. A. Schulte




Figure 4. (a) Time series corresponding to Eq. (19). (b) Corresponding wavelet power spectrum.

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte




Figure 5. (a) Wavelet-based autobicoherence spectrum of the ideal time series. Thick contours enclose regions of 5% pointwise significance after controlling the FDR. The diagonal line separates the spectrum into two symmetric regions. (b) The diagonal slice of the autobicoherence spectrum at $s_1 = s_2 = s$. The critical level for the test represented by the dotted line was calculated using Monte Carlo methods.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte




Figure 7. (a) The local autobicoherence and (b) local biphas corresponding to (30, 30) in the full autobicoherence spectrum shown in Fig. 5a. Biphases differing from 90° indicate that the nonlinear interaction resulted in a waveform with skewness.

**Wavelet analysis for
non-stationary,
non-linear time series**

J. A. Schulte




Figure 8. Same as Fig. 7 except at (28, 28) in the auto-coherence spectrum of the QBO index. Biphases differing from 90° indicate that the nonlinear interaction resulted in a waveform with skewness.

Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte




Figure 9. Same as Fig. 8 except at the point (16, 26).

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures

⏪
⏩

◀
▶

Back	Close
------	-------

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wavelet analysis for non-stationary, non-linear time series

J. A. Schulte




Figure 10. Same as Fig. 5b except for the QBO index for the period 1950–2013.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

