



Springtime contribution of dinitrogen fixation to primary production across the Mediterranean Sea

E. Rahav¹, B. Herut², A. Levi¹, M. R. Mulholland³, and I. Berman-Frank¹

¹Mina and Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat Gan 52900, Israel

²Israel Oceanographic and Limnological Research, National Institute of Oceanography, Haifa 31080, Israel

³Department of Ocean, Earth and Atmospheric Sciences, Old Dominion University, 4600 Elkhorn Avenue, Norfolk, Virginia 23 529-0276, USA

Correspondence to: I. Berman-Frank (ilana.berman-frank@biu.ac.il)

Received: 6 December 2012 – Published in Ocean Sci. Discuss.: 4 January 2013

Revised: 5 April 2013 – Accepted: 5 April 2013 – Published: 14 May 2013

Abstract. Dinitrogen (N_2) fixation rates were measured during early spring across the different provinces of Mediterranean Sea surface waters. N_2 fixation rates, measured using $^{15}N_2$ enriched seawater, were lowest in the eastern basin and increased westward with a maximum at the Strait of Gibraltar (0.10 to 2.35 nmol $NL^{-1} d^{-1}$, respectively). These rates were 3–7 fold higher than N_2 fixation rates measured previously in the Mediterranean Sea during summertime and we estimated that methodological differences alone did not account for the seasonal changes we observed. Higher contribution of N_2 fixation to primary production (4–8 %) was measured in the western basin compared to the eastern basin (~2 %). Our data indicates that these differences between basins may be attributed to changes in N_2 -fixing planktonic communities and that heterotrophic diazotrophy may play a significant role in the eastern Mediterranean while autotrophic diazotrophy has a more dominant role in the western basin.

lem, is the export of underlying, nutrient-rich intermediate-depth water to the North Atlantic Ocean through the Strait of Gibraltar (Moutin and Raimbault, 2002; Krom et al., 2010).

Dissolved inorganic nitrogen (NO_3^- , NO_2^- , NH_4^+) is considered the proximate limiting nutrient for primary productivity in many oceanic regions (Falkowski, 1998), especially in low nutrient, low chlorophyll (LNLC) environments. While traditionally the MS has been considered phosphorus (P) limited (Krom et al., 1991; Thingstad et al., 1998), more recent publications demonstrate nitrogen (N) limitation or N and P co-limitation across the two sub-basins within the MS (Thingstad et al., 2005, Tanaka et al., 2011). Diazotrophs (i.e. N_2 fixers) are likely to have an advantage in N-limited environments because they are able to utilize the abundant dissolved N_2 , unavailable to most organisms, as an N source for growth (Capone and Montoya, 2001; Zehr and Ward, 2002).

Prokaryotic dinitrogen (N_2) fixation is now recognized as a globally important input of new oceanic N (reviewed in Gruber, 2008) that can be subsequently transferred to other planktonic groups (Mulholland et al., 2004; Mulholland and Capone, 2009). However, reported rates of N_2 fixation from the MS are limited to a few studies from the last six years and most are restricted to surface waters and the summer season. Typical rates of N_2 fixation during summer from both the eastern and western basins of the MS are generally low, ranging from undetectable to ~0.15 nmol $NL^{-1} d^{-1}$ (Ibello et al., 2010; Ridame et al., 2011; Yogeve et al., 2011; Rahav et al., 2013); however, N_2 fixation rates at the central zone of the Ligurian Sea station in the NW Mediterranean (DYnamique des

1 Introduction

The Mediterranean Sea (MS) is frequently described as a “blue desert” with low phytoplankton biomass and primary production (Berman et al., 1984; Bosc et al., 2004; Ignatiades et al., 2009; Siokou-Frangou et al., 2010). The low primary production is due to the low concentration and supply of dissolved nutrients in surface waters during most of the year and this is exacerbated during spring through late fall when the water column is thermally stratified. Compounding the prob-




Fig. 1. Map of the sampling locations (triangles): NW Levantine basin (St. 290), anticyclonic Shikmona eddy (St. 294), Ionian Sea (St. 304), Adriatic Sea (St. 312), Tyrrhenian Sea (St. 316), Alboran Sea (St. 333), Strait of Gibraltar (St. 338) and Gulf of Cadiz (St. 339). Background (circle): spatial distribution of chlorophyll *a* concentrations in surface waters (6–8 m) along the R/V *Meteor* M84/3 cruise track ($n = 94$).

Flux de matière en MEDiterranée- DYFAMED) are higher ranging from 2–17 nmol NL⁻¹ d⁻¹ (Garcia et al., 2006; Sandroni et al., 2007).

Diazotrophs contributing to N₂ fixation in the MS have been partially characterized (Man-Aharonovich et al., 2007; Bar Zeev et al., 2008; Le Moal and Biegala, 2009; Le Moal et al., 2011; Yogeve et al., 2011). In the MS organisms expressing *nifH*, a gene encoding part of the nitrogenase complex, include unicellular cyanobacteria, diatom-diazotroph assemblages, proteobacteria, methanogenic archaea, anaerobic bacteria, and purple sulfur bacteria. (Man-Aharonovich et al., 2007; Yogeve et al., 2011). The filamentous cyanobacterium *Trichodesmium* has been sporadically observed in extremely low abundances (Yogeve et al., 2011); one bloom of this genus was recorded from the Aegean Sea near Lesbos Island (Spatharis et al., 2012).

The contribution of N₂ fixation to new primary productivity in the MS was mostly examined during the stratified period in summer and appears to vary between the eastern and western basins. In the western basin, N₂ fixation was shown to contribute up to 35% to new primary production during the stratified period (Bonnet et al., 2011), while in the Levantine basin and the eastern Mediterranean Sea (EMS), N₂ fixation contributed only ~0.5–2% to the new production (Yogeve et al., 2011, Rahav et al., 2013). Yearly variability in the contribution of N₂ fixation to new primary productivity was also observed in the DYFAMED station ranging from 1% to 28% (Sandroni et al., 2007).

Here we present N₂ fixation and carbon uptake rate measurements from surface waters collected from a transect across the Mediterranean Sea during spring (before summer stratification). We calculate the contribution of diazotrophy to primary production in spring and compare these with similar measurements made during the stratified summer period to provide a more comprehensive seasonal assessment of N₂ fixation in the Mediterranean Sea. Additionally, we assessed

the relative contribution of heterotrophic versus autotrophic diazotrophy during springtime across the MS.

2 Material and methods

2.1 Sampling locations

This research was carried out aboard the R/V *Meteor* (cruise M84/3) between 4 and 28 April 2011. Eight stations were sampled along an east to west transect across the Mediterranean Sea, each representing a different water mass with associated mesoscale characteristics. Stations included: the NW Levantine basin (St. 290), the anti-cyclonic Shikmona eddy (St. 294), the Ionian Sea (St. 304), the Adriatic Sea (St. 312), the Tyrrhenian Sea (St. 316), the Alboran Sea (St. 333), Strait of Gibraltar (St. 338), and the Gulf of Cadiz (St. 339) (Fig. 1 and Table 1). Seawater samples collected east of the Sicily strait were defined as eastern Mediterranean (EMS) stations, whereas samples collected to the west of Sicily strait were defined as western Mediterranean (WMS) stations. More details on the physical, chemical and biogeochemical characteristics of the water column during the cruise can be found in Tanhua et al. (2013a, b).

2.2 Experimental design

Subsurface seawater (6–8 m depth) was collected using a low pressure pump and placed in triplicate 4.6 L polycarbonate Nalgene bottles. NaH¹³CO₃ (Sigma) was added to obtain an enrichment of approximately 1% of the ambient dissolved inorganic carbon (460 µL of 200 mmol L⁻¹ NaH¹³CO₃) (Mulholland and Bernhardt, 2005). ¹⁵N₂ uptake measurements were measured using a newly developed ¹⁵N-enriched seawater protocol (Mohr et al., 2010). Enriched seawater was prepared by first degassing filtered (0.2 µm) natural seawater collected at the same site and depth using a

Table 1. Physical and chemical characteristics of the surface seawater (6–8 m) of the MS stations sampled during April 2011. BD- below detection limit; MLD- mixed layer depth.

Station number	290	294	304	312	316	333	338	339
Location	Levantine basin	Shikmona Eddy	Ionian Sea	Adriatic Sea	Tyrrhenian Sea	Alboran Sea	Strait of Gibraltar	Gulf of Cadiz
Position	34°20' N, 27°30' E	34°00' N, 34°25' E	35°36' N, 17°15' E	41°15' N, 18°00' E	38°36' N, 11°30' E	36°06' N, 2°48' E	35°57' N, 4°45' W	35°54' N, 7°00' W
Temperature (°C)	17.0	18.1	17.1	14.7	16.2	16.7	17.8	17.7
Salinity	39.0	39.0	38.3	38.5	37.2	36.3	36.3	36.4
MLD (m)	46	49	30	28	21	45	44	38
NO ₂ +NO ₃ (μM)	0.86 ± 0.05	0.07 ± 0.01	BD	0.39 ± 0.09	0.54 ± 0.16	0.63 ± 0	0.56 ± 0.23	1.39 ± 0.84
PO ₄ (μM)	0.05 ± 0.01	0.05	0.01	0.02 ± 0.02	0.02 ± 0.01	0.24 ± 0.18	0.07 ± 0.02	0.06 ± 0.03
Si(OH) ₄ (μM)	1.10 ± 0.18	0.97 ± 0.06	0.79	0.95 ± 0.17	0.81 ± 0.32	0.61 ± 0.08	0.48 ± 0.13	0.44 ± 0.04

vacuum (250 mbar) applied to continuously stirred seawater for ~ 1 h. The degassed water was transferred into septum capped Nalgene bottles with no headspace, and 1 ml of ¹⁵N₂ gas (99 %) was injected per 100 mL of seawater. The bottles were shaken vigorously until the bubble disappeared. Aliquots of this ¹⁵N₂-sea enriched water were then added to the incubation bottles. The enriched water constituting 5 % of the total sample volume (i.e. 230 mL). Similar enriched seawater additions from the oligotrophic North Pacific Subtropical Gyre (NPSG) resulted in a final ¹⁵N₂ enrichment of 1.5 atom % after adding 50 mL of ¹⁵N₂-enriched water to a 4.5 L bottle (Wilson et al., 2012).

After the enriched seawater and ¹³C were added (i.e. double labeling), the bottles were well shaken, and incubated on deck at ambient surface seawater temperatures, maintained with running surface water pumped on board. Incubations began early in the morning (~ 7 a.m. local time) and the incubators were covered with either neutral density screening to simulate ambient light, or under complete darkness for 48 h incubations. We also compared the obtained rates with 24 h incubations (conducted in parallel) and obtained no significant difference between the rates ($R^2 = 0.91$ $n = 24$ $P < 0.05$, see Supplement Fig. S1). The incubations under ambient irradiance (representative of a full diel cycle) record the activities of both autotrophic and heterotrophic diazotrophs. Whereas, we assume that the 48 h dark incubations reflected the activity of mainly heterotrophic diazotrophs who do not require light energy for dinitrogen fixation. We estimated heterotrophic contribution to N₂ fixation by comparing the dark incubations versus the bottles incubated under ambient diel irradiance.

Incubations were terminated by filtering water onto pre-combusted 25 mm GF/F filters (nominal pore size of 0.7 μm). Filters were then dried in an oven at 60 °C and stored in a dessicator until analysis. In the laboratory, samples for ¹⁵N and ¹³C analyses were pelletized in tin disks and then analyzed on a Europa 20/20 mass spectrometer equipped with an automated nitrogen and carbon analyzer. For isotope ratio mass spectrometry, standard curves to determine N and

C mass were done with each sample run. Samples were run only when standard curves had R_2 values > 0.99. At masses > 4.7 μg N, the precision for the atom percent ¹⁵N measurement was 0.0001 % based on daily calibrations made in association with sample runs and calibrations averaged over runs made over several years. For most of the results reported here, the masses were > 4.7 μg N. However, samples with < 4.7 μg N were only used if the precision was 0.0001 % for that sample run. Standard masses ranged from 1.2 to 100 μg N and from 9.4 to 800 μg C. In addition to daily standard curves, reference standards and standards run as samples were run every six to eight samples.

The percent contribution of N₂ fixation to primary productivity was calculated based on the measured particulate carbon (POC) and nitrogen (PON) in each sample. Although the measured POC and PON are representative of the whole planktonic community and are not specific to diazotrophs, our previous experience in the EMS suggests higher POC : PON ratio than the conventional 106 : 16 Redfield ratio (Yogev et al., 2011, Rahav et al., 2013) and thus were used to calculate the % contribution.

2.3 Physical measurements

Measurements of temperature and salinity were taken at each station along the cruise track using an in situ conductivity, temperature and depth (CTD) sensor (Seabird 19 Plus).

2.4 Inorganic nutrients

Nutrient concentrations were determined for the same seawater used for the N₂ fixation measurements. Duplicate water samples were collected in 15 mL acid-washed plastic scintillation vials from surface (6–8 m) using a low pressure pump (see Sect. 2.1) and immediately frozen at –20 °C. Nutrients were determined in the laboratory ~ 4 months after the cruise using a segmented flow Skalar SANplus System Instrument as detailed in Kress and Herut (2001). The precision of the nitrate + nitrite, orthophosphate and silicic acid measurements were 0.02, 0.003 and 0.06 μM, respectively.

Table 2. Biological characteristics of the surface seawater (6–8 m) of the MS stations sampled during April 2011.

Parameter/station number	290	294	304	312	316	333	338	339
Chlorophyll ($\mu\text{g L}^{-1}$)	0.04 ± 0.01	0.03 ± 0	0.02 ± 0.01	0.11 ± 0.03	0.04 ± 0	0.18 ± 0.01	0.31 ± 0.01	0.07 ± 0.02
<i>Synechococcus</i> (cell L^{-1})	1.33×10^7	2.26×10^6	3.86×10^6	1.78×10^7	1.16×10^7	2.68×10^7	3.27×10^7	4.94×10^6
<i>Prochlorococcus</i> (cell L^{-1})	1.17×10^6	8.32×10^4	3.17×10^5	1.14×10^6	1.24×10^6	2.60×10^6	1.60×10^6	3.57×10^6
picoeukaryotes (cell L^{-1})	4.36×10^5	2.08×10^4	7.53×10^4	2.23×10^5	7.35×10^5	2.53×10^6	3.69×10^6	1.46×10^6
<i>Synechococcus</i> (ng CL^{-1})	2328	396	676	3115	2030	4690	5723	865
<i>Prochlorococcus</i> (ng CL^{-1})	62	4	17	60	66	138	85	19
pico-eukaryotes (ng CL^{-1})	916	44	158	468	1544	5313	7749	3066
POC : PON	9.3 ± 2.5	9.2 ± 0.8	8.3 ± 0.7	7.6 ± 0.7	7.4 ± 0.5	8.2 ± 1.7	8.6 ± 1.6	6.4 ± 0.3
Primary productivity ($\mu\text{g CL}^{-1} \text{d}^{-1}$)	0.74 ± 0.01	0.53 ± 0.02	0.21 ± 0.01	1.39 ± 0.87	0.76 ± 0.13	0.78 ± 0.26	15.04 ± 1.61	8.01 ± 1.79
N_2 fixation ($\text{nmol NL}^{-1} \text{d}^{-1}$)	0.15 ± 0.01	0.12 ± 0.02	0.10 ± 0.02	0.29 ± 0.02	0.22 ± 0.03	0.86 ± 0.17	2.35 ± 1.12	0.39 ± 0.14

The limits of quantification were $0.075 \mu\text{M}$, $0.008 \mu\text{M}$ and $0.07 \mu\text{M}$ for nitrate+ nitrite, orthophosphate and silicic acid, respectively (note that for silicic acid the limit of quantification is similar to the precision). For full nutrients profiles of samples collected from Niskin bottles see Tanhua et al. (2013b).

2.5 Chlorophyll *a* extraction

Duplicate seawater samples (500 mL) taken twice a day across the MS ($n = 94$) were filtered onto glass fiber filters. The filters were stored at -20°C in a dark box until analysis within 2–3 days. Samples were extracted in 5 mL 90 % acetone overnight, at 4°C in dark. Chlorophyll *a* (Chl *a*) concentrations were determined with a Turner Designs (TD-700) fluorometer, using a 436 nm excitation filter and a 680 nm emission filter (Holm-Hansen, 1965). A blank filter was also stored in 90 % acetone under the same conditions as the samples.

2.6 Picophytoplankton abundance

The abundance of picophytoplankton was determined by flow cytometry. Taxonomic discrimination was based on the following parameters: cell side scatter – a proxy of cell volume; forward scatter – a proxy of cell size; and orange and red fluorescence of phycoerythrin and of chlorophyll *a* (585 nm and 630 nm, respectively). Samples of 1.8 mL were fixed immediately at room temperature with $23 \mu\text{L}$ of 25 % glutaraldehyde (Sigma G-5882) retained at room temperature for 10 min, subsequently frozen in liquid nitrogen, and kept

at -80°C until analyzed. Samples were fast thawed at 37°C , and counted using a FACScan Becton Dickinson flow cytometer, fitted with an Argon laser (488 nm) for 10 to 15 min or until 30 000 cells were counted (Vaulot et al., 1989). Pico/nano phytoplankton carbon (C) biomass was calculated from cell counts assuming $175 \text{ fg C cell}^{-1}$ for *Synechococcus* cells $53 \text{ fg C cell}^{-1}$ for *Prochlorococcus* cells, and $2100 \text{ fg C cell}^{-1}$ for pico-eukaryotes (Campbell and Yentsch, 1989).

3 Results

3.1 East–west distribution of physical, chemical and phytoplankton parameters

The physical, chemical and biological parameters of the surface waters at each station are provided in Tables 1 and 2. Overall, surface temperatures and salinities increased from west to east from 14.7 to 18.1°C and 36.3 to 39 , respectively. $\text{NO}_2^- + \text{NO}_3^-$ (DIN) increased from east to west from below detection in the Ionian Sea to $1.39 \mu\text{M}$ at the Gulf of Cadiz station (Table 1). In contrast, Station 290 (NW Levantine Basin) had high surface concentrations of DIN ($0.86 \mu\text{M}$), probably due to upwelling of deeper waters within the cyclonic Rhodes Gyre. Dissolved inorganic phosphorus (DIP) ranged from 0.01 to $0.24 \mu\text{M}$ in surface waters across the entire Mediterranean Sea (MS) (Table 1). Silicic acid ($\text{Si}(\text{OH})_4$) concentration was lowest in the westernmost stations – at the entrance to the MS ($0.44 \mu\text{M}$), and increased toward the east with highest concentration observed at the easternmost station ($1.10 \mu\text{M}$) (Table 1).




Fig. 2. Picophytoplankton distribution of *Synechococcus* (A), *Prochlorococcus* (B) and pico-eukaryotes (C) in the surface waters (6–8 m) of the eastern (black circle) and western (white circle) Mediterranean Sea. $n = 21$ and $n = 12$ for the eastern and western basins, respectively.

Chlorophyll (Chl *a*) concentrations increased from east to west across the MS. Surface Chl *a* concentrations were $\sim 0.03 \mu\text{g L}^{-1}$ at the eastern basin stations and up to $0.31 \mu\text{g L}^{-1}$ at the Strait of Gibraltar – the westernmost station (Fig. 1). *Synechococcus* dominated the picophytoplankton ranging from as low as 2.26×10^6 cells L^{-1} to 3.27×10^7 cells L^{-1} in the eastern and western basin, respectively (Fig. 2, Table 2). Using a cell : carbon conversion ratio of $175 \text{ fg C cell}^{-1}$ (see methods), this represents a range of 396 ng CL^{-1} to 5723 ng CL^{-1} . In the eastern basin, the picoeukaryote abundances ($\sim 2.1 \times 10^4$ to 7.5×10^4 cell L^{-1}) and biomass (44 to 158 ng CL^{-1}) were low except in the Levantine basin (Station 290) where higher abundances (4.36×10^5 cell L^{-1}) and biomass (916 ng CL^{-1}) were measured (Fig. 2, Table 2). *Prochlorococcus* abundances and biomass from the surface waters were generally low throughout the whole MS, especially at the Shikmona Eddy (Station 294) and the Ionian Sea (station 304) (Fig. 2, Table 2).

3.2 Primary productivity and N_2 fixation rates

Photosynthetic carbon fixation rates ranged from 0.21 to $0.74 \mu\text{g CL}^{-1} \text{ d}^{-1}$ in the eastern basin, and 0.76 to $1.39 \mu\text{g CL}^{-1} \text{ d}^{-1}$ at the western Mediterranean stations. Much higher rates were measured at the Strait of Gibraltar ($15.04 \pm 1.6 \mu\text{g CL}^{-1} \text{ d}^{-1}$) and in the Gulf of Cadiz ($8.22 \mu\text{g CL}^{-1} \text{ d}^{-1}$) (Table 2).

N_2 fixation rates obtained across the MS exhibited a strong zonal gradient from the eastern to western basins (Fig. 3a and Table 2). The lowest N_2 fixation rates were measured in the eastern basin, ranging from $0.10 \pm 0.02 \text{ nmol NL}^{-1} \text{ d}^{-1}$ in the Ionian Sea, to $0.15 \pm 0.01 \text{ nmol NL}^{-1} \text{ d}^{-1}$ at Station 290 (affected by the Rhodes Gyre) (Fig. 3a and Table 2). N_2 fixation rates increased gradually toward the west ranging from 0.22 ± 0.03 in the Tyrranean Sea to $2.35 \pm 1.12 \text{ nmol NL}^{-1} \text{ d}^{-1}$ at the westernmost station at the Strait of Gibraltar (Fig. 3a and Table 2). The springtime rates of N_2 fixation at all stations were 3–7 fold higher than measurements published previously during summertime (Fig. 3b).

In addition to total N_2 fixation (measured in light bottles under ambient diel irradiance), we examined N_2 fixation rates in bottles incubated for 48 h in the dark. While some unicellular cyanobacteria fix N_2 during the dark hours, they require light energy to fuel the process. We assumed that after 48 h in the dark, the contribution by these diazotrophs will be negligible and most N_2 fixation would be due to heterotrophic diazotrophs that do not require light for the N_2 fixing process (Postage, 1979). The N_2 fixation rates from 48 h dark incubations showed a similar east-west trend as observed in light bottle incubations (Fig. 4a); within the eastern basin, N_2 fixation in dark incubations were lowest at the easternmost Station 290 ($0.11 \pm 0.02 \text{ nmol NL}^{-1} \text{ d}^{-1}$) and highest at Station 294 in the Shikmona Eddy ($0.16 \pm 0.01 \text{ nmol NL}^{-1} \text{ d}^{-1}$) (Figure 4A). In the western basin N_2 fixation rates in dark incubation bottles rates ranged from 0.20 ± 0.05 to $0.40 \pm 0.11 \text{ nmol NL}^{-1} \text{ d}^{-1}$ (Fig. 4a).

We compared rates of light and dark N_2 fixation (Fig. 4b) to estimate the relative contribution of autotrophic versus heterotrophic N_2 fixation. In the western basin, light : dark estimates of N_2 fixation were always > 1 , suggesting the predominance of autotrophic N_2 fixation. In the eastern basin light : dark N_2 fixation rates were < 1 suggesting a preponderance of heterotrophic diazotrophs (Fig. 4).

3.3 The contribution of N_2 fixation to primary productivity

We calculated the percent contribution of N_2 fixation to total primary productivity during springtime based on rates of N_2 fixation measured in the light bottle incubations and the associated C fixation estimated using an the average particulate C : N ratio obtained at each station (Table 2, and see Yogeve et al., 2011; Rahav et al., 2013). New production due to N_2 fixation was $\sim 2\%$ of the total primary productivity at the EMS stations and increased by a factor of 2 to 4 in the western Mediterranean Sea (WMS), ranging from 3.5% in the Adriatic Sea to 8.5% in the Alboran Sea. The percent contribution of N_2 fixation to primary production in the Gulf of




Fig. 3. Seasonal variations of N_2 fixation in the surface waters of the Mediterranean Sea. (A) Springtime rates measured in this study. (B) Summer data compiled from Rahav et al., 2013; Yogeve et al., 2011; Ibello et al., 2010; Bonnet et al., 2011.

Cadiz, near the Strait of Gibraltar that connects the Mediterranean Sea with the Atlantic Ocean, was 2.3 % (Fig. 5).

4 Discussion

This study provides the first springtime measurements of N_2 fixation in surface waters along an east–west transect across the Mediterranean Sea (MS). We focused sampling at representative stations from different water provinces in the MS (Fig. 1, Table 1). Our results yielded N_2 fixation rates in surface waters that are 3–7 fold higher (Fig. 3a, Table 2) than published rates from two summertime basin-wide N_2 fixation studies (Ibello et al., 2010; Bonnet et al., 2011), routine stations off the Israeli coast (Yogeve et al., 2011), and a Levantine Basin transect (Rahav et al., 2013). Moreover, the gradient of increasing N_2 fixation rates from east to west coincide with the east–west gradient in surface Chl *a* (Fig. 1) and primary productivity (Table 2).

Seasonal measurements of N_2 fixation rates in the MS have been made at two monitoring stations, one located west of the Israeli coastline (Levantine Basin) (Yogeve et al., 2011) and the other off the coast of France, the DYFAMED station (Ligurian Sea) (Garcia et al., 2006; Sandroni et al., 2007). Rates of N_2 fixation in surface waters from the Levantine Basin were uniformly low (~ 0.01 nmol $N L^{-1} d^{-1}$) and did not show any seasonality (Yogeve et al., 2011). In contrast, at the WMS time series station (DYFAMED), higher rates of N_2 fixation were measured during April and August ($4\text{--}7.5$ nmol $N L^{-1} d^{-1}$, 10 m) relative to other months

(< 2 nmol $N L^{-1} d^{-1}$, 10 m), which were associated with higher primary productivity rates (Sandroni et al., 2007).

The Shikmona Eddy (Station 294) and the Ionian Sea (Station 304), representing ultraoligotrophic conditions, had lower nutrient and Chl *a* concentrations than the more productive cyclonic Rhodes Gyre station (Station 290). Yet similar N_2 fixation rates were measured at all three stations (Fig. 3a, Table 2) and there was no correlation between N_2 fixation and primary production ($R^2 = 0.18$, $n = 9$, t test, $P > 0.05$). This suggests that N_2 fixation is attributed mainly to heterotrophic bacteria or that diazotrophs and nondiazotrophic phytoplankton are limited or co-limited by different nutrients. Heterotrophic bacteria are known to compete for N with autotrophs in the nutrient-depleted surface waters of the EMS (Thingstad et al., 2005; Tanaka et al., 2007) and molecular fingerprinting suggests a highly diverse heterotrophic community of *nifH* phylotypes (Man-Aharonovich et al., 2007; Yogeve et al., 2011). Heterotrophic diazotrophs may outcompete other bacteria in an N-impoverished system because they can acquire N from the abundant N_2 pool. Evidence for heterotrophic diazotrophy was found in both surface and aphotic depths in the EMS (Rahav et al., 2013).

Higher DIN (Table 1) and Chl *a* concentrations were measured in the more productive WMS compared to the EMS (Fig. 1, Table 2). Concurrently, N_2 fixation rates in the WMS were also higher (ANOVA, $P < 0.05$) ranging from 0.22 to 0.86 nmol $N L^{-1} d^{-1}$ (Fig. 3a, Table 2) and correlated with PP ($R^2 = 0.82$, $n = 12$, t test, $P < 0.05$), suggesting photoautotrophic associated N_2 fixation. Indeed, relatively high diatom abundances were detected in surface waters of the WMS (> 100 cells L^{-1}) associated with a small spring bloom (A. Oviedo, personal communication, 2013). *Richelia intracellularis*, a symbiotic N_2 fixing cyanobacterium, has been found associated with diatoms in the EMS previously (Bar-Zeev et al., 2008) and may have contributed to N_2 fixation in the WMS.

The highest N_2 fixation rates during this spring transect were observed at the westernmost station in the Strait of Gibraltar (Fig. 3a, Table 2). Moreover, these springtime N_2 fixation rates were 7-fold higher than those measured previously during summer by Ibello et al. (2010) (2.35 nmol $N L^{-1} d^{-1}$ vs. 0.3 nmol $N L^{-1} d^{-1}$, respectively). These differences suggest seasonality of N_2 fixation and/or the abundance or activity of diazotrophic populations, or seasonal exchange of water and resident planktonic populations between the eastern Atlantic Ocean and the MS through the Strait of Gibraltar.

During this study N_2 fixation rates were only measured in surface waters (upper 6–8 m) and therefore depth-integrated N_2 fixation rates could not be calculated. It is therefore conceivable that many autotrophic and heterotrophic diazotrophic groups populating other depths, such as the deep Chl *a* maximum (DCM), were not accounted for in our rate measurements. In addition, seasonal changes in the vertical




Fig. 4. (A) N_2 fixation rates of surface waters from stations across the Mediterranean Sea for bottles incubated under ambient lighting (white bars) and in complete darkness (dark bars). The asterisk above the columns represents statistically significant differences (one-way ANOVA, $P < 0.05$) for mean values of N_2 fixation rates in each station, and (B) the resulting ratio between rates of N_2 fixation from ambient lighting and dark incubations. $n = 3$ for each incubation type at each station.




Fig. 5. The percent contribution of N_2 fixation to primary productivity (PP) of surface waters sampled across the MS during the spring period. The letters above the columns represent statistically significant differences (one-way ANOVA and a Fisher LSD means comparison test, $P < 0.05$) for mean values of % contribution between stations.

distribution of diazotrophic microbes were not considered here. For example, a recent study from the eastern basin found no statistical difference in N_2 fixation rates measured in water collected from below the pycnocline at the DCM compared to surface waters during the stratified period, while during the winter mixing period, when the water column was mixed up to 150 m, the N_2 fixation rates were 2–3 fold higher at the DCM than in surface waters (Yogev et al., 2011).

Another methodological contribution to the higher N_2 fixation rates during spring throughout the MS was our use of the newly enriched ($^{15}N_2$) seawater addition method (Mohr

et al., 2010) rather than the gas bubble $^{15}N_2$ addition method (Montoya et al., 1996). The gas bubble enrichment method may underestimate N_2 fixation rates by a factor of 2 or more in some circumstances (Großkopf et al., 2012; Wilson et al., 2012). Our preliminary comparison of both methods in MS waters demonstrated a 2–3 fold increase in rates using the enriched seawater method ($n = 18$). However, in long incubations (> 24 h), the underestimate of N_2 fixation using the bubble method was reduced because the gas bubble should have equilibrated within the first several hours of the incubation (Mohr et al., 2010; Mulholland et al., 2012). While it is impossible to convert from one method to another using a constant conversion factor, if we assume a two-fold underestimate of previously reported summer N_2 fixation rates, we still observe significant seasonal differences in N_2 fixation rates between the early spring and fully stratified summer periods. This suggests that methodological differences alone cannot account for the seasonal changes we observed.

We examined the relative contribution of autotrophic and heterotrophic diazotrophs to the measured N_2 fixation rates using parallel natural light and dark bottle incubations. It has generally been assumed that diazotrophy in surface waters is dominated by photoautotrophic cyanobacteria that use light energy to satisfy the energetic demands of N_2 fixation and to acquire carbon (Karl et al., 2002). Yet, research demonstrates that the abundant and widely distributed unicellular group A cyanobacteria are photoheterotrophs (Zehr et al., 2008; Tripp et al., 2010). Further, many heterotrophic diazotrophs are present in surface waters (Riemann et al., 2010; Zehr and Kudela, 2011; Mulholland et al., 2012). Our results show that in the eastern basin stations, the ratio of light : dark bottle N_2 fixation was usually < 1 (Fig. 4b) suggesting that heterotrophic diazotrophs may be the dominant N_2 fixers, although we cannot exclude that some of the

dark N_2 fixation was performed by unicellular cyanobacteria. In the western basin, this ratio was generally > 1 suggesting that autotrophic diazotrophs predominated (Fig. 4b). We acknowledge that some phototrophic diazotrophs fix N_2 during the dark, to avoid the inhibitory effects of oxygen, but we assume that our long incubation time in the dark (48 h) would have diminished their impact as they require light energy to fix N_2 .

Phylogenetic characterizations of diazotrophs in surface waters across this Mediterranean transect are currently unavailable. However, a diverse group of auto- and heterotrophic diazotrophs have been reported from the eastern basin with $\sim 40\%$ of the *nifH* transcripts attributed to heterotrophic bacteria (Man-Aharonovich et al., 2007; Bar-Zeev et al., 2008; Yogev et al., 2011). In the WMS, unicellular cyanobacteria (including UCYN-A) are present in low abundances year round and short blooms of 2000–5000 cells mL^{-1} have been reported from a coastal station off Marseille during June and July (Le Moal and Biegala, 2009). Another recent study suggested that cells $< 0.7 \mu m$ in size, usually ignored during routine sampling, can contribute 50% of the N_2 fixation (Konno et al., 2010). In this study we used GF/F filters to measure planktonic N_2 fixation (nominal pore size of $\sim 0.7 \mu m$, see methods), as is a common practice. Thus, it is possible we could have missed N_2 fixation by very small bacteria diazotrophs and thereby underestimated total planktonic N_2 fixation.

Based on results from studies conducted during summer in the EMS, N_2 fixation accounted for only 0.7–2% of primary productivity at stations in the Levantine basin (Yogev et al., 2011, Rahav et al., 2013), but increased to $\sim 6\%$ in the more productive Rhodes Gyre and Cyprus Eddy (Rahav et al., 2013). Consistent with these results, during a summer transect across the Mediterranean (BOUM campaign), N_2 fixation accounted for 6 to 35% of new production at stations in the more productive western basin but only 0 to 0.3% at the more oligotrophic eastern basin (Bonnet et al., 2011). Our springtime results show higher N_2 fixation rates (2–4 fold) at both basins and a similar spatial trend. A higher contribution of N_2 fixation to primary production (4–8%) was measured in the western basin compared to the eastern basin ($\sim 2\%$, Fig. 5). These differences between the two basins are probably attributed to changes in N_2 -fixing planktonic communities and other environmental aspects. Summertime data from the EMS demonstrated a significant positive correlation between N_2 fixation rates and bacterial production suggesting a higher involvement of heterotrophic diazotrophs in the ultraoligotrophic EMS (Rahav et al., 2013).

5 Conclusions

This study provides the first direct measurements of N_2 fixation rates in surface waters across the MS during springtime. N_2 fixation rates were measured using the newly modified

$^{15}N_2$ -uptake method (Mohr et al., 2010) during a spring transect and were 3–7 fold higher than measurements made in surface waters during the stratified summer period. Methodological differences cannot fully account for the higher rates of N_2 fixation observed during this cruise and we suggest that the higher rates are due to seasonal variability in primary productivity and environmental factors. N_2 fixation was higher and contributed more to total primary production in the western basin than in the eastern basin. While our data suggests that N_2 fixation rates across the MS are higher during spring than in the summer stratified period, our measurements were constrained to surface waters and thus we cannot provide depth integrated estimates of N_2 fixation during spring. We suggest that future investigations should include N_2 fixation rate measurements and phylogenetic identity of diazotrophs at both photic and aphotic depths to better constrain the contribution of N_2 fixation to N budgets as well as the total and new production within the Mediterranean Sea.

Supplementary material related to this article is available online at: <http://www.ocean-sci.net/9/489/2013/os-9-489-2013-supplement.pdf>.

Acknowledgements. Many thanks for the help provided by the captain and crew of the R/V *Meteor*. We would also like to thank Toste Tanhua, the Chief Scientist of this cruise, for allowing us to take part in this campaign. This research was supported by the Israel Science Foundation grant (996/08) to I. B-F and B. H. This study is in partial fulfillment of a Ph.D. thesis for Eyal Rahav from Bar Ilan University.

Edited by: T. Tanhua

References

- Bar Zeev, E., Yogev, T., Man-Aharonovich, D., Kress, N., Herut, B., Beja, O., and Berman-Frank, I.: Seasonal dynamics of the endosymbiotic, nitrogen-fixing cyanobacterium *Richelia intracellularis* in the Eastern Mediterranean Sea, *ISME J.*, 2, 911–92. 2008.
- Berman, T., Townsend, D. W., Elsayed, S. Z., Trees, C. C., and Azov, Y.: Optical transparency, chlorophyll and primary productivity in the Eastern Mediterranean near the Israeli coast, *Oceanol. Acta.*, 7, 367–372, 1984.
- Bonnet, S., Grosso, O., and Moutin, T.: Planktonic dinitrogen fixation along a longitudinal gradient across the Mediterranean Sea during the stratified period (BOUM cruise), *Biogeosciences*, 8, 2257–2267, doi:10.5194/bg-8-2257-2011, 2011.
- Bosc, E., Bricaud, A., and Antoine, D.: Seasonal and inter-annual variability in algal biomass and primary production in the Mediterranean Sea, as derived from 4 years of SeaWiFS observations, *Global Biogeochem. Cy.*, 18, GB1005, doi:10.1029/2003GB002034, 2004.

- Campbell, J. W. and Yentsch, C. M.: Variance within homogeneous phytoplankton populations .1. Theoretical framework for interpreting histograms, *Cytometry*, 10, 587–595, 1989.
- Capone, D. G. and Montoya, J. P.: Nitrogen fixation and denitrification, *Methods Microbiol.*, 30, 501–51, 2001.
- Falkowski, P. G., Barber, R. T., and Smetacek, V.: Biogeochemical controls and feedbacks on ocean primary production, *Science*, 281, 200–206, 1998.
- Garcia, N., Raimbault, P., Gouze E., and Sandroni, V.: Nitrogen fixation and primary production in Western Mediterranean, *Comptes. Rendus. Biologies.*, 329, 742–750, 2006.
- Großkopf, T., Mohr, W., Baustian, T., Schunck, H., Gill, D., Kuypers, M. M. M., Lavik, G., Schmitz, R. A., Wallace, D. W. R., and LaRoche, J.: Doubling of marine dinitrogen-fixation rates based on direct measurements, *Nature*, 488, 361–364, 2012.
- Gruber, N.: The marine nitrogen cycle: Overview of distributions and processes, in: *Nitrogen in the marine environment*, edited by: Capone, D., Bronk, D., Mulholland, M., and Carpenter, E., Elsevier, Amsterdam, 1–50, 2008.
- Holm-Hansen, O., Lorenzen, C. J., Homes, R. W., and Strickland, J. D. H.: Fluorometric determination of chlorophyll., *Journal du Conseil Permanent Internationale pour L'Exploration de la Mer.*, 30, 2–15, 1965.
- Ibello, V., Cantoni, C., Cozzi, S., and Civitarese, G.: First basin-wide experimental results on N₂ fixation in the open Mediterranean Sea, *Geophys. Res. Lett.*, 37, L03608, doi:10.1029/2009GL041635, 2010.
- Ignatiades, L., Gotsis-Skretas, O., Pagou, K., and Krasakopoulou, E.: Diversification of phytoplankton community structure and related parameters along a large-scale longitudinal east-west transect of the Mediterranean Sea, *J. Plankton. Res.*, 31, 411–428, 2009.
- Karl, D., Michaels, A., Bergman, B., Capone D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., and Stal, L.: Dinitrogen fixation in the world's oceans, *Biogeochemistry*, 57, 47–98, 2002.
- Konno, U., Tsunogai, U., Komatsu, D. D., Daita, S., Nakagawa, F., Tsuda, A., Matsui, T., Eum, Y.-J., and Suzuki, K.: Determination of total N₂ fixation rates in the ocean taking into account both the particulate and filtrate fractions, *Biogeosciences*, 7, 2369–2377, doi:10.5194/bg-7-2369-2010, 2010.
- Kress, N. and Herut, B.: Spatial and seasonal evolution of dissolved oxygen and nutrients in the Southern Levantine Basin (Eastern Mediterranean Sea): chemical characterization of the water masses and inferences on the N:P ratios, *Deep-Sea Res. Pt. I*, 48, 2347–2372, 2001.
- Krom, M. D., Brenner, S., Kress, N., and Gordon, L. I.: Phosphorous limitation of primary productivity in the Eastern Mediterranean Sea, *Limnol. Oceanogr.*, 36, 424–43, 1991.
- Krom, M. D., Emeis, K. C., and Van Cappellen, P.: Why is the Eastern Mediterranean phosphorus limited?, *Prog. Oceanogr.*, 85, 236–244, 2010.
- Le Moal, M., and Biegala, I. C.: Diazotrophic unicellular cyanobacteria in the Northwestern Mediterranean Sea: A seasonal cycle, *Limnol. Oceanogr.*, 54, 845–855, 2009.
- Le Moal, M., Collin, H., and Biegala, I. C.: Intriguing diversity among diazotrophic picoplankton along a Mediterranean transect: a dominance of rhizobia, *Biogeosciences*, 8, 827–840, doi:10.5194/bg-8-827-2011, 2011.
- Man-Aharonovich, D., Kress, N., Bar Zeev, E., Berman-Frank, I., and Beja, O.: Molecular ecology of *nifH* genes and transcripts in the Eastern Mediterranean Sea, *Environ. Microbiol.*, 9, 2354–236, 2007.
- Moisander, P. H., Beinart, R. A., Hewson, I., White, A. E., Johnson, K. S., Carlson, C. A., Montoya, J. P., and Zehr, J. P.: Unicellular cyanobacterial distributions broaden the oceanic N₂ fixation domain, *Science*, 327, 1512–1514, 2010.
- Mohr, W., Grosskopf, T., Wallace, D. W. R., and LaRoche, J.: Methodological underestimation of oceanic nitrogen fixation rates, *PLoS One*, 5, e12583. doi:10.1371/journal.pone.0012583, 2010.
- Montoya, J. P., Voss, M., Kahler, P., and Capone, D. G.: A simple, high-precision, high-sensitivity tracer assay for N₂ fixation, *Appl. Environ. Microbiol.*, 62, 986–993, 1996.
- Moutin, T. and Raimbault, P.: Primary production, carbon export and nutrients availability in western and eastern Mediterranean Sea in early summer 1996 (MINOS cruise), *J. Marine. Syst.*, 33, 273–288, 2002.
- Mulholland, M. R., Boneillo, G., and Minor, E. C.: A comparison of N and C uptake during brown tide (*Aureococcus anophagefferens*) blooms from two coastal bays on the east coast of the USA, *Harmful Algae*, 3, 361–376, 2004.
- Mulholland, M. R. and Capone, D. G.: Dinitrogen fixation in the Indian Ocean, in: *Indian Ocean Biogeochemical Processes and Ecological Variability*, edited by: Wiggert, J. D., Hood, R. R., Naqvi, S. W. A., Brink, K. H., and Smith, S. L., American Geophysical Union, Washington, DC, 167–186, 2009.
- Mulholland, M. R. and Bernhardt, P. W.: The effect of growth rate, phosphorus concentration, and temperature on N₂ fixation, carbon fixation, and nitrogen release in continuous cultures of *Trichodesmium* IMS101, *Limnol. Oceanogr.*, 50, 839–849, 2005.
- Postgate, J.: Nitrogen-fixation and the future of the worlds food-supply, *Biologist*, 26, 165–166, 1979.
- Rahav, E., Herut, B., Stambler, N., Bar Zeev, E., Mulholland, M. R., and Berman-Frank, I.: Uncoupling between dinitrogen fixation and primary productivity in the Eastern Mediterranean Sea, *J. Geophys. Res-Bioge.*, 118, 1–8, doi:10.1002/jgrg.20023, 2013.
- Ridame, C., Le Moal, M., Guieu, C., Ternon, E., Biegala, I. C., L'Helguen, S., and Pujo-Pay, M.: Nutrient control of N₂ fixation in the oligotrophic Mediterranean Sea and the impact of Saharan dust events, *Biogeosciences*, 8, 2773–2783, doi:10.5194/bg-8-2773-2011, 2011.
- Riemann, L., Farnelid, H., and Steward, G. F.: Nitrogenase genes in non-cyanobacterial plankton: prevalence, diversity and regulation in marine waters, *Aquat. Microb. Ecol.*, 61, 225–237, 2010.
- Sandroni, V., Raimbault, P., Migon, C., Garcia, N., and Gouze, E.: Dry atmospheric deposition and diazotrophy as sources of new nitrogen to northwestern Mediterranean oligotrophic surface waters, *Deep-Sea Res. Pt. I*, 54, 1859–1870, 2007.
- Siokou-Frangou, I., Christaki, U., Mazzocchi, M. G., Montresor, M., Ribera d'Alcalá, M., Vaqué, D., and Zingone, A.: Plankton in the open Mediterranean Sea: a review, *Biogeosciences*, 7, 1543–1586, doi:10.5194/bg-7-1543-2010, 2010.
- Spatharis, S., Skliris, N., Meziti, A., and Kormas, K.: First record of a *Trichodesmium erythraeum* bloom in the Mediterranean Sea: a result of climate change?, *Can. J. Fish. Aquat. Sci.*, 69, 1444–1455, 2012.

- Tanaka, T., Zohary, T., Krom, M. D., Lawe, C. S., Pitta, P., and Psarra, S.: Microbial community structure and function in the Levantine Basin of the Eastern Mediterranean, *Deep-Sea Res. I*, 54, 1721–1743, 2007.
- Tanaka, T., Thingstad, T. F., Christaki, U., Colombet, J., Cornet-Barthaux, V., Courties, C., Grattepanche, J.-D., Lagaria, A., Nedoma, J., Oriol, L., Psarra, S., Pujo-Pay, M., and Van Wambeke, F.: Lack of P-limitation of phytoplankton and heterotrophic prokaryotes in surface waters of three anticyclonic eddies in the stratified Mediterranean Sea, *Biogeosciences*, 8, 525–538, doi:10.5194/bg-8-525-2011, 2011.
- Tanhua, T., Hainbucher, D., Cardin, V., Álvarez, M., and Civitarese, G.: Repeat hydrography in the Mediterranean Sea, data from the Meteor cruise 84/3 in 2011, *Earth Syst. Sci. Data Discuss.*, 6, 59–71, doi:10.5194/essdd-6-59-2013, 2013.
- Tanhua, T., Hainbucher, D., Schröder, K., Cardin, V., Álvarez, M., and Civitarese, G.: The Mediterranean Sea system: a review and an introduction to the special issue, *Ocean Sci. Discuss.*, 10, 581–617, doi:10.5194/osd-10-581-2013, 2013.
- Thingstad, F. T., Zweifel, U. L., and Rassoulzadegan, F.: P limitation of heterotrophic bacteria and phytoplankton in the northwest Mediterranean, *Limnol. Oceanogr.*, 43, 88–94, 1998.
- Thingstad, T. F., Krom, M. D., Mantoura, R. F., Flaten, G. A., Groom, S., and Herut, B.: Nature of phosphorus limitation in the ultraoligotrophic Eastern Mediterranean, *Science*, 309, 1068–1071, 2005.
- Tripp, H. J., Bench, S. R., Turk, K. A., Foster, R. A., Desany, B. A., Niazi, F., and Zehr, J. P.: Metabolic streamlining in an open-ocean nitrogen-fixing cyanobacterium, *Nature*, 464, 90–94, 2010.
- Vaulot, D., Courties, C., and Partensky, F.: A simple method to preserve oceanic phytoplankton for flow cytometric analyses, *Cytometry*, 10, 629–635, 1989.
- Wilson, S. T., Bottjer, D., Church, M. J., and Karl, D. M.: Comparative assessment of nitrogen fixation methodologies, conducted in the oligotrophic North Pacific Ocean, *Appl. Environ. Microb.*, 78, 6516–6523, 2012.
- Yogev, T., Rahav, E., Bar-Zeev, E., Man-Aharonovich, D., Stambler, N., Kress, N., Beja, O., Mulholland, M. R., Herut, B., and Berman-Frank, I.: Is dinitrogen fixation significant in the Levantine Basin, East Mediterranean Sea?, *Environ. Microbiol.*, 13, 854–87, 2011.
- Zehr, J. P. and Kudela, R. M.: Nitrogen cycle of the open ocean: from genes to ecosystems, in: *Annual Review of Marine Science*, Vol. 3, edited by: Carlson, C. A. and Giovannoni, S. J., 197–225, 2011.
- Zehr, J. P. and Ward, B. B.: Nitrogen cycling in the ocean: New perspectives on processes and paradigms, *Appl. Environ. Microb.*, 68, 1015–1024, 2002.
- Zehr, J. P., Bench, S. R., Carter, B. J., Hewson, I., Niazi, F., Shi, T., and Affourtit, J. P.: Globally Distributed Uncultivated Oceanic N-2-Fixing Cyanobacteria Lack Oxygenic Photosystem II, *Science*, 322, 1110–1112, doi:10.1126/science.1165340, 2008.