

BUILDING AN UPPER TRIASSIC CARBON ISOTOPE REFERENCE CURVE

Sylvain RICHZOZ¹, Leopold KRISTYN¹, Christoph SPÖTL² &
Wolfram M. KÜRSCHNER³

¹ Department of Palaeontology, University of Vienna, Althanstrasse 14, A-1090 Vienna.
sylvain.richoz@univie.ac.at, leopold.krystyn@univie.ac.at

² Institute of Geology & Paleontology, Leopold Franzens University of Innsbruck, Innrain 52, A-6020 Innsbruck.
christoph.spoetl@uibk.ac.at

³ Inst. of Environ. Biology, Laboratory of Palaeobotany & Palynology, Utrecht University, NL-3584 CD Utrecht.
w.m.kuerschner@uu.nl

During the Upper Triassic, it appears that despite new originations, the general decline in biodiversity was punctuated by a series of accelerated steps between the Carnian and the Rhaetian, while the T-J boundary event may have been the final strike (McROBERTS & NEWTON, 1995, HALLAM, 2002). How these changes in the biosphere were related to oceanographic and/or geochemical changes during the Late Triassic biotic crisis events is a question of primary importance. In order to solve this issue, two principal questions have to be addressed: (1) are these extinctions best explained by a gradual process of environmental change or by (a series of) abrupt or even catastrophic events? and (2) how do Late Triassic patterns of biotic turnover correlate and couple with oceanographic geochemistry? An expansion of a well-calibrated carbon isotope reference curve for the whole Upper Triassic is an important first step to address these questions.

While a comprehensive isotopic data set is available for the T-J boundary (e.g. MORANTE & HALLAM, 1996; GUÉX et al., 2004; KÜRSCHNER et al., 2007; WILLIFORD et al., 2007) only a few data are available for the Upper Triassic. To establish a carbon isotope reference curve and in addition of the sparse literature (ATUDOREI, 1999; GAWLICK & BÖHM, 2000; HAUSER et al., 2001; MUTTONI et al. 2004; HORNUNG & BRANDNER 2005; KORTE et al., 2005; HORNUNG et al., 2007a,b, WARD et al., 2004; SEPHTON et al., 2002), several Tethyan and Peritethyan sections were measured in the Austrian Alps, Slovakia, Turkey, Oman and the Indian.

The Upper Ladinian samples record an increase in $d^{13}C_{carb}$ until the Lower Carnian, followed by stable values until the Upper Carnian (MIETTO et al., 2007; RICHZOZ et al., 2007a). This stability is, however, disturbed by some small negative excursions in the isotopic signal near the Reingraben event (Lower Carnian-Upper Carnian boundary, ATUDOREI, 1999; HAUSER et al., 2001; HORNUNG & BRANDNER, 2005 and HORNUNG et al., 2007a,b). We present here evidence from the Spiti valley, Indian Himalaya; Mayerling, Austrian Alps and several sections in Taurus. The Carnian-Norian boundary interval in Turkey and Slovakia is marked by a minor increase in the C isotope value (less than 1‰, MUTTONI et al., 2004; GAWLICK & BÖHM, 2000; RICHZOZ et al., 2007b). The isotopic values then show an increase until the

Middle Norian followed by a decrease recorded in Oman, Sicily and Austria (MUTTONI et al., 2004; RICHOZ et al., 2007b). In the Upper Norian the isotopic values are relatively stable, and show no shift across the newly proposed Norian/Rhaetian boundary (Steinbergkogel, Austria and Oman, KRYSSTYN et al., 2007) before increasing again through the classical Norian/Rhaetian boundary (Oman, Turkey, Austria, this study; British Columbia, WARD et al., 2004, SEPHTON 2004) into the Lower Rhaetian. The isotopic record then remains constant until the top of the Rhaetian and the significant negative shift of approximately 2.0 to 3.0‰, identified in a number of marine sections in close proximity to the Rhaetian-Hettangian boundary (e.g. KUERSCHNER et al. 2007; GUEX et al., 2004; WARD et al., 2004, VAN de SCHOOTBRUGGE et al 2008). These excursions begin below the highest occurrence of conodonts and Triassic ammonites and the lowest occurrence of Jurassic ammonites, allowing very good correlation, and demonstrating unequivocally that the base of the shift lies below the Triassic-Jurassic boundary. The isotopic trends could be compared to the one at the PT boundary.

The Reingraben event is marked by a disturbance of the carbon cycle. The Lower to Middle Norian crisis is marked by a turning point from slowly increasing carbon isotopic values to gradually decreasing values. The Upper Norian (in the classical sense) is marked by a shift from decreasing to increasing isotope values. From an isotopic point of view, only the Reingraben event (Lower Carnian-Upper Carnian boundary) and the Triassic-Jurassic Boundary can be interpreted as event, whereas other biotic crises of the Late Triassic seem to have occurred during periods of gradual changes in the carbon isotopic composition of the marine seawater.

References

- ATUDOREI, N.-V., 1999, Constraints on the Upper Permian to Upper Triassic marine carbon isotope curve. Case studies from the Tethys. -PhD Thesis: Lausanne University, 155p.
- GAWLICK, H.J., BÖHM, F., 2000, Sequence and isotope stratigraphy of Late Triassic distal periplatform limestones from the Northern Calcareous Alps (Kälberstein Quarry, Berchtesgaden Hallstatt Zone). - *Int. J. Earth Sciences*, **89**, p. 108-129
- GUEX, J., BARTOLINI, A., ATUDOREI, V., AND TAYLOR, D., 2004, High-resolution ammonite and carbon isotope stratigraphy across the Triassic-Jurassic boundary at New York Canyon (Nevada). - *Earth and Planetary Science Letters*, **225**, p. 29-41. 367.
- HALLAM, A., 2002, How catastrophic was the end-Triassic mass extinction. - *Lethaia*, **35**, p. 147-157.
- HAUSER, M., MARTINI, R., BURNS, S., DUMITRICA, P., KRYSSTYN, L., MATTER, A., PETERS, T. & ZANINETTI, L., 2001, Triassic stratigraphic evolution of the Arabian-Greater India embayment of the southern Tethys margin. - *Eclogae Geologicae Helveticae*, **94**(1), p. 29-62.
- HORNUNG, T., BRANDNER, R., 2005, Biochronostratigraphy of the Reingraben Turnover (Hallstatt Facies Belt): Local black shale events controlled by regional tectonics, climatic change and plate tectonics. - *Facies* **51**, p. 460-479.
- HORNUNG, T. BRANDNER, R., KRYSSTYN, L., 2007a, A Tethys-wide mid-Carnian (Upper Triassic) carbonate productivity crisis: Evidence for the Alpine Reingraben Event from Spiti (Indian Himalaya)? - *J. of Asian Earth Sciences*, **30**/ 2, p. 285-302.

- HORNUNG, T. BRANDNER, R., KRYSZYN, L., M. JOACHIMSKI AND L. KEIM 2007c: Multistratigraphic constraints on the NW Tethyan " Carnian Crisis". - *New Mexico Mus. Natural Hist. & Science, Bulletin* **41**, 59-67.
- KORTE, C., KOZUR, H.W., VEIZER, J., 2005, $\delta^{13}C$ and $\delta^{18}O$ values of Triassic brachiopods and carbonate rocks as proxies for coeval seawater and palaeotemperature. - *Palaeogeography, Palaeoclimatology, Palaeoecology*, **226**, p. 287–306.
- KÜRSCHNER, W., BONIS, N. AND KRYSZYN, L., 2007, High resolution carbon isotope stratigraphy and palynostratigraphy of the Triassic–Jurassic transition in the Tiefengraben section – Northern Calcareous Alps (Austria). - *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. **244**, p. 257-280.
- KRYSZYN, L., BOUQUEREL, H., KUERSCHNER, W., S. RICHOSZ & GALLET, Y., (2007): Proposal for a candidate GSSP for the base of the Rhaetian stage. - *New Mexico Museum Nat. Hist. & Science, Bull.* **41**, 189-199.
- McROBERTS, C. A., AND NEWTON, C. R., 1995, Selective extinction among end-Triassic European bivalves. - *Geology*, **23**, no. 2, p. 102-104.
- MORANTE, R., HALLAM, A., 1996, Organic carbon isotopic record across the Triassic–Jurassic boundary in Austria and its bearing on the cause of the mass extinction. – *Geology*, **24**, p. 391-394.
- MIETTO P., ANDREETTA R., BROGLIO LORIGA C., BURATTI N., CIRILLI S., DE ZANCHE V., FURIN S., GIANOLLA P., MANFRIN S., MUTTONI G., NERI C., NICORA A., POSENATO R., PRETO N., RIGO M., ROGHI G. AND SPÖTL C., 2007, A Candidate Of The Global Boundary Stratotype Section And Point For The Base Of The Carnian Stage (Upper Triassic): GSSP at the base of the canadensis Subzone (FAD of Daxatina) in the Prati di Stuares/Stuares Wiesen section (Southern Alps, NE Italy). - *Albertiana*, **36**, p. 78-97.
- MORANTE, R., HALLAM, A., 1996, Organic carbon isotopic record across the Triassic–Jurassic boundary in Austria and its bearing on the cause of the mass extinction. – *Geology*, **24**, p. 391-394.
- MUTTONI, G., KENT, D.V., OLSEN, P., DI STEFANO, P., LOWRIE, W., BERNASCONI, S., MARTIN HERNANDEZ, F., 2004, Tethyan magnetostratigraphy from Pizzo Mondello (Sicily) and correlation to the Late Triassic astrochronological polarity time scale.- *Geol. Soc. Am. Bull.* **116**, p. 1043-1058.
- RICHOSZ, S., KRYSZYN, L. AND SPÖTL, C., 2007a, First detailed carbon isotope curve through the Ladinian-Carnian boundary: The Weissenbach section (Austria). - *Albertiana*, **36**, 98-101.
- RICHOSZ, S., KRYSZYN, L. AND SPÖTL, C., 2007b, Towards a carbon isotope reference curve of the Upper Triassic. - *New Mexico Museum of Natural History and Science, Bulletin* **41**, 366-367.
- SEPTON, M.A., AMOR, K., FRANCHI, I.A., WIGNALL, P.B., NEWTON, R., ZONNEVELD, J.-P., 2002, Carbon and Nitrogen isotope disturbances and an end-Norian extinction event. - *Geology*, **30**, p. 1119–1122.
- van de SCHOOTBRUGGE, B., J. L. PAYNE, A. TOMASOVYCH, J. PROSS, J. FIEBIG, M. BENBRAHIM, K. B. FOLLMI, & T. M. QUAN 2008, Carbon cycle perturbation and stabilization in the wake of the Triassic-Jurassic boundary mass-extinction event. - *Geochem. Geophys. Geosyst.*, **9**, Q04028, doi:10.1029/2007GC001914
- WARD, P.D., GARRISON, G.H., HAGGART, J.W., KRING, D.A., BEATTIE, M.J., 2004, Isotopic evidence bearing on Late Triassic extinction events, Queen Charlotte Islands, British Columbia, and implications for the duration and cause of the Triassic-Jurassic mass extinction. - *Earth & Plan. Sc. Letters* v. **224**, 589-600.
- WILLIFORD, K., WARD, P., GARRISON, G., AND BUICK, R. 2007, An extended stable organic carbon isotope record across the Triassic-Jurassic boundary in the Queen Charlotte Islands, British Columbia, Canada. - *Palaeogeography, Palaeoclimatology, Palaeoecology*, **244**, p. 290-296.